

**(19) World Intellectual Property Organization  
International Bureau**



**(43) International Publication Date**  
**23 May 2002 (23.05.2002)**

**PCT**

**(10) International Publication Number**  
**WO 02/40683 A2**

- (51) **International Patent Classification<sup>7</sup>:** C12N 15/54, 9/12, G01N 33/573, C12Q 1/68, A61K 38/45
- (21) **International Application Number:** PCT/US01/32616
- (22) **International Filing Date:** 22 October 2001 (22.10.2001)
- (25) **Filing Language:** English
- (26) **Publication Language:** English
- (30) **Priority Data:**
- |            |                               |    |
|------------|-------------------------------|----|
| 09/711,134 | 14 November 2000 (14.11.2000) | US |
| 09/858,664 | 17 May 2001 (17.05.2001)      | US |
- (71) **Applicant:** PE CORPORATION (NY) [US/US]; 761 Main Avenue, Norwalk, CT 06859 (US).
- (72) **Inventors:** WEI, Ming-Hui; c/o Celera Genomics, 45 West Gude Drive C2-4#21, Rockville, MD 20850 (US). KETCHUM, Kären; c/o Celera Genomics, 45 West Gude Drive C2-4#21, Rockville, MD 20850 (US). DI FRANCESCO, Valentina; c/o Celera Genomics, 45 West Gude Drive C2-4#21, Rockville, MD 20850 (US).
- BEASLEY, Ellen, M.; c/o Celera Genomics, 45 West Gude Drive C2-4#21, Rockville, MD 20850 (US).
- (74) **Agent:** MILLMAN, Robert, A.; Celera Genomics, Chief Intellectual Property Counsel, 45 West Gude Drive C2-4#21, Rockville, MD 20850 (US).
- (81) **Designated States (national):** AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NO, NZ, PI, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, TZ, UA, UG, UZ, VN, YU, ZA, ZW.
- (84) **Designated States (regional):** ARIPO patent (GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE, TR), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG).

*[Continued on next page]*

- (54) Title:** ISOLATED HUMAN KINASE PROTEINS, NUCLEIC ACID MOLECULES ENCODING HUMAN KINASE PROTEINS, AND USES THEREOF

[illegible][illegible]

**WO 02/40683 A2**

**(57) Abstract:** The present invention provides amino acid sequences of peptides that are encoded by genes within the human genome, the kinase peptides of the present invention. The present invention specifically provides isolated peptide and nucleic acid molecules, methods of identifying orthologs and paralogs of the transporter peptides, and methods of identifying modulators of the kinase peptides.



**Published:**

— without international search report and to be republished upon receipt of that report

*For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.*

# ISOLATED HUMAN KINASE PROTEINS, NUCLEIC ACID MOLECULES ENCODING HUMAN KINASE PROTEINS, AND USES THEREOF

## RELATED APPLICATIONS

5       The present application is a Continuation-In-Part of US Serial No. 09/711,134, filed November 14, 2000 (Atty. Docket CL000927), and U.S. Serial No. 09/858,664 filed May 17, 2001 (Atty. Docket CL000927-CIP).

## FIELD OF THE INVENTION

10       The present invention is in the field of kinase proteins that are related to the myosin light chain kinase subfamily, recombinant DNA molecules, and protein production. The present invention specifically provides novel peptides and proteins that effect protein phosphorylation and nucleic acid molecules encoding such peptide and protein molecules, all of which are useful in the development of human therapeutics and diagnostic compositions and methods.

## BACKGROUND OF THE INVENTION

### 15       Protein Kinases

Kinases regulate many different cell proliferation, differentiation, and signaling processes by adding phosphate groups to proteins. Uncontrolled signaling has been implicated in a variety of disease conditions including inflammation, cancer, arteriosclerosis, and psoriasis. Reversible protein phosphorylation is the main strategy for controlling activities of eukaryotic cells. It is estimated that more than 1000 of the 10,000 proteins active in a typical mammalian cell are phosphorylated. The high energy phosphate, which drives activation, is generally transferred from adenosine triphosphate molecules (ATP) to a particular protein by protein kinases and removed from that protein by protein phosphatases. Phosphorylation occurs in response to extracellular signals (hormones, neurotransmitters, growth and differentiation factors, etc), cell cycle checkpoints, and environmental or nutritional stresses and is roughly analogous to turning on a molecular switch. When the switch goes on, the appropriate protein kinase activates a metabolic enzyme, regulatory protein, receptor, cytoskeletal protein, ion channel or pump, or transcription factor.

25       The kinases comprise the largest known protein group, a superfamily of enzymes with  
30       widely varied functions and specificities. They are usually named after their substrate, their

regulatory molecules, or some aspect of a mutant phenotype. With regard to substrates, the protein kinases may be roughly divided into two groups; those that phosphorylate tyrosine residues (protein tyrosine kinases, PTK) and those that phosphorylate serine or threonine residues (serine/threonine kinases, STK). A few protein kinases have dual specificity and phosphorylate threonine and tyrosine residues. Almost all kinases contain a similar 250-300 amino acid catalytic domain. The N-terminal domain, which contains subdomains I-IV, generally folds into a two-lobed structure, which binds and orients the ATP (or GTP) donor molecule. The larger C terminal lobe, which contains subdomains VI A-XI, binds the protein substrate and carries out the transfer of the gamma phosphate from ATP to the hydroxyl group of a serine, threonine, or tyrosine residue. Subdomain V spans the two lobes.

The kinases may be categorized into families by the different amino acid sequences (generally between 5 and 100 residues) located on either side of, or inserted into loops of, the kinase domain. These added amino acid sequences allow the regulation of each kinase as it recognizes and interacts with its target protein. The primary structure of the kinase domains is conserved and can be further subdivided into 11 subdomains. Each of the 11 subdomains contains specific residues and motifs or patterns of amino acids that are characteristic of that subdomain and are highly conserved (Hardie, G. and Hanks, S. (1995) *The Protein Kinase Facts Books*, Vol I:7-20 Academic Press, San Diego, Calif.).

The second messenger dependent protein kinases primarily mediate the effects of second messengers such as cyclic AMP (cAMP), cyclic GMP, inositol triphosphate, phosphatidylinositol, 3,4,5-triphosphate, cyclic-ADPribose, arachidonic acid, diacylglycerol and calcium-calmodulin. The cyclic-AMP dependent protein kinases (PKA) are important members of the STK family. Cyclic-AMP is an intracellular mediator of hormone action in all prokaryotic and animal cells that have been studied. Such hormone-induced cellular responses include thyroid hormone secretion, cortisol secretion, progesterone secretion, glycogen breakdown, bone resorption, and regulation of heart rate and force of heart muscle contraction. PKA is found in all animal cells and is thought to account for the effects of cyclic-AMP in most of these cells. Altered PKA expression is implicated in a variety of disorders and diseases including cancer, thyroid disorders, diabetes, atherosclerosis, and cardiovascular disease (Isselbacher, K. J. *et al.* (1994) *Harrison's Principles of Internal Medicine*, McGraw-Hill, New York, N.Y., pp. 416-431, 1887).

Calcium-calmodulin (CaM) dependent protein kinases are also members of STK family. Calmodulin is a calcium receptor that mediates many calcium regulated processes by binding to



target proteins in response to the binding of calcium. The principle target protein in these processes is CaM dependent protein kinases. CaM-kinases are involved in regulation of smooth muscle contraction (MLC kinase), glycogen breakdown (phosphorylase kinase), and neurotransmission (CaM kinase I and CaM kinase II). CaM kinase I phosphorylates a variety of substrates including the neurotransmitter related proteins synapsin I and II, the gene transcription regulator, CREB, and the cystic fibrosis conductance regulator protein, CFTR (Haribabu, B. *et al.* (1995) *EMBO Journal* 14:3679-86). CaM II kinase also phosphorylates synapsin at different sites, and controls the synthesis of catecholamines in the brain through phosphorylation and activation of tyrosine hydroxylase. Many of the CaM kinases are activated by phosphorylation in addition to binding to CaM. The kinase may autophosphorylate itself, or be phosphorylated by another kinase as part of a "kinase cascade".

Another ligand-activated protein kinase is 5'-AMP-activated protein kinase (AMPK) (Gao, G. *et al.* (1996) *J. Biol Chem.* 15:8675-81). Mammalian AMPK is a regulator of fatty acid and sterol synthesis through phosphorylation of the enzymes acetyl-CoA carboxylase and hydroxymethylglutaryl-CoA reductase and mediates responses of these pathways to cellular stresses such as heat shock and depletion of glucose and ATP. AMPK is a heterotrimeric complex comprised of a catalytic alpha subunit and two non-catalytic beta and gamma subunits that are believed to regulate the activity of the alpha subunit. Subunits of AMPK have a much wider distribution in non-lipogenic tissues such as brain, heart, spleen, and lung than expected. This distribution suggests that its role may extend beyond regulation of lipid metabolism alone.

The mitogen-activated protein kinases (MAP) are also members of the STK family. MAP kinases also regulate intracellular signaling pathways. They mediate signal transduction from the cell surface to the nucleus via phosphorylation cascades. Several subgroups have been identified, and each manifests different substrate specificities and responds to distinct extracellular stimuli (Egan, S. E. and Weinberg, R. A. (1993) *Nature* 365:781-783). MAP kinase signaling pathways are present in mammalian cells as well as in yeast. The extracellular stimuli that activate mammalian pathways include epidermal growth factor (EGF), ultraviolet light, hyperosmolar medium, heat shock, endotoxic lipopolysaccharide (LPS), and pro-inflammatory cytokines such as tumor necrosis factor (TNF) and interleukin-1 (IL-1).

PRK (proliferation-related kinase) is a serum/cytokine inducible STK that is involved in regulation of the cell cycle and cell proliferation in human megakaryotic cells (Li, B. *et al.* (1996) *J. Biol. Chem.* 271:19402-8). PRK is related to the polo (derived from humans polo gene) family of STKs implicated in cell division. PRK is downregulated in lung tumor tissue and may

be a proto-oncogene whose deregulated expression in normal tissue leads to oncogenic transformation. Altered MAP kinase expression is implicated in a variety of disease conditions including cancer, inflammation, immune disorders, and disorders affecting growth and development.

5        The cyclin-dependent protein kinases (CDKs) are another group of STKs that control the progression of cells through the cell cycle. Cyclins are small regulatory proteins that act by binding to and activating CDKs that then trigger various phases of the cell cycle by phosphorylating and activating selected proteins involved in the mitotic process. CDKs are unique in that they require multiple inputs to become activated. In addition to the binding of  
10       cyclin, CDK activation requires the phosphorylation of a specific threonine residue and the dephosphorylation of a specific tyrosine residue.

      Protein tyrosine kinases, PTKs, specifically phosphorylate tyrosine residues on their target proteins and may be divided into transmembrane, receptor PTKs and nontransmembrane, non-receptor PTKs. Transmembrane protein-tyrosine kinases are receptors for most growth  
15       factors. Binding of growth factor to the receptor activates the transfer of a phosphate group from ATP to selected tyrosine side chains of the receptor and other specific proteins. Growth factors (GF) associated with receptor PTKs include; epidermal GF, platelet-derived GF, fibroblast GF, hepatocyte GF, insulin and insulin-like GFs, nerve GF, vascular endothelial GF, and macrophage colony stimulating factor.

20       Non-receptor PTKs lack transmembrane regions and, instead, form complexes with the intracellular regions of cell surface receptors. Such receptors that function through non-receptor PTKs include those for cytokines, hormones (growth hormone and prolactin) and antigen-specific receptors on T and B lymphocytes.

      Many of these PTKs were first identified as the products of mutant oncogenes in cancer  
25       cells where their activation was no longer subject to normal cellular controls. In fact, about one third of the known oncogenes encode PTKs, and it is well known that cellular transformation (oncogenesis) is often accompanied by increased tyrosine phosphorylation activity (Carbonneau H and Tonks NK (1992) *Annu. Rev. Cell. Biol.* 8:463-93). Regulation of PTK activity may therefore be an important strategy in controlling some types of cancer.

### 30       Myosin Light Chain Kinase

      Activation of smooth/nonmuscle myosin light chain kinase (MLCK) by Ca/calmodulin results in phosphorylation of myosin regulatory light chain that plays important roles in initiation of smooth muscle contraction, endothelial cell retraction, secretion, and other cellular processes

(Stull et al., in *International Symposium on Regulation of the Contractile Cycle in Smooth Muscle*, April 26, 1995, Mie, Japan). The same myosin light chain kinases are present in smooth and nonmuscle tissues.(Gallagher et al., *J Biol Chem* 1991 Dec 15;266(35):23936-44, Published erratum appears in *J Biol Chem* 1992 May 5;267(13):9450). The phosphorylation of myosin light chains by myosin light chain kinase is a key event in agonist-mediated endothelial cell gap formation and vascular permeability. Amino acid sequence analysis indicates endothelial MLCK consensus sequences for a variety of protein kinases including highly conserved potential phosphorylation sites for cAMP-dependent protein kinase A (PKA) in the CaM-binding region. Augmentation of intracellular cAMP levels markedly enhanced MLCK phosphorylation (2.5- fold increase) and reduced kinase activity in MLCK immunoprecipitates (4-fold decreases) (Garcia et al., *Am J Respir Cell Mol Biol* 1997 May;16(5):489-94). The smooth/nonmuscle myosin light chain kinase contains a catalytic core homologous to that of other protein kinases and a carboxyl-terminal regulatory domain consisting of both an inhibitory sequence and a calmodulin-binding sequence (Kemp et al., *Trends Biochem. Sci.* 19, 440-444, 1994; Stull et al., 1995). Initially, inspection of the linear sequence within the regulatory domain revealed a similar number and sequential arrangement of 4 basic residues with those shown to be important substrate determinants in a synthetic peptide containing residues 11-23 of the myosin regulatory light chain . Thus, it has been proposed that the regulatory domain contained a pseudosubstrate inhibitory sequence whereby 4 specific basic residues in myosin light chain kinase mimic the basic substrate determinants in the light chain peptide substrate . Binding of the pseudosubstrate sequence to the active site inhibited activity. Intrasteric inhibition involves an autoinhibitory sequence that folds back on the catalytic site to inhibit kinase activity as opposed to an allosteric mechanism whereby a conformational change induced at a site distinct from the active site would be responsible for regulation of enzyme activity (Kemp et al., *Biochim. Biophys. Acta.* 1094, 67-76, 1991). The sequence comprising the pseudosubstrate region was later expanded to include overlap with the complete amino terminus of the light chain (Faux et al., *Mol. Cell. Biochem.* 128, 81-91, 1993). However, these additional residues(1, 2, 3, 4, 5, 6, 7, 8, 9, 10) are not important for substrate binding and thus are not part of the consensus phosphorylation sequence (Kemp et al., *Trends Biochem. Sci.* 15, 342-346, 1990).

Kinase proteins, particularly members of the myosin light chain kinase subfamily, are a major target for drug action and development. Accordingly, it is valuable to the field of pharmaceutical development to identify and characterize previously unknown members of this subfamily of kinase proteins. The present invention advances the state of the art by providing

previously unidentified human kinase proteins that have homology to members of the myosin light chain kinase subfamily.

### **SUMMARY OF THE INVENTION**

5 The present invention is based in part on the identification of amino acid sequences of human kinase peptides and proteins that are related to the myosin light chain kinase subfamily, as well as allelic variants and other mammalian orthologs thereof. These unique peptide sequences, and nucleic acid sequences that encode these peptides, can be used as models for the development of human therapeutic targets, aid in the identification of therapeutic proteins, and serve as targets for the development of human therapeutic agents that modulate kinase activity in  
10 cells and tissues that express the kinase. Experimental data as provided in Figure 1 indicates expression in the human placenta, kidney, lung, skeletal muscle, heart, fetal brain, and colon carcinoma.

### **DESCRIPTION OF THE FIGURE SHEETS**

FIGURE 1 provides the nucleotide sequence of a cDNA molecule or transcript sequence  
15 that encodes the kinase protein of the present invention. (SEQ ID NO:1) In addition, structure and functional information is provided, such as ATG start, stop and tissue distribution, where available, that allows one to readily determine specific uses of inventions based on this molecular sequence. Experimental data as provided in Figure 1 indicates expression in the human placenta, kidney, lung, skeletal muscle, heart, fetal brain, and colon carcinoma.

20 FIGURE 2 provides the predicted amino acid sequence of the kinase of the present invention. (SEQ ID NO:2) In addition structure and functional information such as protein family, function, and modification sites is provided where available, allowing one to readily determine specific uses of inventions based on this molecular sequence.

FIGURE 3 provides genomic sequences that span the gene encoding the kinase protein of  
25 the present invention. (SEQ ID NO:3) In addition structure and functional information, such as intron/exon structure, promoter location, etc., is provided where available, allowing one to readily determine specific uses of inventions based on this molecular sequence. 6 SNPs, have been identified in the gene encoding the kinase protein provided by the present invention and are given in Figure 3.

## DETAILED DESCRIPTION OF THE INVENTION

### General Description

The present invention is based on the sequencing of the human genome. During the sequencing and assembly of the human genome, analysis of the sequence information revealed previously unidentified fragments of the human genome that encode peptides that share structural and/or sequence homology to protein/peptide/domains identified and characterized within the art as being a kinase protein or part of a kinase protein and are related to the myosin light chain kinase subfamily. Utilizing these sequences, additional genomic sequences were assembled and transcript and/or cDNA sequences were isolated and characterized. Based on this analysis, the present invention provides amino acid sequences of human kinase peptides and proteins that are related to the myosin light chain kinase subfamily, nucleic acid sequences in the form of transcript sequences, cDNA sequences and/or genomic sequences that encode these kinase peptides and proteins, nucleic acid variation (allelic information), tissue distribution of expression, and information about the closest art known protein/peptide/domain that has structural or sequence homology to the kinase of the present invention.

In addition to being previously unknown, the peptides that are provided in the present invention are selected based on their ability to be used for the development of commercially important products and services. Specifically, the present peptides are selected based on homology and/or structural relatedness to known kinase proteins of the myosin light chain kinase subfamily and the expression pattern observed. Experimental data as provided in Figure 1 indicates expression in the human placenta, kidney, lung, skeletal muscle, heart, fetal brain, and colon carcinoma. The art has clearly established the commercial importance of members of this family of proteins and proteins that have expression patterns similar to that of the present gene. Some of the more specific features of the peptides of the present invention, and the uses thereof, are described herein, particularly in the Background of the Invention and in the annotation provided in the Figures, and/or are known within the art for each of the known myosin light chain kinase family or subfamily of kinase proteins.

### Specific Embodiments

#### Peptide Molecules

The present invention provides nucleic acid sequences that encode protein molecules that have been identified as being members of the kinase family of proteins and are related to the

myosin light chain kinase subfamily (protein sequences are provided in Figure 2, transcript/cDNA sequences are provided in Figure 1 and genomic sequences are provided in Figure 3). The peptide sequences provided in Figure 2, as well as the obvious variants described herein, particularly allelic variants as identified herein and using the information in Figure 3, will be referred herein as the kinase peptides of the present invention, kinase peptides, or peptides/proteins of the present invention.

The present invention provides isolated peptide and protein molecules that consist of, consist essentially of, or comprise the amino acid sequences of the kinase peptides disclosed in the Figure 2, (encoded by the nucleic acid molecule shown in Figure 1, transcript/cDNA or Figure 3, genomic sequence), as well as all obvious variants of these peptides that are within the art to make and use. Some of these variants are described in detail below.

As used herein, a peptide is said to be "isolated" or "purified" when it is substantially free of cellular material or free of chemical precursors or other chemicals. The peptides of the present invention can be purified to homogeneity or other degrees of purity. The level of purification will be based on the intended use. The critical feature is that the preparation allows for the desired function of the peptide, even if in the presence of considerable amounts of other components (the features of an isolated nucleic acid molecule is discussed below).

In some uses, "substantially free of cellular material" includes preparations of the peptide having less than about 30% (by dry weight) other proteins (i.e., contaminating protein), less than about 20% other proteins, less than about 10% other proteins, or less than about 5% other proteins. When the peptide is recombinantly produced, it can also be substantially free of culture medium, i.e., culture medium represents less than about 20% of the volume of the protein preparation.

The language "substantially free of chemical precursors or other chemicals" includes preparations of the peptide in which it is separated from chemical precursors or other chemicals that are involved in its synthesis. In one embodiment, the language "substantially free of chemical precursors or other chemicals" includes preparations of the kinase peptide having less than about 30% (by dry weight) chemical precursors or other chemicals, less than about 20% chemical precursors or other chemicals, less than about 10% chemical precursors or other chemicals, or less than about 5% chemical precursors or other chemicals.

The isolated kinase peptide can be purified from cells that naturally express it, purified from cells that have been altered to express it (recombinant), or synthesized using known protein synthesis methods. Experimental data as provided in Figure 1 indicates expression in the human placenta, kidney, lung, skeletal muscle, heart, fetal brain, and colon carcinoma. For example, a

nucleic acid molecule encoding the kinase peptide is cloned into an expression vector, the expression vector introduced into a host cell and the protein expressed in the host cell. The protein can then be isolated from the cells by an appropriate purification scheme using standard protein purification techniques. Many of these techniques are described in detail below.

5        Accordingly, the present invention provides proteins that consist of the amino acid sequences provided in Figure 2 (SEQ ID NO:2), for example, proteins encoded by the transcript/cDNA nucleic acid sequences shown in Figure 1 (SEQ ID NO:1) and the genomic sequences provided in Figure 3 (SEQ ID NO:3). The amino acid sequence of such a protein is provided in Figure 2. A protein consists of an amino acid sequence when the amino acid sequence  
10        is the final amino acid sequence of the protein.

      The present invention further provides proteins that consist essentially of the amino acid sequences provided in Figure 2 (SEQ ID NO:2), for example, proteins encoded by the transcript/cDNA nucleic acid sequences shown in Figure 1 (SEQ ID NO:1) and the genomic sequences provided in Figure 3 (SEQ ID NO:3). A protein consists essentially of an amino acid  
15        sequence when such an amino acid sequence is present with only a few additional amino acid residues, for example from about 1 to about 100 or so additional residues, typically from 1 to about 20 additional residues in the final protein.

      The present invention further provides proteins that comprise the amino acid sequences provided in Figure 2 (SEQ ID NO:2), for example, proteins encoded by the transcript/cDNA nucleic acid sequences shown in Figure 1 (SEQ ID NO:1) and the genomic sequences provided in Figure 3  
20        (SEQ ID NO:3). A protein comprises an amino acid sequence when the amino acid sequence is at least part of the final amino acid sequence of the protein. In such a fashion, the protein can be only the peptide or have additional amino acid molecules, such as amino acid residues (contiguous encoded sequence) that are naturally associated with it or heterologous amino acid residues/peptide  
25        sequences. Such a protein can have a few additional amino acid residues or can comprise several hundred or more additional amino acids. The preferred classes of proteins that are comprised of the kinase peptides of the present invention are the naturally occurring mature proteins. A brief description of how various types of these proteins can be made/isolated is provided below.

      The kinase peptides of the present invention can be attached to heterologous sequences to  
30        form chimeric or fusion proteins. Such chimeric and fusion proteins comprise a kinase peptide operatively linked to a heterologous protein having an amino acid sequence not substantially homologous to the kinase peptide. "Operatively linked" indicates that the kinase peptide and the

heterologous protein are fused in-frame. The heterologous protein can be fused to the N-terminus or C-terminus of the kinase peptide.

In some uses, the fusion protein does not affect the activity of the kinase peptide *per se*. For example, the fusion protein can include, but is not limited to, enzymatic fusion proteins, for example beta-galactosidase fusions, yeast two-hybrid GAL fusions, poly-His fusions, MYC-tagged, HI-tagged and Ig fusions. Such fusion proteins, particularly poly-His fusions, can facilitate the purification of recombinant kinase peptide. In certain host cells (e.g., mammalian host cells), expression and/or secretion of a protein can be increased by using a heterologous signal sequence.

A chimeric or fusion protein can be produced by standard recombinant DNA techniques. For example, DNA fragments coding for the different protein sequences are ligated together in-frame in accordance with conventional techniques. In another embodiment, the fusion gene can be synthesized by conventional techniques including automated DNA synthesizers. Alternatively, PCR amplification of gene fragments can be carried out using anchor primers which give rise to complementary overhangs between two consecutive gene fragments which can subsequently be annealed and re-amplified to generate a chimeric gene sequence (see Ausubel *et al.*, *Current Protocols in Molecular Biology*, 1992). Moreover, many expression vectors are commercially available that already encode a fusion moiety (e.g., a GST protein). A kinase peptide-encoding nucleic acid can be cloned into such an expression vector such that the fusion moiety is linked in-frame to the kinase peptide.

As mentioned above, the present invention also provides and enables obvious variants of the amino acid sequence of the proteins of the present invention, such as naturally occurring mature forms of the peptide, allelic/sequence variants of the peptides, non-naturally occurring recombinantly derived variants of the peptides, and orthologs and paralogs of the peptides. Such variants can readily be generated using art-known techniques in the fields of recombinant nucleic acid technology and protein biochemistry. It is understood, however, that variants exclude any amino acid sequences disclosed prior to the invention.

Such variants can readily be identified/made using molecular techniques and the sequence information disclosed herein. Further, such variants can readily be distinguished from other peptides based on sequence and/or structural homology to the kinase peptides of the present invention. The degree of homology/identity present will be based primarily on whether the peptide is a functional variant or non-functional variant, the amount of divergence present in the paralog family and the evolutionary distance between the orthologs.



To determine the percent identity of two amino acid sequences or two nucleic acid sequences, the sequences are aligned for optimal comparison purposes (e.g., gaps can be introduced in one or both of a first and a second amino acid or nucleic acid sequence for optimal alignment and non-homologous sequences can be disregarded for comparison purposes). In a preferred embodiment, at least 30%, 40%, 50%, 60%, 70%, 80%, or 90% or more of the length of a reference sequence is aligned for comparison purposes. The amino acid residues or nucleotides at corresponding amino acid positions or nucleotide positions are then compared. When a position in the first sequence is occupied by the same amino acid residue or nucleotide as the corresponding position in the second sequence, then the molecules are identical at that position (as used herein amino acid or nucleic acid "identity" is equivalent to amino acid or nucleic acid "homology"). The percent identity between the two sequences is a function of the number of identical positions shared by the sequences, taking into account the number of gaps, and the length of each gap, which need to be introduced for optimal alignment of the two sequences.

The comparison of sequences and determination of percent identity and similarity between two sequences can be accomplished using a mathematical algorithm. (*Computational Molecular Biology*, Lesk, A.M., ed., Oxford University Press, New York, 1988; *Biocomputing: Informatics and Genome Projects*, Smith, D.W., ed., Academic Press, New York, 1993; *Computer Analysis of Sequence Data, Part 1*, Griffin, A.M., and Griffin, H.G., eds., Humana Press, New Jersey, 1994; *Sequence Analysis in Molecular Biology*, von Heinje, G., Academic Press, 1987; and *Sequence Analysis Primer*, Gribskov, M. and Devereux, J., eds., M Stockton Press, New York, 1991). In a preferred embodiment, the percent identity between two amino acid sequences is determined using the Needleman and Wunsch (*J. Mol. Biol.* (48):444-453 (1970)) algorithm which has been incorporated into the GAP program in the GCG software package (available at <http://www.gcg.com>), using either a Blossom 62 matrix or a PAM250 matrix, and a gap weight of 16, 14, 12, 10, 8, 6, or 4 and a length weight of 1, 2, 3, 4, 5, or 6. In yet another preferred embodiment, the percent identity between two nucleotide sequences is determined using the GAP program in the GCG software package (Devereux, J., *et al.*, *Nucleic Acids Res.* 12(1):387 (1984)) (available at <http://www.gcg.com>), using a NWSgapdna.CMP matrix and a gap weight of 40, 50, 60, 70, or 80 and a length weight of 1, 2, 3, 4, 5, or 6. In another embodiment, the percent identity between two amino acid or nucleotide sequences is determined using the algorithm of E. Myers and W. Miller (CABIOS, 4:11-17 (1989)) which has been incorporated

into the ALIGN program (version 2.0), using a PAM120 weight residue table, a gap length penalty of 12 and a gap penalty of 4.

The nucleic acid and protein sequences of the present invention can further be used as a "query sequence" to perform a search against sequence databases to, for example, identify other family members or related sequences. Such searches can be performed using the NBLAST and XBLAST programs (version 2.0) of Altschul, *et al.* (*J. Mol. Biol.* 215:403-10 (1990)). BLAST nucleotide searches can be performed with the NBLAST program, score = 100, wordlength = 12 to obtain nucleotide sequences homologous to the nucleic acid molecules of the invention. BLAST protein searches can be performed with the XBLAST program, score = 50, wordlength = 3 to obtain amino acid sequences homologous to the proteins of the invention. To obtain gapped alignments for comparison purposes, Gapped BLAST can be utilized as described in Altschul *et al.* (*Nucleic Acids Res.* 25(17):3389-3402 (1997)). When utilizing BLAST and gapped BLAST programs, the default parameters of the respective programs (e.g., XBLAST and NBLAST) can be used.

Full-length pre-processed forms, as well as mature processed forms, of proteins that comprise one of the peptides of the present invention can readily be identified as having complete sequence identity to one of the kinase peptides of the present invention as well as being encoded by the same genetic locus as the kinase peptide provided herein. As indicated by the data presented in Figure 3, the map position was determined to be on chromosome 1 by ePCR, and confirmed with radiation hybrid mapping. As indicated by the data presented in Figure 3, the gene provided by the present invention encoding a novel phosphatase maps to public BAC AC AC023889, which is known to be located on human chromosome 1.

Allelic variants of a kinase peptide can readily be identified as being a human protein having a high degree (significant) of sequence homology/identity to at least a portion of the kinase peptide as well as being encoded by the same genetic locus as the kinase peptide provided herein. Genetic locus can readily be determined based on the genomic information provided in Figure 3, such as the genomic sequence mapped to the reference human. As indicated by the data presented in Figure 3, the map position was determined to be on chromosome 1 by ePCR, and confirmed with radiation hybrid mapping. As indicated by the data presented in Figure 3, the gene provided by the present invention encoding a novel phosphatase maps to public BAC AC AC023889, which is known to be located on human chromosome 1. As used herein, two proteins (or a region of the proteins) have significant homology when the amino acid sequences are typically at least about 70-80%, 80-90%, and more typically at least about 90-95% or more homologous. A significantly

homologous amino acid sequence, according to the present invention, will be encoded by a nucleic acid sequence that will hybridize to a kinase peptide encoding nucleic acid molecule under stringent conditions as more fully described below.

Figure 3 provides information on SNPs that have been identified in a gene encoding the kinase protein of the present invention. 6 SNP variants were found, and all SNPs in exons, of which 3 of these cause changes in the amino acid sequence (i.e., nonsynonymous SNPs). The changes in the amino acid sequence that these SNPs cause is indicated in Figure 3 and can readily be determined using the universal genetic code and the protein sequence provided in Figure 2 as a reference.

Paralogs of a kinase peptide can readily be identified as having some degree of significant sequence homology/identity to at least a portion of the kinase peptide, as being encoded by a gene from humans, and as having similar activity or function. Two proteins will typically be considered paralogs when the amino acid sequences are typically at least about 60% or greater, and more typically at least about 70% or greater homology through a given region or domain. Such paralogs will be encoded by a nucleic acid sequence that will hybridize to a kinase peptide encoding nucleic acid molecule under moderate to stringent conditions as more fully described below.

Orthologs of a kinase peptide can readily be identified as having some degree of significant sequence homology/identity to at least a portion of the kinase peptide as well as being encoded by a gene from another organism. Preferred orthologs will be isolated from mammals, preferably primates, for the development of human therapeutic targets and agents. Such orthologs will be encoded by a nucleic acid sequence that will hybridize to a kinase peptide encoding nucleic acid molecule under moderate to stringent conditions, as more fully described below, depending on the degree of relatedness of the two organisms yielding the proteins.

Non-naturally occurring variants of the kinase peptides of the present invention can readily be generated using recombinant techniques. Such variants include, but are not limited to deletions, additions and substitutions in the amino acid sequence of the kinase peptide. For example, one class of substitutions are conserved amino acid substitution. Such substitutions are those that substitute a given amino acid in a kinase peptide by another amino acid of like characteristics. Typically seen as conservative substitutions are the replacements, one for another, among the aliphatic amino acids Ala, Val, Leu, and Ile; interchange of the hydroxyl residues Ser and Thr; exchange of the acidic residues Asp and Glu; substitution between the amide residues Asn and Gln; exchange of the basic residues Lys and Arg; and replacements among the aromatic residues Phe and Tyr. Guidance

concerning which amino acid changes are likely to be phenotypically silent are found in Bowie *et al.*, *Science* 247:1306-1310 (1990).

Variant kinase peptides can be fully functional or can lack function in one or more activities, e.g. ability to bind substrate, ability to phosphorylate substrate, ability to mediate signaling, etc.

5 Fully functional variants typically contain only conservative variation or variation in non-critical residues or in non-critical regions. Figure 2 provides the result of protein analysis and can be used to identify critical domains/regions. Functional variants can also contain substitution of similar amino acids that result in no change or an insignificant change in function. Alternatively, such substitutions may positively or negatively affect function to some degree.

10 Non-functional variants typically contain one or more non-conservative amino acid substitutions, deletions, insertions, inversions, or truncation or a substitution, insertion, inversion, or deletion in a critical residue or critical region.

Amino acids that are essential for function can be identified by methods known in the art, such as site-directed mutagenesis or alanine-scanning mutagenesis (Cunningham *et al.*, *Science* 15 244:1081-1085 (1989)), particularly using the results provided in Figure 2. The latter procedure introduces single alanine mutations at every residue in the molecule. The resulting mutant molecules are then tested for biological activity such as kinase activity or in assays such as an *in vitro* proliferative activity. Sites that are critical for binding partner/substrate binding can also be determined by structural analysis such as crystallization, nuclear magnetic resonance or 20 photoaffinity labeling (Smith *et al.*, *J. Mol. Biol.* 224:899-904 (1992); de Vos *et al.* *Science* 255:306-312 (1992)).

The present invention further provides fragments of the kinase peptides, in addition to proteins and peptides that comprise and consist of such fragments, particularly those comprising the residues identified in Figure 2. The fragments to which the invention pertains, however, are not to 25 be construed as encompassing fragments that may be disclosed publicly prior to the present invention.

As used herein, a fragment comprises at least 8, 10, 12, 14, 16, or more contiguous amino acid residues from a kinase peptide. Such fragments can be chosen based on the ability to retain one or more of the biological activities of the kinase peptide or could be chosen for the ability to 30 perform a function, e.g. bind a substrate or act as an immunogen. Particularly important fragments are biologically active fragments, peptides that are, for example, about 8 or more amino acids in length. Such fragments will typically comprise a domain or motif of the kinase peptide, e.g., active site, a transmembrane domain or a substrate-binding domain. Further, possible fragments include,

but are not limited to, domain or motif containing fragments, soluble peptide fragments, and fragments containing immunogenic structures. Predicted domains and functional sites are readily identifiable by computer programs well known and readily available to those of skill in the art (e.g., PROSITE analysis). The results of one such analysis are provided in Figure 2.

5 Polypeptides often contain amino acids other than the 20 amino acids commonly referred to as the 20 naturally occurring amino acids. Further, many amino acids, including the terminal amino acids, may be modified by natural processes, such as processing and other post-translational modifications, or by chemical modification techniques well known in the art. Common modifications that occur naturally in kinase peptides are described in basic texts, detailed  
10 monographs, and the research literature, and they are well known to those of skill in the art (some of these features are identified in Figure 2).

Known modifications include, but are not limited to, acetylation, acylation, ADP-ribosylation, amidation, covalent attachment of flavin, covalent attachment of a heme moiety, covalent attachment of a nucleotide or nucleotide derivative, covalent attachment of a lipid or lipid  
15 derivative, covalent attachment of phosphatidylinositol, cross-linking, cyclization, disulfide bond formation, demethylation, formation of covalent crosslinks, formation of cystine, formation of pyroglutamate, formylation, gamma carboxylation, glycosylation, GPI anchor formation, hydroxylation, iodination, methylation, myristoylation, oxidation, proteolytic processing, phosphorylation, prenylation, racemization, selenoylation, sulfation, transfer-RNA mediated  
20 addition of amino acids to proteins such as arginylation, and ubiquitination.

Such modifications are well known to those of skill in the art and have been described in great detail in the scientific literature. Several particularly common modifications, glycosylation, lipid attachment, sulfation, gamma-carboxylation of glutamic acid residues, hydroxylation and ADP-ribosylation, for instance, are described in most basic texts, such as *Proteins - Structure and*  
25 *Molecular Properties*, 2nd Ed., T.E. Creighton, W. H. Freeman and Company, New York (1993). Many detailed reviews are available on this subject, such as by Wold, F., *Posttranslational Covalent Modification of Proteins*, B.C. Johnson, Ed., Academic Press, New York 1-12 (1983); Seifter *et al.* (*Meth. Enzymol.* 182: 626-646 (1990)) and Rattan *et al.* (*Ann. N.Y. Acad. Sci.* 663:48-62 (1992)).

Accordingly, the kinase peptides of the present invention also encompass derivatives or  
30 analogs in which a substituted amino acid residue is not one encoded by the genetic code, in which a substituent group is included, in which the mature kinase peptide is fused with another compound, such as a compound to increase the half-life of the kinase peptide (for example, polyethylene glycol), or in which the additional amino acids are fused to the mature kinase peptide, such as a

leader or secretory sequence or a sequence for purification of the mature kinase peptide or a pro-protein sequence.

#### Protein/Peptide Uses

5       The proteins of the present invention can be used in substantial and specific assays related to the functional information provided in the Figures; to raise antibodies or to elicit another immune response; as a reagent (including the labeled reagent) in assays designed to quantitatively determine levels of the protein (or its binding partner or ligand) in biological fluids; and as markers for tissues in which the corresponding protein is preferentially expressed  
10 (either constitutively or at a particular stage of tissue differentiation or development or in a disease state). Where the protein binds or potentially binds to another protein or ligand (such as, for example, in a kinase-effector protein interaction or kinase-ligand interaction), the protein can be used to identify the binding partner/ligand so as to develop a system to identify inhibitors of the binding interaction. Any or all of these uses are capable of being developed into reagent  
15 grade or kit format for commercialization as commercial products.

Substantial chemical and structural homology exists between the kinase protein of the present invention described herein and myosin light chain kinase (see Figure 1). As discussed in the background, myosin light chain kinase are known in the art to be involved in smooth muscle contraction, endothelial cell retraction, secretion, and other cellular process. Accordingly, the  
20 myosin light chain kinase, and the encoding gene, provided by the present invention is useful for treating, preventing, and/or diagnosing disorders associated with muscle, endothelial cells.

Methods for performing the uses listed above are well known to those skilled in the art. References disclosing such methods include "Molecular Cloning: A Laboratory Manual", 2d ed., Cold Spring Harbor Laboratory Press, Sambrook, J., E. F. Fritsch and T. Maniatis eds., 1989,  
25 and "Methods in Enzymology: Guide to Molecular Cloning Techniques", Academic Press, Berger, S. L. and A. R. Kimmel eds., 1987.

The potential uses of the peptides of the present invention are based primarily on the source of the protein as well as the class/action of the protein. For example, kinases isolated from humans and their human/mammalian orthologs serve as targets for identifying agents for  
30 use in mammalian therapeutic applications, e.g. a human drug, particularly in modulating a biological or pathological response in a cell or tissue that expresses the kinase. Experimental data as provided in Figure 1 indicates that kinase proteins of the present invention are expressed in the human placenta, kidney, lung, skeletal muscle, heart, fetal brain, and colon carcinoma.

Specifically, a virtual northern blot shows expression in human colon carcinoma. In addition, PCR-based tissue screening panel indicates expression in human placenta, kidney, lung, skeletal muscle, heart, and fetal brain. A large percentage of pharmaceutical agents are being developed that modulate the activity of kinase proteins, particularly members of the myosin light chain  
5 kinase subfamily (see Background of the Invention). The structural and functional information provided in the Background and Figures provide specific and substantial uses for the molecules of the present invention, particularly in combination with the expression information provided in Figure 1. Experimental data as provided in Figure 1 indicates expression in the human placenta, kidney, lung, skeletal muscle, heart, fetal brain, and colon carcinoma. Such uses can readily be  
10 determined using the information provided herein, that which is known in the art, and routine experimentation.

The proteins of the present invention (including variants and fragments that may have been disclosed prior to the present invention) are useful for biological assays related to kinases that are related to members of the myosin light chain kinase subfamily. Such assays involve any of the  
15 known kinase functions or activities or properties useful for diagnosis and treatment of kinase-related conditions that are specific for the subfamily of kinases that the one of the present invention belongs to, particularly in cells and tissues that express the kinase. Experimental data as provided in Figure 1 indicates that kinase proteins of the present invention are expressed in the human placenta, kidney, lung, skeletal muscle, heart, fetal brain, and colon carcinoma. Specifically, a virtual northern  
20 blot shows expression in human colon carcinoma. In addition, PCR-based tissue screening panel indicates expression in human placenta, kidney, lung, skeletal muscle, heart, and fetal brain.

The proteins of the present invention are also useful in drug screening assays, in cell-based or cell-free systems. Cell-based systems can be native, i.e., cells that normally express the kinase, as a biopsy or expanded in cell culture. Experimental data as provided in Figure 1 indicates  
25 expression in the human placenta, kidney, lung, skeletal muscle, heart, fetal brain, and colon carcinoma. In an alternate embodiment, cell-based assays involve recombinant host cells expressing the kinase protein.

The polypeptides can be used to identify compounds that modulate kinase activity of the protein in its natural state or an altered form that causes a specific disease or pathology associated  
30 with the kinase. Both the kinases of the present invention and appropriate variants and fragments can be used in high-throughput screens to assay candidate compounds for the ability to bind to the kinase. These compounds can be further screened against a functional kinase to determine the effect of the compound on the kinase activity. Further, these compounds can be tested in animal or

invertebrate systems to determine activity/effectiveness. Compounds can be identified that activate (agonist) or inactivate (antagonist) the kinase to a desired degree.

Further, the proteins of the present invention can be used to screen a compound for the ability to stimulate or inhibit interaction between the kinase protein and a molecule that normally interacts with the kinase protein, e.g. a substrate or a component of the signal pathway that the kinase protein normally interacts (for example, another kinase). Such assays typically include the steps of combining the kinase protein with a candidate compound under conditions that allow the kinase protein, or fragment, to interact with the target molecule, and to detect the formation of a complex between the protein and the target or to detect the biochemical consequence of the interaction with the kinase protein and the target, such as any of the associated effects of signal transduction such as protein phosphorylation, cAMP turnover, and adenylate cyclase activation, etc.

Candidate compounds include, for example, 1) peptides such as soluble peptides, including Ig-tailed fusion peptides and members of random peptide libraries (see, e.g., Lam *et al.*, *Nature* 354:82-84 (1991); Houghten *et al.*, *Nature* 354:84-86 (1991)) and combinatorial chemistry-derived molecular libraries made of D- and/or L- configuration amino acids; 2) phosphopeptides (e.g., members of random and partially degenerate, directed phosphopeptide libraries, see, e.g., Songyang *et al.*, *Cell* 72:767-778 (1993)); 3) antibodies (e.g., polyclonal, monoclonal, humanized, anti-idiotypic, chimeric, and single chain antibodies as well as Fab, F(ab')<sub>2</sub>, Fab expression library fragments, and epitope-binding fragments of antibodies); and 4) small organic and inorganic molecules (e.g., molecules obtained from combinatorial and natural product libraries).

One candidate compound is a soluble fragment of the receptor that competes for substrate binding. Other candidate compounds include mutant kinases or appropriate fragments containing mutations that affect kinase function and thus compete for substrate. Accordingly, a fragment that competes for substrate, for example with a higher affinity, or a fragment that binds substrate but does not allow release, is encompassed by the invention.

The invention further includes other end point assays to identify compounds that modulate (stimulate or inhibit) kinase activity. The assays typically involve an assay of events in the signal transduction pathway that indicate kinase activity. Thus, the phosphorylation of a substrate, activation of a protein, a change in the expression of genes that are up- or down-regulated in response to the kinase protein dependent signal cascade can be assayed.

Any of the biological or biochemical functions mediated by the kinase can be used as an endpoint assay. These include all of the biochemical or biochemical/biological events described herein, in the references cited herein, incorporated by reference for these endpoint assay targets, and



other functions known to those of ordinary skill in the art or that can be readily identified using the information provided in the Figures, particularly Figure 2. Specifically, a biological function of a cell or tissues that expresses the kinase can be assayed. Experimental data as provided in Figure 1 indicates that kinase proteins of the present invention are expressed in the human placenta, kidney, lung, skeletal muscle, heart, fetal brain, and colon carcinoma. Specifically, a virtual northern blot shows expression in human colon carcinoma. In addition, PCR-based tissue screening panel indicates expression in human placenta, kidney, lung, skeletal muscle, heart, and fetal brain.

Binding and/or activating compounds can also be screened by using chimeric kinase proteins in which the amino terminal extracellular domain, or parts thereof, the entire transmembrane domain or subregions, such as any of the seven transmembrane segments or any of the intracellular or extracellular loops and the carboxy terminal intracellular domain, or parts thereof, can be replaced by heterologous domains or subregions. For example, a substrate-binding region can be used that interacts with a different substrate than that which is recognized by the native kinase. Accordingly, a different set of signal transduction components is available as an endpoint assay for activation. This allows for assays to be performed in other than the specific host cell from which the kinase is derived.

The proteins of the present invention are also useful in competition binding assays in methods designed to discover compounds that interact with the kinase (e.g. binding partners and/or ligands). Thus, a compound is exposed to a kinase polypeptide under conditions that allow the compound to bind or to otherwise interact with the polypeptide. Soluble kinase polypeptide is also added to the mixture. If the test compound interacts with the soluble kinase polypeptide, it decreases the amount of complex formed or activity from the kinase target. This type of assay is particularly useful in cases in which compounds are sought that interact with specific regions of the kinase. Thus, the soluble polypeptide that competes with the target kinase region is designed to contain peptide sequences corresponding to the region of interest.

To perform cell free drug screening assays, it is sometimes desirable to immobilize either the kinase protein, or fragment, or its target molecule to facilitate separation of complexes from uncomplexed forms of one or both of the proteins, as well as to accommodate automation of the assay.

Techniques for immobilizing proteins on matrices can be used in the drug screening assays. In one embodiment, a fusion protein can be provided which adds a domain that allows the protein to be bound to a matrix. For example, glutathione-S-transferase fusion proteins can be adsorbed onto glutathione sepharose beads (Sigma Chemical, St. Louis, MO) or glutathione derivatized microtitre

plates, which are then combined with the cell lysates (e.g., <sup>35</sup>S-labeled) and the candidate compound, and the mixture incubated under conditions conducive to complex formation (e.g., at physiological conditions for salt and pH). Following incubation, the beads are washed to remove any unbound label, and the matrix immobilized and radiolabel determined directly, or in the  
5 supernatant after the complexes are dissociated. Alternatively, the complexes can be dissociated from the matrix, separated by SDS-PAGE, and the level of kinase-binding protein found in the bead fraction quantitated from the gel using standard electrophoretic techniques. For example, either the polypeptide or its target molecule can be immobilized utilizing conjugation of biotin and streptavidin using techniques well known in the art. Alternatively, antibodies reactive with the  
10 protein but which do not interfere with binding of the protein to its target molecule can be derivatized to the wells of the plate, and the protein trapped in the wells by antibody conjugation. Preparations of a kinase-binding protein and a candidate compound are incubated in the kinase protein-presenting wells and the amount of complex trapped in the well can be quantitated. Methods for detecting such complexes, in addition to those described above for the GST-  
15 immobilized complexes, include immunodetection of complexes using antibodies reactive with the kinase protein target molecule, or which are reactive with kinase protein and compete with the target molecule, as well as enzyme-linked assays which rely on detecting an enzymatic activity associated with the target molecule.

Agents that modulate one of the kinases of the present invention can be identified using one  
20 or more of the above assays, alone or in combination. It is generally preferable to use a cell-based or cell free system first and then confirm activity in an animal or other model system. Such model systems are well known in the art and can readily be employed in this context.

Modulators of kinase protein activity identified according to these drug screening assays can be used to treat a subject with a disorder mediated by the kinase pathway, by treating cells or tissues  
25 that express the kinase. Experimental data as provided in Figure 1 indicates expression in the human placenta, kidney, lung, skeletal muscle, heart, fetal brain, and colon carcinoma. These methods of treatment include the steps of administering a modulator of kinase activity in a pharmaceutical composition to a subject in need of such treatment, the modulator being identified as described herein.

30 In yet another aspect of the invention, the kinase proteins can be used as "bait proteins" in a two-hybrid assay or three-hybrid assay (see, e.g., U.S. Patent No. 5,283,317; Zervos *et al.* (1993) *Cell* 72:223-232; Madura *et al.* (1993) *J. Biol. Chem.* 268:12046-12054; Bartel *et al.* (1993) *Biotechniques* 14:920-924; Iwabuchi *et al.* (1993) *Oncogene* 8:1693-1696; and Brent

WO94/10300), to identify other proteins, which bind to or interact with the kinase and are involved in kinase activity. Such kinase-binding proteins are also likely to be involved in the propagation of signals by the kinase proteins or kinase targets as, for example, downstream elements of a kinase-mediated signaling pathway. Alternatively, such kinase-binding proteins are likely to be kinase inhibitors.

The two-hybrid system is based on the modular nature of most transcription factors, which consist of separable DNA-binding and activation domains. Briefly, the assay utilizes two different DNA constructs. In one construct, the gene that codes for a kinase protein is fused to a gene encoding the DNA binding domain of a known transcription factor (e.g., GAL-4). In the other construct, a DNA sequence, from a library of DNA sequences, that encodes an unidentified protein ("prey" or "sample") is fused to a gene that codes for the activation domain of the known transcription factor. If the "bait" and the "prey" proteins are able to interact, *in vivo*, forming a kinase-dependent complex, the DNA-binding and activation domains of the transcription factor are brought into close proximity. This proximity allows transcription of a reporter gene (e.g., LacZ) which is operably linked to a transcriptional regulatory site responsive to the transcription factor. Expression of the reporter gene can be detected and cell colonies containing the functional transcription factor can be isolated and used to obtain the cloned gene which encodes the protein which interacts with the kinase protein.

This invention further pertains to novel agents identified by the above-described screening assays. Accordingly, it is within the scope of this invention to further use an agent identified as described herein in an appropriate animal model. For example, an agent identified as described herein (e.g., a kinase-modulating agent, an antisense kinase nucleic acid molecule, a kinase-specific antibody, or a kinase-binding partner) can be used in an animal or other model to determine the efficacy, toxicity, or side effects of treatment with such an agent. Alternatively, an agent identified as described herein can be used in an animal or other model to determine the mechanism of action of such an agent. Furthermore, this invention pertains to uses of novel agents identified by the above-described screening assays for treatments as described herein.

The kinase proteins of the present invention are also useful to provide a target for diagnosing a disease or predisposition to disease mediated by the peptide. Accordingly, the invention provides methods for detecting the presence, or levels of, the protein (or encoding mRNA) in a cell, tissue, or organism. Experimental data as provided in Figure 1 indicates expression in the human placenta, kidney, lung, skeletal muscle, heart, fetal brain, and colon carcinoma. The method involves contacting a biological sample with a compound capable of

interacting with the kinase protein such that the interaction can be detected. Such an assay can be provided in a single detection format or a multi-detection format such as an antibody chip array.

One agent for detecting a protein in a sample is an antibody capable of selectively binding to protein. A biological sample includes tissues, cells and biological fluids isolated from a subject, as well as tissues, cells and fluids present within a subject.

The peptides of the present invention also provide targets for diagnosing active protein activity, disease, or predisposition to disease, in a patient having a variant peptide, particularly activities and conditions that are known for other members of the family of proteins to which the present one belongs. Thus, the peptide can be isolated from a biological sample and assayed for the presence of a genetic mutation that results in aberrant peptide. This includes amino acid substitution, deletion, insertion, rearrangement, (as the result of aberrant splicing events), and inappropriate post-translational modification. Analytic methods include altered electrophoretic mobility, altered tryptic peptide digest, altered kinase activity in cell-based or cell-free assay, alteration in substrate or antibody-binding pattern, altered isoelectric point, direct amino acid sequencing, and any other of the known assay techniques useful for detecting mutations in a protein. Such an assay can be provided in a single detection format or a multi-detection format such as an antibody chip array.

*In vitro* techniques for detection of peptide include enzyme linked immunosorbent assays (ELISAs), Western blots, immunoprecipitations and immunofluorescence using a detection reagent, such as an antibody or protein binding agent. Alternatively, the peptide can be detected *in vivo* in a subject by introducing into the subject a labeled anti-peptide antibody or other types of detection agent. For example, the antibody can be labeled with a radioactive marker whose presence and location in a subject can be detected by standard imaging techniques. Particularly useful are methods that detect the allelic variant of a peptide expressed in a subject and methods which detect fragments of a peptide in a sample.

The peptides are also useful in pharmacogenomic analysis. Pharmacogenomics deal with clinically significant hereditary variations in the response to drugs due to altered drug disposition and abnormal action in affected persons. See, e.g., Eichelbaum, M. (*Clin. Exp. Pharmacol. Physiol.* 23(10-11):983-985 (1996)), and Linder, M.W. (*Clin. Chem.* 43(2):254-266 (1997)). The clinical outcomes of these variations result in severe toxicity of therapeutic drugs in certain individuals or therapeutic failure of drugs in certain individuals as a result of individual variation in metabolism. Thus, the genotype of the individual can determine the way a therapeutic compound acts on the body or the way the body metabolizes the compound. Further, the activity of drug metabolizing

enzymes effects both the intensity and duration of drug action. Thus, the pharmacogenomics of the individual permit the selection of effective compounds and effective dosages of such compounds for prophylactic or therapeutic treatment based on the individual's genotype. The discovery of genetic polymorphisms in some drug metabolizing enzymes has explained why some patients do not obtain the expected drug effects, show an exaggerated drug effect, or experience serious toxicity from standard drug dosages. Polymorphisms can be expressed in the phenotype of the extensive metabolizer and the phenotype of the poor metabolizer. Accordingly, genetic polymorphism may lead to allelic protein variants of the kinase protein in which one or more of the kinase functions in one population is different from those in another population. The peptides thus allow a target to ascertain a genetic predisposition that can affect treatment modality. Thus, in a ligand-based treatment, polymorphism may give rise to amino terminal extracellular domains and/or other substrate-binding regions that are more or less active in substrate binding, and kinase activation. Accordingly, substrate dosage would necessarily be modified to maximize the therapeutic effect within a given population containing a polymorphism. As an alternative to genotyping, specific polymorphic peptides could be identified.

The peptides are also useful for treating a disorder characterized by an absence of, inappropriate, or unwanted expression of the protein. Experimental data as provided in Figure 1 indicates expression in the human placenta, kidney, lung, skeletal muscle, heart, fetal brain, and colon carcinoma. Accordingly, methods for treatment include the use of the kinase protein or fragments.

### Antibodies

The invention also provides antibodies that selectively bind to one of the peptides of the present invention, a protein comprising such a peptide, as well as variants and fragments thereof. As used herein, an antibody selectively binds a target peptide when it binds the target peptide and does not significantly bind to unrelated proteins. An antibody is still considered to selectively bind a peptide even if it also binds to other proteins that are not substantially homologous with the target peptide so long as such proteins share homology with a fragment or domain of the peptide target of the antibody. In this case, it would be understood that antibody binding to the peptide is still selective despite some degree of cross-reactivity.

As used herein, an antibody is defined in terms consistent with that recognized within the art: they are multi-subunit proteins produced by a mammalian organism in response to an antigen challenge. The antibodies of the present invention include polyclonal antibodies and monoclonal

antibodies, as well as fragments of such antibodies, including, but not limited to, Fab or F(ab')<sub>2</sub>, and Fv fragments.

Many methods are known for generating and/or identifying antibodies to a given target peptide. Several such methods are described by Harlow, *Antibodies*, Cold Spring Harbor Press, (1989).

In general, to generate antibodies, an isolated peptide is used as an immunogen and is administered to a mammalian organism, such as a rat, rabbit or mouse. The full-length protein, an antigenic peptide fragment or a fusion protein can be used. Particularly important fragments are those covering functional domains, such as the domains identified in Figure 2, and domain of sequence homology or divergence amongst the family, such as those that can readily be identified using protein alignment methods and as presented in the Figures.

Antibodies are preferably prepared from regions or discrete fragments of the kinase proteins. Antibodies can be prepared from any region of the peptide as described herein. However, preferred regions will include those involved in function/activity and/or kinase/binding partner interaction. Figure 2 can be used to identify particularly important regions while sequence alignment can be used to identify conserved and unique sequence fragments.

An antigenic fragment will typically comprise at least 8 contiguous amino acid residues. The antigenic peptide can comprise, however, at least 10, 12, 14, 16 or more amino acid residues. Such fragments can be selected on a physical property, such as fragments correspond to regions that are located on the surface of the protein, e.g., hydrophilic regions or can be selected based on sequence uniqueness (see Figure 2).

Detection on an antibody of the present invention can be facilitated by coupling (i.e., physically linking) the antibody to a detectable substance. Examples of detectable substances include various enzymes, prosthetic groups, fluorescent materials, luminescent materials, bioluminescent materials, and radioactive materials. Examples of suitable enzymes include horseradish peroxidase, alkaline phosphatase,  $\beta$ -galactosidase, or acetylcholinesterase; examples of suitable prosthetic group complexes include streptavidin/biotin and avidin/biotin; examples of suitable fluorescent materials include umbelliferone, fluorescein, fluorescein isothiocyanate, rhodamine, dichlorotriazinylamine fluorescein, dansyl chloride or phycoerythrin; an example of a luminescent material includes luminol; examples of bioluminescent materials include luciferase, luciferin, and aequorin, and examples of suitable radioactive material include <sup>125</sup>I, <sup>131</sup>I, <sup>35</sup>S or <sup>3</sup>H.

### Antibody Uses

The antibodies can be used to isolate one of the proteins of the present invention by standard techniques, such as affinity chromatography or immunoprecipitation. The antibodies can facilitate the purification of the natural protein from cells and recombinantly produced protein expressed in host cells. In addition, such antibodies are useful to detect the presence of one of the proteins of the present invention in cells or tissues to determine the pattern of expression of the protein among various tissues in an organism and over the course of normal development. Experimental data as provided in Figure 1 indicates that kinase proteins of the present invention are expressed in the human placenta, kidney, lung, skeletal muscle, heart, fetal brain, and colon carcinoma. Specifically, a virtual northern blot shows expression in human colon carcinoma. In addition, PCR-based tissue screening panel indicates expression in human placenta, kidney, lung, skeletal muscle, heart, and fetal brain. Further, such antibodies can be used to detect protein *in situ*, *in vitro*, or in a cell lysate or supernatant in order to evaluate the abundance and pattern of expression. Also, such antibodies can be used to assess abnormal tissue distribution or abnormal expression during development or progression of a biological condition. Antibody detection of circulating fragments of the full length protein can be used to identify turnover.

Further, the antibodies can be used to assess expression in disease states such as in active stages of the disease or in an individual with a predisposition toward disease related to the protein's function. When a disorder is caused by an inappropriate tissue distribution, developmental expression, level of expression of the protein, or expressed/processed form, the antibody can be prepared against the normal protein. Experimental data as provided in Figure 1 indicates expression in the human placenta, kidney, lung, skeletal muscle, heart, fetal brain, and colon carcinoma. If a disorder is characterized by a specific mutation in the protein, antibodies specific for this mutant protein can be used to assay for the presence of the specific mutant protein.

The antibodies can also be used to assess normal and aberrant subcellular localization of cells in the various tissues in an organism. Experimental data as provided in Figure 1 indicates expression in the human placenta, kidney, lung, skeletal muscle, heart, fetal brain, and colon carcinoma. The diagnostic uses can be applied, not only in genetic testing, but also in monitoring a treatment modality. Accordingly, where treatment is ultimately aimed at correcting expression level or the presence of aberrant sequence and aberrant tissue distribution or developmental expression, antibodies directed against the protein or relevant fragments can be used to monitor therapeutic efficacy.

Additionally, antibodies are useful in pharmacogenomic analysis. Thus, antibodies prepared against polymorphic proteins can be used to identify individuals that require modified treatment modalities. The antibodies are also useful as diagnostic tools as an immunological marker for aberrant protein analyzed by electrophoretic mobility, isoelectric point, tryptic peptide digest, and other physical assays known to those in the art.

The antibodies are also useful for tissue typing. Experimental data as provided in Figure 1 indicates expression in the human placenta, kidney, lung, skeletal muscle, heart, fetal brain, and colon carcinoma. Thus, where a specific protein has been correlated with expression in a specific tissue, antibodies that are specific for this protein can be used to identify a tissue type.

The antibodies are also useful for inhibiting protein function, for example, blocking the binding of the kinase peptide to a binding partner such as a substrate. These uses can also be applied in a therapeutic context in which treatment involves inhibiting the protein's function. An antibody can be used, for example, to block binding, thus modulating (agonizing or antagonizing) the peptides activity. Antibodies can be prepared against specific fragments containing sites required for function or against intact protein that is associated with a cell or cell membrane. See Figure 2 for structural information relating to the proteins of the present invention.

The invention also encompasses kits for using antibodies to detect the presence of a protein in a biological sample. The kit can comprise antibodies such as a labeled or labelable antibody and a compound or agent for detecting protein in a biological sample; means for determining the amount of protein in the sample; means for comparing the amount of protein in the sample with a standard; and instructions for use. Such a kit can be supplied to detect a single protein or epitope or can be configured to detect one of a multitude of epitopes, such as in an antibody detection array. Arrays are described in detail below for nucleic acid arrays and similar methods have been developed for antibody arrays.

### Nucleic Acid Molecules

The present invention further provides isolated nucleic acid molecules that encode a kinase peptide or protein of the present invention (cDNA, transcript and genomic sequence). Such nucleic acid molecules will consist of, consist essentially of, or comprise a nucleotide sequence that encodes one of the kinase peptides of the present invention, an allelic variant thereof, or an ortholog or paralog thereof.



As used herein, an "isolated" nucleic acid molecule is one that is separated from other nucleic acid present in the natural source of the nucleic acid. Preferably, an "isolated" nucleic acid is free of sequences which naturally flank the nucleic acid (i.e., sequences located at the 5' and 3' ends of the nucleic acid) in the genomic DNA of the organism from which the nucleic acid is  
5 derived. However, there can be some flanking nucleotide sequences, for example up to about 5KB, 4KB, 3KB, 2KB, or 1KB or less, particularly contiguous peptide encoding sequences and peptide encoding sequences within the same gene but separated by introns in the genomic sequence. The important point is that the nucleic acid is isolated from remote and unimportant flanking sequences such that it can be subjected to the specific manipulations described herein such as recombinant  
10 expression, preparation of probes and primers, and other uses specific to the nucleic acid sequences.

Moreover, an "isolated" nucleic acid molecule, such as a transcript/cDNA molecule, can be substantially free of other cellular material, or culture medium when produced by recombinant techniques, or chemical precursors or other chemicals when chemically synthesized. However, the nucleic acid molecule can be fused to other coding or regulatory sequences and still be considered  
15 isolated.

For example, recombinant DNA molecules contained in a vector are considered isolated. Further examples of isolated DNA molecules include recombinant DNA molecules maintained in heterologous host cells or purified (partially or substantially) DNA molecules in solution. Isolated RNA molecules include *in vivo* or *in vitro* RNA transcripts of the isolated DNA molecules of the  
20 present invention. Isolated nucleic acid molecules according to the present invention further include such molecules produced synthetically.

Accordingly, the present invention provides nucleic acid molecules that consist of the nucleotide sequence shown in Figure 1 or 3 (SEQ ID NO:1, transcript sequence and SEQ ID NO:3, genomic sequence), or any nucleic acid molecule that encodes the protein provided in Figure 2,  
25 SEQ ID NO:2. A nucleic acid molecule consists of a nucleotide sequence when the nucleotide sequence is the complete nucleotide sequence of the nucleic acid molecule.

The present invention further provides nucleic acid molecules that consist essentially of the nucleotide sequence shown in Figure 1 or 3 (SEQ ID NO:1, transcript sequence and SEQ ID NO:3, genomic sequence), or any nucleic acid molecule that encodes the protein provided in Figure 2,  
30 SEQ ID NO:2. A nucleic acid molecule consists essentially of a nucleotide sequence when such a nucleotide sequence is present with only a few additional nucleic acid residues in the final nucleic acid molecule.

The present invention further provides nucleic acid molecules that comprise the nucleotide sequences shown in Figure 1 or 3 (SEQ ID NO:1, transcript sequence and SEQ ID NO:3, genomic sequence), or any nucleic acid molecule that encodes the protein provided in Figure 2, SEQ ID NO:2. A nucleic acid molecule comprises a nucleotide sequence when the nucleotide sequence is at least part of the final nucleotide sequence of the nucleic acid molecule. In such a fashion, the nucleic acid molecule can be only the nucleotide sequence or have additional nucleic acid residues, such as nucleic acid residues that are naturally associated with it or heterologous nucleotide sequences. Such a nucleic acid molecule can have a few additional nucleotides or can comprises several hundred or more additional nucleotides. A brief description of how various types of these nucleic acid molecules can be readily made/isolated is provided below.

In Figures 1 and 3, both coding and non-coding sequences are provided. Because of the source of the present invention, humans genomic sequence (Figure 3) and cDNA/transcript sequences (Figure 1), the nucleic acid molecules in the Figures will contain genomic intronic sequences, 5' and 3' non-coding sequences, gene regulatory regions and non-coding intergenic sequences. In general such sequence features are either noted in Figures 1 and 3 or can readily be identified using computational tools known in the art. As discussed below, some of the non-coding regions, particularly gene regulatory elements such as promoters, are useful for a variety of purposes, e.g. control of heterologous gene expression, target for identifying gene activity modulating compounds, and are particularly claimed as fragments of the genomic sequence provided herein.

The isolated nucleic acid molecules can encode the mature protein plus additional amino or carboxyl-terminal amino acids, or amino acids interior to the mature peptide (when the mature form has more than one peptide chain, for instance). Such sequences may play a role in processing of a protein from precursor to a mature form, facilitate protein trafficking, prolong or shorten protein half-life or facilitate manipulation of a protein for assay or production, among other things. As generally is the case *in situ*, the additional amino acids may be processed away from the mature protein by cellular enzymes.

As mentioned above, the isolated nucleic acid molecules include, but are not limited to, the sequence encoding the kinase peptide alone, the sequence encoding the mature peptide and additional coding sequences, such as a leader or secretory sequence (e.g., a pre-pro or pro-protein sequence), the sequence encoding the mature peptide, with or without the additional coding sequences, plus additional non-coding sequences, for example introns and non-coding 5' and 3' sequences such as transcribed but non-translated sequences that play a role in transcription, mRNA

processing (including splicing and polyadenylation signals), ribosome binding and stability of mRNA. In addition, the nucleic acid molecule may be fused to a marker sequence encoding, for example, a peptide that facilitates purification.

Isolated nucleic acid molecules can be in the form of RNA, such as mRNA, or in the form of DNA, including cDNA and genomic DNA obtained by cloning or produced by chemical synthetic techniques or by a combination thereof. The nucleic acid, especially DNA, can be double-stranded or single-stranded. Single-stranded nucleic acid can be the coding strand (sense strand) or the non-coding strand (anti-sense strand):

The invention further provides nucleic acid molecules that encode fragments of the peptides of the present invention as well as nucleic acid molecules that encode obvious variants of the kinase proteins of the present invention that are described above. Such nucleic acid molecules may be naturally occurring, such as allelic variants (same locus), paralogs (different locus), and orthologs (different organism), or may be constructed by recombinant DNA methods or by chemical synthesis. Such non-naturally occurring variants may be made by mutagenesis techniques, including those applied to nucleic acid molecules, cells, or organisms. Accordingly, as discussed above, the variants can contain nucleotide substitutions, deletions, inversions and insertions. Variation can occur in either or both the coding and non-coding regions. The variations can produce both conservative and non-conservative amino acid substitutions.

The present invention further provides non-coding fragments of the nucleic acid molecules provided in Figures 1 and 3. Preferred non-coding fragments include, but are not limited to, promoter sequences, enhancer sequences, gene modulating sequences and gene termination sequences. Such fragments are useful in controlling heterologous gene expression and in developing screens to identify gene-modulating agents. A promoter can readily be identified as being 5' to the ATG start site in the genomic sequence provided in Figure 3.

A fragment comprises a contiguous nucleotide sequence greater than 12 or more nucleotides. Further, a fragment could be at least 30, 40, 50, 100, 250 or 500 nucleotides in length. The length of the fragment will be based on its intended use. For example, the fragment can encode epitope bearing regions of the peptide, or can be useful as DNA probes and primers. Such fragments can be isolated using the known nucleotide sequence to synthesize an oligonucleotide probe. A labeled probe can then be used to screen a cDNA library, genomic DNA library, or mRNA to isolate nucleic acid corresponding to the coding region. Further, primers can be used in PCR reactions to clone specific regions of gene.

A probe/primer typically comprises substantially a purified oligonucleotide or oligonucleotide pair. The oligonucleotide typically comprises a region of nucleotide sequence that hybridizes under stringent conditions to at least about 12, 20, 25, 40, 50 or more consecutive nucleotides.

5       Orthologs, homologs, and allelic variants can be identified using methods well known in the art. As described in the Peptide Section, these variants comprise a nucleotide sequence encoding a peptide that is typically 60-70%, 70-80%, 80-90%, and more typically at least about 90-95% or more homologous to the nucleotide sequence shown in the Figure sheets or a fragment of this sequence. Such nucleic acid molecules can readily be identified as being able to hybridize under  
10   moderate to stringent conditions, to the nucleotide sequence shown in the Figure sheets or a fragment of the sequence. Allelic variants can readily be determined by genetic locus of the encoding gene. As indicated by the data presented in Figure 3, the map position was determined to be on chromosome 1 by ePCR, and confirmed with radiation hybrid mapping. As indicated by the data presented in Figure 3, the gene provided by the present invention encoding a novel phosphatase  
15   maps to public BAC AC AC023889, which is known to be located on human chromosome 1.

Figure 3 provides information on SNPs that have been identified in a gene encoding the kinase protein of the present invention. 6 SNP variants were found, and all SNPs in exons, of which 3 of these cause changes in the amino acid sequence (i.e., nonsynonymous SNPs). The changes in the amino acid sequence that these SNPs cause is indicated in Figure 3 and can readily be  
20   determined using the universal genetic code and the protein sequence provided in Figure 2 as a reference.

As used herein, the term "hybridizes under stringent conditions" is intended to describe conditions for hybridization and washing under which nucleotide sequences encoding a peptide at least 60-70% homologous to each other typically remain hybridized to each other. The conditions  
25   can be such that sequences at least about 60%, at least about 70%, or at least about 80% or more homologous to each other typically remain hybridized to each other. Such stringent conditions are known to those skilled in the art and can be found in *Current Protocols in Molecular Biology*, John Wiley & Sons, N.Y. (1989), 6.3.1-6.3.6. One example of stringent hybridization conditions are hybridization in 6X sodium chloride/sodium citrate (SSC) at about 45C, followed by one or more  
30   washes in 0.2 X SSC, 0.1% SDS at 50-65C. Examples of moderate to low stringency hybridization conditions are well known in the art.

### Nucleic Acid Molecule Uses

The nucleic acid molecules of the present invention are useful for probes, primers, chemical intermediates, and in biological assays. The nucleic acid molecules are useful as a hybridization probe for messenger RNA, transcript/cDNA and genomic DNA to isolate full-length cDNA and genomic clones encoding the peptide described in Figure 2 and to isolate cDNA and genomic clones that correspond to variants (alleles, orthologs, etc.) producing the same or related peptides shown in Figure 2. 6 SNPs, have been identified in the gene encoding the kinase protein provided by the present invention and are given in Figure 3.

The probe can correspond to any sequence along the entire length of the nucleic acid molecules provided in the Figures. Accordingly, it could be derived from 5' noncoding regions, the coding region, and 3' noncoding regions. However, as discussed, fragments are not to be construed as encompassing fragments disclosed prior to the present invention.

The nucleic acid molecules are also useful as primers for PCR to amplify any given region of a nucleic acid molecule and are useful to synthesize antisense molecules of desired length and sequence.

The nucleic acid molecules are also useful for constructing recombinant vectors. Such vectors include expression vectors that express a portion of, or all of, the peptide sequences. Vectors also include insertion vectors, used to integrate into another nucleic acid molecule sequence, such as into the cellular genome, to alter *in situ* expression of a gene and/or gene product. For example, an endogenous coding sequence can be replaced via homologous recombination with all or part of the coding region containing one or more specifically introduced mutations.

The nucleic acid molecules are also useful for expressing antigenic portions of the proteins.

The nucleic acid molecules are also useful as probes for determining the chromosomal positions of the nucleic acid molecules by means of *in situ* hybridization methods. As indicated by the data presented in Figure 3, the map position was determined to be on chromosome 1 by ePCR, and confirmed with radiation hybrid mapping. As indicated by the data presented in Figure 3, the gene provided by the present invention encoding a novel phosphatase maps to public BAC AC AC023889, which is known to be located on human chromosome 1.

The nucleic acid molecules are also useful in making vectors containing the gene regulatory regions of the nucleic acid molecules of the present invention.

The nucleic acid molecules are also useful for designing ribozymes corresponding to all, or a part, of the mRNA produced from the nucleic acid molecules described herein.

The nucleic acid molecules are also useful for making vectors that express part, or all, of the peptides.

The nucleic acid molecules are also useful for constructing host cells expressing a part, or all, of the nucleic acid molecules and peptides.

5       The nucleic acid molecules are also useful for constructing transgenic animals expressing all, or a part, of the nucleic acid molecules and peptides.

      The nucleic acid molecules are also useful as hybridization probes for determining the presence, level, form and distribution of nucleic acid expression. Experimental data as provided in Figure 1 indicates that kinase proteins of the present invention are expressed in the human placenta, kidney, lung, skeletal muscle, heart, fetal brain, and colon carcinoma. Specifically, a virtual northern blot shows expression in human colon carcinoma. In addition, PCR-based tissue screening panel indicates expression in human placenta, kidney, lung, skeletal muscle, heart, and fetal brain. Accordingly, the probes can be used to detect the presence of, or to determine levels of, a specific nucleic acid molecule in cells, tissues, and in organisms. The nucleic acid whose level is  
15       determined can be DNA or RNA. Accordingly, probes corresponding to the peptides described herein can be used to assess expression and/or gene copy number in a given cell, tissue, or organism. These uses are relevant for diagnosis of disorders involving an increase or decrease in kinase protein expression relative to normal results.

*In vitro* techniques for detection of mRNA include Northern hybridizations and *in situ*  
20       hybridizations. *In vitro* techniques for detecting DNA includes Southern hybridizations and *in situ* hybridization.

      Probes can be used as a part of a diagnostic test kit for identifying cells or tissues that express a kinase protein, such as by measuring a level of a kinase-encoding nucleic acid in a sample of cells from a subject e.g., mRNA or genomic DNA, or determining if a kinase gene has been  
25       mutated. Experimental data as provided in Figure 1 indicates that kinase proteins of the present invention are expressed in the human placenta, kidney, lung, skeletal muscle, heart, fetal brain, and colon carcinoma. Specifically, a virtual northern blot shows expression in human colon carcinoma. In addition, PCR-based tissue screening panel indicates expression in human placenta, kidney, lung, skeletal muscle, heart, and fetal brain.

30       Nucleic acid expression assays are useful for drug screening to identify compounds that modulate kinase nucleic acid expression.

      The invention thus provides a method for identifying a compound that can be used to treat a disorder associated with nucleic acid expression of the kinase gene, particularly biological and

pathological processes that are mediated by the kinase in cells and tissues that express it.

Experimental data as provided in Figure 1 indicates expression in the human placenta, kidney, lung, skeletal muscle, heart, fetal brain, and colon carcinoma. The method typically includes assaying the ability of the compound to modulate the expression of the kinase nucleic acid and thus identifying a compound that can be used to treat a disorder characterized by undesired kinase nucleic acid expression. The assays can be performed in cell-based and cell-free systems. Cell-based assays include cells naturally expressing the kinase nucleic acid or recombinant cells genetically engineered to express specific nucleic acid sequences.

The assay for kinase nucleic acid expression can involve direct assay of nucleic acid levels, such as mRNA levels, or on collateral compounds involved in the signal pathway. Further, the expression of genes that are up- or down-regulated in response to the kinase protein signal pathway can also be assayed. In this embodiment the regulatory regions of these genes can be operably linked to a reporter gene such as luciferase.

Thus, modulators of kinase gene expression can be identified in a method wherein a cell is contacted with a candidate compound and the expression of mRNA determined. The level of expression of kinase mRNA in the presence of the candidate compound is compared to the level of expression of kinase mRNA in the absence of the candidate compound. The candidate compound can then be identified as a modulator of nucleic acid expression based on this comparison and be used, for example to treat a disorder characterized by aberrant nucleic acid expression. When expression of mRNA is statistically significantly greater in the presence of the candidate compound than in its absence, the candidate compound is identified as a stimulator of nucleic acid expression. When nucleic acid expression is statistically significantly less in the presence of the candidate compound than in its absence, the candidate compound is identified as an inhibitor of nucleic acid expression.

The invention further provides methods of treatment, with the nucleic acid as a target, using a compound identified through drug screening as a gene modulator to modulate kinase nucleic acid expression in cells and tissues that express the kinase. Experimental data as provided in Figure 1 indicates that kinase proteins of the present invention are expressed in the human placenta, kidney, lung, skeletal muscle, heart, fetal brain, and colon carcinoma. Specifically, a virtual northern blot shows expression in human colon carcinoma. In addition, PCR-based tissue screening panel indicates expression in human placenta, kidney, lung, skeletal muscle, heart, and fetal brain. Modulation includes both up-regulation (i.e. activation or agonization) or down-regulation (suppression or antagonization) or nucleic acid expression.

Alternatively, a modulator for kinase nucleic acid expression can be a small molecule or drug identified using the screening assays described herein as long as the drug or small molecule inhibits the kinase nucleic acid expression in the cells and tissues that express the protein. Experimental data as provided in Figure 1 indicates expression in the human placenta, kidney, lung, skeletal muscle, heart, fetal brain, and colon carcinoma.

The nucleic acid molecules are also useful for monitoring the effectiveness of modulating compounds on the expression or activity of the kinase gene in clinical trials or in a treatment regimen. Thus, the gene expression pattern can serve as a barometer for the continuing effectiveness of treatment with the compound, particularly with compounds to which a patient can develop resistance. The gene expression pattern can also serve as a marker indicative of a physiological response of the affected cells to the compound. Accordingly, such monitoring would allow either increased administration of the compound or the administration of alternative compounds to which the patient has not become resistant. Similarly, if the level of nucleic acid expression falls below a desirable level, administration of the compound could be commensurately decreased.

The nucleic acid molecules are also useful in diagnostic assays for qualitative changes in kinase nucleic acid expression, and particularly in qualitative changes that lead to pathology. The nucleic acid molecules can be used to detect mutations in kinase genes and gene expression products such as mRNA. The nucleic acid molecules can be used as hybridization probes to detect naturally occurring genetic mutations in the kinase gene and thereby to determine whether a subject with the mutation is at risk for a disorder caused by the mutation. Mutations include deletion, addition, or substitution of one or more nucleotides in the gene, chromosomal rearrangement, such as inversion or transposition, modification of genomic DNA, such as aberrant methylation patterns or changes in gene copy number, such as amplification. Detection of a mutated form of the kinase gene associated with a dysfunction provides a diagnostic tool for an active disease or susceptibility to disease when the disease results from overexpression, underexpression, or altered expression of a kinase protein.

Individuals carrying mutations in the kinase gene can be detected at the nucleic acid level by a variety of techniques. Figure 3 provides information on SNPs that have been identified in a gene encoding the kinase protein of the present invention. 6 SNP variants were found, and all SNPs in exons, of which 3 of these cause changes in the amino acid sequence (i.e., nonsynonymous SNPs). The changes in the amino acid sequence that these SNPs cause is indicated in Figure 3 and can readily be determined using the universal genetic code and the protein sequence provided in Figure



2 as a reference. As indicated by the data presented in Figure 3, the map position was determined to be on chromosome 1 by ePCR, and confirmed with radiation hybrid mapping. As indicated by the data presented in Figure 3, the gene provided by the present invention encoding a novel phosphatase maps to public BAC AC AC023889, which is known to be located on human chromosome 1.

5 Genomic DNA can be analyzed directly or can be amplified by using PCR prior to analysis. RNA or cDNA can be used in the same way. In some uses, detection of the mutation involves the use of a probe/primer in a polymerase chain reaction (PCR) (see, e.g. U.S. Patent Nos. 4,683,195 and 4,683,202), such as anchor PCR or RACE PCR, or, alternatively, in a ligation chain reaction (LCR) (see, e.g., Landegran *et al.*, *Science* 241:1077-1080 (1988); and Nakazawa *et al.*, *PNAS* 91:360-364  
10 (1994)), the latter of which can be particularly useful for detecting point mutations in the gene (see Abravaya *et al.*, *Nucleic Acids Res.* 23:675-682 (1995)). This method can include the steps of collecting a sample of cells from a patient, isolating nucleic acid (e.g., genomic, mRNA or both) from the cells of the sample, contacting the nucleic acid sample with one or more primers which specifically hybridize to a gene under conditions such that hybridization and amplification of the  
15 gene (if present) occurs, and detecting the presence or absence of an amplification product, or detecting the size of the amplification product and comparing the length to a control sample. Deletions and insertions can be detected by a change in size of the amplified product compared to the normal genotype. Point mutations can be identified by hybridizing amplified DNA to normal RNA or antisense DNA sequences.

20 Alternatively, mutations in a kinase gene can be directly identified, for example, by alterations in restriction enzyme digestion patterns determined by gel electrophoresis.

Further, sequence-specific ribozymes (U.S. Patent No. 5,498,531) can be used to score for the presence of specific mutations by development or loss of a ribozyme cleavage site. Perfectly matched sequences can be distinguished from mismatched sequences by nuclease cleavage  
25 digestion assays or by differences in melting temperature.

Sequence changes at specific locations can also be assessed by nuclease protection assays such as RNase and S1 protection or the chemical cleavage method. Furthermore, sequence differences between a mutant kinase gene and a wild-type gene can be determined by direct DNA sequencing. A variety of automated sequencing procedures can be utilized when performing the  
30 diagnostic assays (Naeve, C.W., (1995) *Biotechniques* 19:448), including sequencing by mass spectrometry (see, e.g., PCT International Publication No. WO 94/16101; Cohen *et al.*, *Adv. Chromatogr.* 36:127-162 (1996); and Griffin *et al.*, *Appl. Biochem. Biotechnol.* 38:147-159 (1993)).

Other methods for detecting mutations in the gene include methods in which protection from cleavage agents is used to detect mismatched bases in RNA/RNA or RNA/DNA duplexes (Myers *et al.*, *Science* 230:1242 (1985)); Cotton *et al.*, *PNAS* 85:4397 (1988); Saleeba *et al.*, *Meth. Enzymol.* 217:286-295 (1992)), electrophoretic mobility of mutant and wild type nucleic acid is compared (Orita *et al.*, *PNAS* 86:2766 (1989); Cotton *et al.*, *Mutat. Res.* 285:125-144 (1993); and Hayashi *et al.*, *Genet. Anal. Tech. Appl.* 9:73-79 (1992)), and movement of mutant or wild-type fragments in polyacrylamide gels containing a gradient of denaturant is assayed using denaturing gradient gel electrophoresis (Myers *et al.*, *Nature* 313:495 (1985)). Examples of other techniques for detecting point mutations include selective oligonucleotide hybridization, selective amplification, and selective primer extension.

The nucleic acid molecules are also useful for testing an individual for a genotype that while not necessarily causing the disease, nevertheless affects the treatment modality. Thus, the nucleic acid molecules can be used to study the relationship between an individual's genotype and the individual's response to a compound used for treatment (pharmacogenomic relationship).

Accordingly, the nucleic acid molecules described herein can be used to assess the mutation content of the kinase gene in an individual in order to select an appropriate compound or dosage regimen for treatment. Figure 3 provides information on SNPs that have been identified in a gene encoding the kinase protein of the present invention. 6 SNP variants were found, and all SNPs in exons, of which 3 of these cause changes in the amino acid sequence (i.e., nonsynonymous SNPs). The changes in the amino acid sequence that these SNPs cause is indicated in Figure 3 and can readily be determined using the universal genetic code and the protein sequence provided in Figure 2 as a reference.

Thus nucleic acid molecules displaying genetic variations that affect treatment provide a diagnostic target that can be used to tailor treatment in an individual. Accordingly, the production of recombinant cells and animals containing these polymorphisms allow effective clinical design of treatment compounds and dosage regimens.

The nucleic acid molecules are thus useful as antisense constructs to control kinase gene expression in cells, tissues, and organisms. A DNA antisense nucleic acid molecule is designed to be complementary to a region of the gene involved in transcription, preventing transcription and hence production of kinase protein. An antisense RNA or DNA nucleic acid molecule would hybridize to the mRNA and thus block translation of mRNA into kinase protein.

Alternatively, a class of antisense molecules can be used to inactivate mRNA in order to decrease expression of kinase nucleic acid. Accordingly, these molecules can treat a disorder

characterized by abnormal or undesired kinase nucleic acid expression. This technique involves cleavage by means of ribozymes containing nucleotide sequences complementary to one or more regions in the mRNA that attenuate the ability of the mRNA to be translated. Possible regions include coding regions and particularly coding regions corresponding to the catalytic and other functional activities of the kinase protein, such as substrate binding.

The nucleic acid molecules also provide vectors for gene therapy in patients containing cells that are aberrant in kinase gene expression. Thus, recombinant cells, which include the patient's cells that have been engineered *ex vivo* and returned to the patient, are introduced into an individual where the cells produce the desired kinase protein to treat the individual.

The invention also encompasses kits for detecting the presence of a kinase nucleic acid in a biological sample. Experimental data as provided in Figure 1 indicates that kinase proteins of the present invention are expressed in the human placenta, kidney, lung, skeletal muscle, heart, fetal brain, and colon carcinoma. Specifically, a virtual northern blot shows expression in human colon carcinoma. In addition, PCR-based tissue screening panel indicates expression in human placenta, kidney, lung, skeletal muscle, heart, and fetal brain. For example, the kit can comprise reagents such as a labeled or labelable nucleic acid or agent capable of detecting kinase nucleic acid in a biological sample; means for determining the amount of kinase nucleic acid in the sample; and means for comparing the amount of kinase nucleic acid in the sample with a standard. The compound or agent can be packaged in a suitable container. The kit can further comprise instructions for using the kit to detect kinase protein mRNA or DNA.

#### Nucleic Acid Arrays

The present invention further provides nucleic acid detection kits, such as arrays or microarrays of nucleic acid molecules that are based on the sequence information provided in Figures 1 and 3 (SEQ ID NOS:1 and 3).

As used herein "Arrays" or "Microarrays" refers to an array of distinct polynucleotides or oligonucleotides synthesized on a substrate, such as paper, nylon or other type of membrane, filter, chip, glass slide, or any other suitable solid support. In one embodiment, the microarray is prepared and used according to the methods described in US Patent 5,837,832, Chee *et al.*, PCT application W095/11995 (Chee *et al.*), Lockhart, D. J. *et al.* (1996; Nat. Biotech. 14: 1675-1680) and Schena, M. *et al.* (1996; Proc. Natl. Acad. Sci. 93: 10614-10619), all of which are incorporated herein in their entirety by reference. In other embodiments, such arrays are produced by the methods described by Brown *et al.*, US Patent No. 5,807,522.

The microarray or detection kit is preferably composed of a large number of unique, single-stranded nucleic acid sequences, usually either synthetic antisense oligonucleotides or fragments of cDNAs, fixed to a solid support. The oligonucleotides are preferably about 6-60 nucleotides in length, more preferably 15-30 nucleotides in length, and most preferably about 20-25 nucleotides in length. For a certain type of microarray or detection kit, it may be preferable to use oligonucleotides that are only 7-20 nucleotides in length. The microarray or detection kit may contain oligonucleotides that cover the known 5', or 3', sequence, sequential oligonucleotides which cover the full length sequence; or unique oligonucleotides selected from particular areas along the length of the sequence. Polynucleotides used in the microarray or detection kit may be oligonucleotides that are specific to a gene or genes of interest.

In order to produce oligonucleotides to a known sequence for a microarray or detection kit, the gene(s) of interest (or an ORF identified from the contigs of the present invention) is typically examined using a computer algorithm which starts at the 5' or at the 3' end of the nucleotide sequence. Typical algorithms will then identify oligomers of defined length that are unique to the gene, have a GC content within a range suitable for hybridization, and lack predicted secondary structure that may interfere with hybridization. In certain situations it may be appropriate to use pairs of oligonucleotides on a microarray or detection kit. The "pairs" will be identical, except for one nucleotide that preferably is located in the center of the sequence. The second oligonucleotide in the pair (mismatched by one) serves as a control. The number of oligonucleotide pairs may range from two to one million. The oligomers are synthesized at designated areas on a substrate using a light-directed chemical process. The substrate may be paper, nylon or other type of membrane, filter, chip, glass slide or any other suitable solid support.

In another aspect, an oligonucleotide may be synthesized on the surface of the substrate by using a chemical coupling procedure and an ink jet application apparatus, as described in PCT application W095/251116 (Baldeschweiler *et al.*) which is incorporated herein in its entirety by reference. In another aspect, a "gridded" array analogous to a dot (or slot) blot may be used to arrange and link cDNA fragments or oligonucleotides to the surface of a substrate using a vacuum system, thermal, UV, mechanical or chemical bonding procedures. An array, such as those described above, may be produced by hand or by using available devices (slot blot or dot blot apparatus), materials (any suitable solid support), and machines (including robotic instruments), and may contain 8, 24, 96, 384, 1536, 6144 or more oligonucleotides, or any other

number between two and one million which lends itself to the efficient use of commercially available instrumentation.

In order to conduct sample analysis using a microarray or detection kit, the RNA or DNA from a biological sample is made into hybridization probes. The mRNA is isolated, and cDNA is produced and used as a template to make antisense RNA (aRNA). The aRNA is amplified in the presence of fluorescent nucleotides, and labeled probes are incubated with the microarray or detection kit so that the probe sequences hybridize to complementary oligonucleotides of the microarray or detection kit. Incubation conditions are adjusted so that hybridization occurs with precise complementary matches or with various degrees of less complementarity. After removal of nonhybridized probes, a scanner is used to determine the levels and patterns of fluorescence. The scanned images are examined to determine degree of complementarity and the relative abundance of each oligonucleotide sequence on the microarray or detection kit. The biological samples may be obtained from any bodily fluids (such as blood, urine, saliva, phlegm, gastric juices, etc.), cultured cells, biopsies, or other tissue preparations. A detection system may be used to measure the absence, presence, and amount of hybridization for all of the distinct sequences simultaneously. This data may be used for large-scale correlation studies on the sequences, expression patterns, mutations, variants, or polymorphisms among samples.

Using such arrays, the present invention provides methods to identify the expression of the kinase proteins/peptides of the present invention. In detail, such methods comprise incubating a test sample with one or more nucleic acid molecules and assaying for binding of the nucleic acid molecule with components within the test sample. Such assays will typically involve arrays comprising many genes, at least one of which is a gene of the present invention and or alleles of the kinase gene of the present invention. Figure 3 provides information on SNPs that have been identified in a gene encoding the kinase protein of the present invention. 6 SNP variants were found, and all SNPs in exons, of which 3 of these cause changes in the amino acid sequence (i.e., nonsynonymous SNPs). The changes in the amino acid sequence that these SNPs cause is indicated in Figure 3 and can readily be determined using the universal genetic code and the protein sequence provided in Figure 2 as a reference.

Conditions for incubating a nucleic acid molecule with a test sample vary. Incubation conditions depend on the format employed in the assay, the detection methods employed, and the type and nature of the nucleic acid molecule used in the assay. One skilled in the art will recognize that any one of the commonly available hybridization, amplification or array assay formats can readily be adapted to employ the novel fragments of the Human genome disclosed

herein. Examples of such assays can be found in Chard, T, *An Introduction to Radioimmunoassay and Related Techniques*, Elsevier Science Publishers, Amsterdam, The Netherlands (1986); Bullock, G. R. *et al.*, *Techniques in Immunocytochemistry*, Academic Press, Orlando, FL Vol. 1 (1982), Vol. 2 (1983), Vol. 3 (1985); Tijssen, P., *Practice and Theory of Enzyme Immunoassays: Laboratory Techniques in Biochemistry and Molecular Biology*, Elsevier Science Publishers, Amsterdam, The Netherlands (1985).

The test samples of the present invention include cells, protein or membrane extracts of cells. The test sample used in the above-described method will vary based on the assay format, nature of the detection method and the tissues, cells or extracts used as the sample to be assayed.

10 Methods for preparing nucleic acid extracts or of cells are well known in the art and can be readily be adapted in order to obtain a sample that is compatible with the system utilized.

In another embodiment of the present invention, kits are provided which contain the necessary reagents to carry out the assays of the present invention.

Specifically, the invention provides a compartmentalized kit to receive, in close  
15 confinement, one or more containers which comprises: (a) a first container comprising one of the nucleic acid molecules that can bind to a fragment of the Human genome disclosed herein; and (b) one or more other containers comprising one or more of the following: wash reagents, reagents capable of detecting presence of a bound nucleic acid.

In detail, a compartmentalized kit includes any kit in which reagents are contained in  
20 separate containers. Such containers include small glass containers, plastic containers, strips of plastic, glass or paper, or arraying material such as silica. Such containers allows one to efficiently transfer reagents from one compartment to another compartment such that the samples and reagents are not cross-contaminated, and the agents or solutions of each container can be added in a quantitative fashion from one compartment to another. Such containers will  
25 include a container which will accept the test sample, a container which contains the nucleic acid probe, containers which contain wash reagents (such as phosphate buffered saline, Tris-buffers, etc.), and containers which contain the reagents used to detect the bound probe. One skilled in the art will readily recognize that the previously unidentified kinase gene of the present invention can be routinely identified using the sequence information disclosed herein can be readily  
30 incorporated into one of the established kit formats which are well known in the art, particularly expression arrays.

### Vectors/host cells

The invention also provides vectors containing the nucleic acid molecules described herein. The term "vector" refers to a vehicle, preferably a nucleic acid molecule, which can transport the nucleic acid molecules. When the vector is a nucleic acid molecule, the nucleic acid molecules are covalently linked to the vector nucleic acid. With this aspect of the invention, the vector includes a plasmid, single or double stranded phage, a single or double stranded RNA or DNA viral vector, or artificial chromosome, such as a BAC, PAC, YAC, OR MAC.

A vector can be maintained in the host cell as an extrachromosomal element where it replicates and produces additional copies of the nucleic acid molecules. Alternatively, the vector may integrate into the host cell genome and produce additional copies of the nucleic acid molecules when the host cell replicates.

The invention provides vectors for the maintenance (cloning vectors) or vectors for expression (expression vectors) of the nucleic acid molecules. The vectors can function in prokaryotic or eukaryotic cells or in both (shuttle vectors).

Expression vectors contain cis-acting regulatory regions that are operably linked in the vector to the nucleic acid molecules such that transcription of the nucleic acid molecules is allowed in a host cell. The nucleic acid molecules can be introduced into the host cell with a separate nucleic acid molecule capable of affecting transcription. Thus, the second nucleic acid molecule may provide a trans-acting factor interacting with the cis-regulatory control region to allow transcription of the nucleic acid molecules from the vector. Alternatively, a trans-acting factor may be supplied by the host cell. Finally, a trans-acting factor can be produced from the vector itself. It is understood, however, that in some embodiments, transcription and/or translation of the nucleic acid molecules can occur in a cell-free system.

The regulatory sequence to which the nucleic acid molecules described herein can be operably linked include promoters for directing mRNA transcription. These include, but are not limited to, the left promoter from bacteriophage  $\lambda$ , the lac, TRP, and TAC promoters from *E. coli*, the early and late promoters from SV40, the CMV immediate early promoter, the adenovirus early and late promoters, and retrovirus long-terminal repeats.

In addition to control regions that promote transcription, expression vectors may also include regions that modulate transcription, such as repressor binding sites and enhancers. Examples include the SV40 enhancer, the cytomegalovirus immediate early enhancer, polyoma enhancer, adenovirus enhancers, and retrovirus LTR enhancers.

In addition to containing sites for transcription initiation and control, expression vectors can also contain sequences necessary for transcription termination and, in the transcribed region a ribosome binding site for translation. Other regulatory control elements for expression include initiation and termination codons as well as polyadenylation signals. The person of ordinary skill in the art would be aware of the numerous regulatory sequences that are useful in expression vectors. Such regulatory sequences are described, for example, in Sambrook *et al.*, *Molecular Cloning: A Laboratory Manual*. 2nd. ed., Cold Spring Harbor Laboratory Press, Cold Spring Harbor, NY, (1989).

A variety of expression vectors can be used to express a nucleic acid molecule. Such vectors include chromosomal, episomal, and virus-derived vectors, for example vectors derived from bacterial plasmids, from bacteriophage, from yeast episomes, from yeast chromosomal elements, including yeast artificial chromosomes, from viruses such as baculoviruses, papovaviruses such as SV40, Vaccinia viruses, adenoviruses, poxviruses, pseudorabies viruses, and retroviruses. Vectors may also be derived from combinations of these sources such as those derived from plasmid and bacteriophage genetic elements, e.g. cosmids and phagemids. Appropriate cloning and expression vectors for prokaryotic and eukaryotic hosts are described in Sambrook *et al.*, *Molecular Cloning: A Laboratory Manual*. 2nd. ed., Cold Spring Harbor Laboratory Press, Cold Spring Harbor, NY, (1989).

The regulatory sequence may provide constitutive expression in one or more host cells (i.e. tissue specific) or may provide for inducible expression in one or more cell types such as by temperature, nutrient additive, or exogenous factor such as a hormone or other ligand. A variety of vectors providing for constitutive and inducible expression in prokaryotic and eukaryotic hosts are well known to those of ordinary skill in the art.

The nucleic acid molecules can be inserted into the vector nucleic acid by well-known methodology. Generally, the DNA sequence that will ultimately be expressed is joined to an expression vector by cleaving the DNA sequence and the expression vector with one or more restriction enzymes and then ligating the fragments together. Procedures for restriction enzyme digestion and ligation are well known to those of ordinary skill in the art.

The vector containing the appropriate nucleic acid molecule can be introduced into an appropriate host cell for propagation or expression using well-known techniques. Bacterial cells include, but are not limited to, *E. coli*, *Streptomyces*, and *Salmonella typhimurium*. Eukaryotic cells include, but are not limited to, yeast, insect cells such as *Drosophila*, animal cells such as COS and CHO cells, and plant cells.



As described herein, it may be desirable to express the peptide as a fusion protein. Accordingly, the invention provides fusion vectors that allow for the production of the peptides. Fusion vectors can increase the expression of a recombinant protein, increase the solubility of the recombinant protein, and aid in the purification of the protein by acting for example as a ligand for affinity purification. A proteolytic cleavage site may be introduced at the junction of the fusion moiety so that the desired peptide can ultimately be separated from the fusion moiety. Proteolytic enzymes include, but are not limited to, factor Xa, thrombin, and enterokinase. Typical fusion expression vectors include pGEX (Smith *et al.*, *Gene* 67:31-40 (1988)), pMAL (New England Biolabs, Beverly, MA) and pRIT5 (Pharmacia, Piscataway, NJ) which fuse glutathione S-transferase (GST), maltose E binding protein, or protein A, respectively, to the target recombinant protein. Examples of suitable inducible non-fusion *E. coli* expression vectors include pTrc (Amann *et al.*, *Gene* 69:301-315 (1988)) and pET 11d (Studier *et al.*, *Gene Expression Technology: Methods in Enzymology* 185:60-89 (1990)).

Recombinant protein expression can be maximized in host bacteria by providing a genetic background wherein the host cell has an impaired capacity to proteolytically cleave the recombinant protein. (Gottesman, S., *Gene Expression Technology: Methods in Enzymology* 185, Academic Press, San Diego, California (1990) 119-128). Alternatively, the sequence of the nucleic acid molecule of interest can be altered to provide preferential codon usage for a specific host cell, for example *E. coli*. (Wada *et al.*, *Nucleic Acids Res.* 20:2111-2118 (1992)).

The nucleic acid molecules can also be expressed by expression vectors that are operative in yeast. Examples of vectors for expression in yeast e.g., *S. cerevisiae* include pYepSec1 (Baldari, *et al.*, *EMBO J.* 6:229-234 (1987)), pMFa (Kurjan *et al.*, *Cell* 30:933-943(1982)), pJRY88 (Schultz *et al.*, *Gene* 54:113-123 (1987)), and pYES2 (Invitrogen Corporation, San Diego, CA).

The nucleic acid molecules can also be expressed in insect cells using, for example, baculovirus expression vectors. Baculovirus vectors available for expression of proteins in cultured insect cells (e.g., Sf9 cells) include the pAc series (Smith *et al.*, *Mol. Cell Biol.* 3:2156-2165 (1983)) and the pVL series (Lucklow *et al.*, *Virology* 170:31-39 (1989)).

In certain embodiments of the invention, the nucleic acid molecules described herein are expressed in mammalian cells using mammalian expression vectors. Examples of mammalian expression vectors include pCDM8 (Seed, B. *Nature* 329:840(1987)) and pMT2PC (Kaufman *et al.*, *EMBO J.* 6:187-195 (1987)).

The expression vectors listed herein are provided by way of example only of the well-known vectors available to those of ordinary skill in the art that would be useful to express the

nucleic acid molecules. The person of ordinary skill in the art would be aware of other vectors suitable for maintenance propagation or expression of the nucleic acid molecules described herein. These are found for example in Sambrook, J., Fritsh, E. F., and Maniatis, T. *Molecular Cloning: A Laboratory Manual. 2nd, ed., Cold Spring Harbor Laboratory, Cold Spring Harbor Laboratory*  
5 Press, Cold Spring Harbor, NY, 1989.

The invention also encompasses vectors in which the nucleic acid sequences described herein are cloned into the vector in reverse orientation, but operably linked to a regulatory sequence that permits transcription of antisense RNA. Thus, an antisense transcript can be produced to all, or to a portion, of the nucleic acid molecule sequences described herein, including both coding and  
10 non-coding regions. Expression of this antisense RNA is subject to each of the parameters described above in relation to expression of the sense RNA (regulatory sequences, constitutive or inducible expression, tissue-specific expression).

The invention also relates to recombinant host cells containing the vectors described herein. Host cells therefore include prokaryotic cells, lower eukaryotic cells such as yeast, other eukaryotic  
15 cells such as insect cells, and higher eukaryotic cells such as mammalian cells.

The recombinant host cells are prepared by introducing the vector constructs described herein into the cells by techniques readily available to the person of ordinary skill in the art. These include, but are not limited to, calcium phosphate transfection, DEAE-dextran-mediated transfection, cationic lipid-mediated transfection, electroporation, transduction, infection,  
20 lipofection, and other techniques such as those found in Sambrook, *et al. (Molecular Cloning: A Laboratory Manual. 2nd, ed., Cold Spring Harbor Laboratory, Cold Spring Harbor Laboratory Press, Cold Spring Harbor, NY, 1989).*

Host cells can contain more than one vector. Thus, different nucleotide sequences can be introduced on different vectors of the same cell. Similarly, the nucleic acid molecules can be  
25 introduced either alone or with other nucleic acid molecules that are not related to the nucleic acid molecules such as those providing trans-acting factors for expression vectors. When more than one vector is introduced into a cell, the vectors can be introduced independently, co-introduced or joined to the nucleic acid molecule vector.

In the case of bacteriophage and viral vectors, these can be introduced into cells as packaged  
30 or encapsulated virus by standard procedures for infection and transduction. Viral vectors can be replication-competent or replication-defective. In the case in which viral replication is defective, replication will occur in host cells providing functions that complement the defects.

Vectors generally include selectable markers that enable the selection of the subpopulation of cells that contain the recombinant vector constructs. The marker can be contained in the same vector that contains the nucleic acid molecules described herein or may be on a separate vector. Markers include tetracycline or ampicillin-resistance genes for prokaryotic host cells and  
5 dihydrofolate reductase or neomycin resistance for eukaryotic host cells. However, any marker that provides selection for a phenotypic trait will be effective.

While the mature proteins can be produced in bacteria, yeast, mammalian cells, and other cells under the control of the appropriate regulatory sequences, cell- free transcription and translation systems can also be used to produce these proteins using RNA derived from the DNA  
10 constructs described herein.

Where secretion of the peptide is desired, which is difficult to achieve with multi-transmembrane domain containing proteins such as kinases, appropriate secretion signals are incorporated into the vector. The signal sequence can be endogenous to the peptides or heterologous to these peptides.

15 Where the peptide is not secreted into the medium, which is typically the case with kinases, the protein can be isolated from the host cell by standard disruption procedures, including freeze thaw, sonication, mechanical disruption, use of lysing agents and the like. The peptide can then be recovered and purified by well-known purification methods including ammonium sulfate precipitation, acid extraction, anion or cationic exchange chromatography, phosphocellulose  
20 chromatography, hydrophobic-interaction chromatography, affinity chromatography, hydroxylapatite chromatography, lectin chromatography, or high performance liquid chromatography.

It is also understood that depending upon the host cell in recombinant production of the peptides described herein, the peptides can have various glycosylation patterns, depending upon the  
25 cell, or maybe non-glycosylated as when produced in bacteria. In addition, the peptides may include an initial modified methionine in some cases as a result of a host-mediated process.

#### Uses of vectors and host cells

The recombinant host cells expressing the peptides described herein have a variety of uses.  
30 First, the cells are useful for producing a kinase protein or peptide that can be further purified to produce desired amounts of kinase protein or fragments. Thus, host cells containing expression vectors are useful for peptide production.

Host cells are also useful for conducting cell-based assays involving the kinase protein or kinase protein fragments, such as those described above as well as other formats known in the art. Thus, a recombinant host cell expressing a native kinase protein is useful for assaying compounds that stimulate or inhibit kinase protein function.

5        Host cells are also useful for identifying kinase protein mutants in which these functions are affected. If the mutants naturally occur and give rise to a pathology, host cells containing the mutations are useful to assay compounds that have a desired effect on the mutant kinase protein (for example, stimulating or inhibiting function) which may not be indicated by their effect on the native kinase protein.

10        Genetically engineered host cells can be further used to produce non-human transgenic animals. A transgenic animal is preferably a mammal, for example a rodent, such as a rat or mouse, in which one or more of the cells of the animal include a transgene. A transgene is exogenous DNA which is integrated into the genome of a cell from which a transgenic animal develops and which remains in the genome of the mature animal in one or more cell types or tissues of the transgenic  
15        animal. These animals are useful for studying the function of a kinase protein and identifying and evaluating modulators of kinase protein activity. Other examples of transgenic animals include non-human primates, sheep, dogs, cows, goats, chickens, and amphibians.

A transgenic animal can be produced by introducing nucleic acid into the male pronuclei of a fertilized oocyte, e.g., by microinjection, retroviral infection, and allowing the oocyte to develop  
20        in a pseudopregnant female foster animal. Any of the kinase protein nucleotide sequences can be introduced as a transgene into the genome of a non-human animal, such as a mouse.

Any of the regulatory or other sequences useful in expression vectors can form part of the transgenic sequence. This includes intronic sequences and polyadenylation signals, if not already included. A tissue-specific regulatory sequence(s) can be operably linked to the transgene to direct  
25        expression of the kinase protein to particular cells.

Methods for generating transgenic animals via embryo manipulation and microinjection, particularly animals such as mice, have become conventional in the art and are described, for example, in U.S. Patent Nos. 4,736,866 and 4,870,009, both by Leder *et al.*, U.S. Patent No. 4,873,191 by Wagner *et al.* and in Hogan, B., *Manipulating the Mouse Embryo*, (Cold Spring  
30        Harbor Laboratory Press, Cold Spring Harbor, N.Y., 1986). Similar methods are used for production of other transgenic animals. A transgenic founder animal can be identified based upon the presence of the transgene in its genome and/or expression of transgenic mRNA in tissues or cells of the animals. A transgenic founder animal can then be used to breed additional animals

carrying the transgene. Moreover, transgenic animals carrying a transgene can further be bred to other transgenic animals carrying other transgenes. A transgenic animal also includes animals in which the entire animal or tissues in the animal have been produced using the homologously recombinant host cells described herein.

5 In another embodiment, transgenic non-human animals can be produced which contain selected systems that allow for regulated expression of the transgene. One example of such a system is the *cre/loxP* recombinase system of bacteriophage P1. For a description of the *cre/loxP* recombinase system, see, e.g., Lakso *et al. PNAS* 89:6232-6236 (1992). Another example of a recombinase system is the FLP recombinase system of *S. cerevisiae* (O'Gorman *et al. Science* 10 251:1351-1355 (1991). If a *cre/loxP* recombinase system is used to regulate expression of the transgene, animals containing transgenes encoding both the *Cre* recombinase and a selected protein is required. Such animals can be provided through the construction of "double" transgenic animals, e.g., by mating two transgenic animals, one containing a transgene encoding a selected protein and the other containing a transgene encoding a recombinase.

15 Clones of the non-human transgenic animals described herein can also be produced according to the methods described in Wilmut, I. *et al. Nature* 385:810-813 (1997) and PCT International Publication Nos. WO 97/07668 and WO 97/07669. In brief, a cell, e.g., a somatic cell, from the transgenic animal can be isolated and induced to exit the growth cycle and enter G<sub>0</sub> phase. The quiescent cell can then be fused, e.g., through the use of electrical pulses, to an enucleated 20 oocyte from an animal of the same species from which the quiescent cell is isolated. The reconstructed oocyte is then cultured such that it develops to morula or blastocyst and then transferred to pseudopregnant female foster animal. The offspring born of this female foster animal will be a clone of the animal from which the cell, e.g., the somatic cell, is isolated.

Transgenic animals containing recombinant cells that express the peptides described herein 25 are useful to conduct the assays described herein in an *in vivo* context. Accordingly, the various physiological factors that are present *in vivo* and that could effect substrate binding, kinase protein activation, and signal transduction, may not be evident from *in vitro* cell-free or cell-based assays. Accordingly, it is useful to provide non-human transgenic animals to assay *in vivo* kinase protein function, including substrate interaction, the effect of specific mutant kinase proteins on kinase 30 protein function and substrate interaction, and the effect of chimeric kinase proteins. It is also possible to assess the effect of null mutations, that is, mutations that substantially or completely eliminate one or more kinase protein functions.

All publications and patents mentioned in the above specification are herein incorporated by reference. Various modifications and variations of the described method and system of the invention will be apparent to those skilled in the art without departing from the scope and spirit of the invention. Although the invention has been described in connection with specific  
5 preferred embodiments, it should be understood that the invention as claimed should not be unduly limited to such specific embodiments. Indeed, various modifications of the above-described modes for carrying out the invention which are obvious to those skilled in the field of molecular biology or related fields are intended to be within the scope of the following claims.

## Claims

That which is claimed is:

1. An isolated peptide consisting of an amino acid sequence selected from the group consisting of:
  - (a) an amino acid sequence shown in SEQ ID NO:2;
  - (b) an amino acid sequence of an allelic variant of an amino acid sequence shown in SEQ ID NO:2, wherein said allelic variant is encoded by a nucleic acid molecule that hybridizes under stringent conditions to the opposite strand of a nucleic acid molecule shown in SEQ ID NOS:1 or 3;
  - (c) an amino acid sequence of an ortholog of an amino acid sequence shown in SEQ ID NO:2, wherein said ortholog is encoded by a nucleic acid molecule that hybridizes under stringent conditions to the opposite strand of a nucleic acid molecule shown in SEQ ID NOS:1 or 3; and
  - (d) a fragment of an amino acid sequence shown in SEQ ID NO:2, wherein said fragment comprises at least 10 contiguous amino acids.
2. An isolated peptide comprising an amino acid sequence selected from the group consisting of:
  - (a) an amino acid sequence shown in SEQ ID NO:2;
  - (b) an amino acid sequence of an allelic variant of an amino acid sequence shown in SEQ ID NO:2, wherein said allelic variant is encoded by a nucleic acid molecule that hybridizes under stringent conditions to the opposite strand of a nucleic acid molecule shown in SEQ ID NOS:1 or 3;
  - (c) an amino acid sequence of an ortholog of an amino acid sequence shown in SEQ ID NO:2, wherein said ortholog is encoded by a nucleic acid molecule that hybridizes under stringent conditions to the opposite strand of a nucleic acid molecule shown in SEQ ID NOS:1 or 3; and
  - (d) a fragment of an amino acid sequence shown in SEQ ID NO:2, wherein said fragment comprises at least 10 contiguous amino acids.
3. An isolated antibody that selectively binds to a peptide of claim 2.

4. An isolated nucleic acid molecule consisting of a nucleotide sequence selected from the group consisting of:

- (a) a nucleotide sequence that encodes an amino acid sequence shown in SEQ ID NO:2;
- (b) a nucleotide sequence that encodes of an allelic variant of an amino acid sequence shown in SEQ ID NO:2, wherein said nucleotide sequence hybridizes under stringent conditions to the opposite strand of a nucleic acid molecule shown in SEQ ID NOS:1 or 3;
- (c) a nucleotide sequence that encodes an ortholog of an amino acid sequence shown in SEQ ID NO:2, wherein said nucleotide sequence hybridizes under stringent conditions to the opposite strand of a nucleic acid molecule shown in SEQ ID NOS:1 or 3;
- (d) a nucleotide sequence that encodes a fragment of an amino acid sequence shown in SEQ ID NO:2, wherein said fragment comprises at least 10 contiguous amino acids; and
- (e) a nucleotide sequence that is the complement of a nucleotide sequence of (a)-(d).

5. An isolated nucleic acid molecule comprising a nucleotide sequence selected from the group consisting of:

- (a) a nucleotide sequence that encodes an amino acid sequence shown in SEQ ID NO:2;
- (b) a nucleotide sequence that encodes of an allelic variant of an amino acid sequence shown in SEQ ID NO:2, wherein said nucleotide sequence hybridizes under stringent conditions to the opposite strand of a nucleic acid molecule shown in SEQ ID NOS:1 or 3;
- (c) a nucleotide sequence that encodes an ortholog of an amino acid sequence shown in SEQ ID NO:2, wherein said nucleotide sequence hybridizes under stringent conditions to the opposite strand of a nucleic acid molecule shown in SEQ ID NOS:1 or 3;
- (d) a nucleotide sequence that encodes a fragment of an amino acid sequence shown in SEQ ID NO:2, wherein said fragment comprises at least 10 contiguous amino acids; and
- (e) a nucleotide sequence that is the complement of a nucleotide sequence of (a)-(d).

6. A gene chip comprising a nucleic acid molecule of claim 5.

7. A transgenic non-human animal comprising a nucleic acid molecule of claim 5.



8. A nucleic acid vector comprising a nucleic acid molecule of claim 5.
9. A host cell containing the vector of claim 8.
10. A method for producing any of the peptides of claim 1 comprising introducing a nucleotide sequence encoding any of the amino acid sequences in (a)-(d) into a host cell, and culturing the host cell under conditions in which the peptides are expressed from the nucleotide sequence.
11. A method for producing any of the peptides of claim 2 comprising introducing a nucleotide sequence encoding any of the amino acid sequences in (a)-(d) into a host cell, and culturing the host cell under conditions in which the peptides are expressed from the nucleotide sequence.
12. A method for detecting the presence of any of the peptides of claim 2 in a sample, said method comprising contacting said sample with a detection agent that specifically allows detection of the presence of the peptide in the sample and then detecting the presence of the peptide.
13. A method for detecting the presence of a nucleic acid molecule of claim 5 in a sample, said method comprising contacting the sample with an oligonucleotide that hybridizes to said nucleic acid molecule under stringent conditions and determining whether the oligonucleotide binds to said nucleic acid molecule in the sample.
14. A method for identifying a modulator of a peptide of claim 2, said method comprising contacting said peptide with an agent and determining if said agent has modulated the function or activity of said peptide.
15. The method of claim 14, wherein said agent is administered to a host cell comprising an expression vector that expresses said peptide.

16. A method for identifying an agent that binds to any of the peptides of claim 2, said method comprising contacting the peptide with an agent and assaying the contacted mixture to determine whether a complex is formed with the agent bound to the peptide.
17. A pharmaceutical composition comprising an agent identified by the method of claim 16 and a pharmaceutically acceptable carrier therefor.
18. A method for treating a disease or condition mediated by a human kinase protein, said method comprising administering to a patient a pharmaceutically effective amount of an agent identified by the method of claim 16.
19. A method for identifying a modulator of the expression of a peptide of claim 2, said method comprising contacting a cell expressing said peptide with an agent, and determining if said agent has modulated the expression of said peptide.
20. An isolated human kinase peptide having an amino acid sequence that shares at least 70% homology with an amino acid sequence shown in SEQ ID NO:2.
21. A peptide according to claim 20 that shares at least 90 percent homology with an amino acid sequence shown in SEQ ID NO:2.
22. An isolated nucleic acid molecule encoding a human kinase peptide, said nucleic acid molecule sharing at least 80 percent homology with a nucleic acid molecule shown in SEQ ID NOS:1 or 3.
23. A nucleic acid molecule according to claim 22 that shares at least 90 percent homology with a nucleic acid molecule shown in SEQ ID NOS:1 or 3.

1 CAGCACGAGG AACTCCTTCT GATCACCTGG CCAGCTGAGG TCAGAGTGGG  
 51 AGAGGCAGTG GTTCCATTGA AGGAGTACTC CTAAGTCA GAAGCCTGGG  
 101 CGGTACAGGAT GGGGTGCTGT CGCTTGGGCT GCGGGGGTG TTCAATTGCC  
 151 CACAGTGTAT CTCAGGGTCT CACCAACCAT CCAAGCATGG TAGGCTGTGG  
 201 CTGGCACCCA GGGTTGTGTG GCTGGGGAGG TGCTCTCCAC AGTTCCTCC  
 251 CTGCCCTCCC AGGGCCCCCA TCCATGCAGG TAACCATCGA GGATGTGCAG  
 301 GCACAGACAG GCGGAACGGC CCAATTCGAG GCTATCATTTG AGGGCGACCC  
 351 ACAGCCCTCG GTGACCTGGT ACAAGGACAG CGTCCAGCTG GTGACAGCA  
 401 CCCGGCTTAG CCAGCAGCAA GAAGGCACCA CATACTCCCT GGTGCTGAGG  
 451 CATGTGGCCT CGAAGGATGC CGGCGTTTAC ACCTGCCTGG CCCAAACAC  
 501 TGGTGGCCAG GTGCTCTGCA AGGCAGAGCT GCTGGTGCTT GGGGGGACA  
 551 ATGAGCCGGA CTCAGAGAAG CAAAGCCACC GGAGGAAGCT GCACTCCTTC  
 601 TATGAGTCA AGGAGGAGAT TGGAGGGGC GTGTTTGGCT TCGTAAAAAG  
 651 AGTGACAGAC AAAGGAAACA AGATCTTGTG CGTGCCCAAG TTCATCCCCC  
 701 TACGGAGCAG AACTCGGGCC CAGGCATACA GGGAGCGAGA CATCCTGGCC  
 751 GCGCTGAGCC ACCCGCTGGT CACGGGGCTG CTGGACCAGT TTGAGACCCG  
 801 CAAGACCCTC ATCCTCATCC TGGAGCTGTG CTCATCCGAG GAGCTGCTGG  
 851 ACCGCTGTGA CAGGAAGGGC GTGGTGACGG AGGCCGAGGT CAAGGTCTAC  
 901 ATCCAGCAGC TGGTGGAGGG GCTGCACTAC CTGCACAGCC ATGGCGTCT  
 951 CCACCTGGAG ATAAAGCCCT CTAACATCCT GATGGTGCAT CCGCCCGGG  
 1001 AAGACATTAA AATCTGCGAC TTTGGCTTTG CCCAGAACAT CACCCAGCA  
 1051 GAGCTGCAGT TCAGCCAGTA CGGCTCCCTT GAGTTCGTCT CCCCCGAGAT  
 1101 CATCCAGCAG AACCTGTGA ACGAAGCCTC CGACATTTGG CCGATGGGTG  
 1151 TCATCTCCTA CCTCAGCCTG ACCTGCTCAT CCCCATTGCG CGGCAGAGT  
 1201 GACCGTGCCA CCTCCTGAA CGTCTGGAG GGGCGCGTGT CATGGAGCAG  
 1251 CCCCATTGGT GCCCACCCTA GCGAAGACGC CAAAGACTTC ATCAAGGCTA  
 1301 CGCTGCAGAG AGCCCTCAG GCCCGGCTTA GTGCGGCCA GTGCTCTCTC  
 1351 CACCCCTGGT TCCTGAAATC CATGCTGCG GAGGAGGCCC ACTTCATCAA  
 1401 CACCAAGCAG CTCAGATTCC TCCTGGCCCG AAGTCGCTGG CAGCGTTCCC  
 1451 TGATGAGCTA CAAGTCCATC CTGGTGATGC GCTCCATCCC TGAGCTGCTG  
 1501 CGGGGCCAC CCGACAGCCC CTCCTCGCG GTAGCCCGGC ACCTCTGCAG  
 1551 GGAACCTGGT GGCTCCTCCA GTTCTCTCTC CTCCTCTGAC AACGAGCTCG  
 1601 CCCCATTGCG CCGGGCTAAG TCACGTCCAC CCTCCCGGT GACACACTCA  
 1651 CCACCTGTGC ACCCCCGGGG CTTCTGCGG CCTCGGCCA GCCTGCCTGA  
 1701 GGAAGCCGAG GCCAGTGAGC GCTCCACCGA GGGCCAGCT GCGCTGCAT  
 1751 CTCCCGAGGG TGCCGGGCCA CCGGCGGCC AGGGCTGCGT GCGCCGAC  
 1801 AGCGTATCC GCAGCCTGTT TACACACAG GCGGGTGAGA GCGCTGAGCA  
 1851 CCGGGCCCTG GCCCGGGGA GCAGGCGGCA CCGGCCCGG CCGCGGAC  
 1901 TGCTGAAGGG CGGCTACATT GCGGGGGCGC TGCCAGGCTT CCGCGGCCA  
 1951 CTGATGGAGC ACCGCGTGCT GGAGGAGGAG GCGGCCAGG AGGAGCAGGC  
 2001 CACCCCTCTG GCCAAAGCCC CCTCATTCGA GACTGCCTTC CCGCTGCCTG  
 2051 CCTCTGGCAC CCACCTGGCC CTTGGCCACA GCGACTCCCT GGAACATGAC  
 2101 TCTCCGAGCA CCCCCGCCC CTCCTCGGAG GCCTGCGGTG AGGCACAGCG  
 2151 ACTGCCCTTA CCCCCCTCCG GGGGGGGCCC TATCAGGGAC ATGGGGCACC  
 2201 CTCAGGGCTC CAAGCAGCTT CCATCCACTG GTGGCCACCC AGGCATGCT  
 2251 CAGCCAGAGA GGCCATCCCC GGACAGCCCT TGGGGGAGC CAGCCCTTT  
 2301 CTGCCACCCC AAGCAGGGTT CTGCCCCCA GGAGGGCTGC AGCCCCCACC  
 2351 CAGCAGTTGC CCCATGCCCT CTTGGCTCCT TCCCTCCAGG ATCTTGCAAA  
 2401 GAGGCCCCCT TAGTACCCTC AAGCCCTTC TTGGGACAGC CCCAGGCACC  
 2451 CCCTGCCCTT GCCAAAGCAA GCGCCCATTT GGAATCTAAG ATGGGGCTG  
 2501 GAGACATCTC TCTTCTGGG AGGCCAAAC CCGGCCCTG CAGTTCCCA  
 2551 GGGTCAGCTT CCCAGGCGAG CTCTTCCCA GTGAGCTCCC TCAGGGTGGG  
 2601 CTCTCCAG GTGGGCACAG AGCCTGGCCC CTCCCTGGAT GCGGAGGGCT  
 2651 GGACCCAGGA GGCTGAGGAT CTGTCCGACT CCACACCCAC CTTGCAGCGG  
 2701 CCTCAGGAAC AGGTGACCAT GCGCAAGTTC TCCCTGGGTG GTCGCGGGG  
 2751 CTACGCAGGC GTGGCTGGCT ATGGCACCTT TGCTTTGGT GGAGATGCAG  
 2801 GGGGCATGCT GGGGCAGGG CCCATGTGG CCAGGATAGC CTGGGCTGTG  
 2851 TCCAGTCCG AGGAGGAGGA GCAGGAGGAG GCCAGGGCTG AGTCCCAGTC  
 2901 GGAGGAGCAG CAGGAGGCCA GGGCTGAGAG CCCACTGCCC CAGGTACGTG  
 2951 CAAGGCTGT GCCTGAGGTC GGCAGGGCTC CCACAGGAG CTCTCCAGAG  
 3001 CCCACCCCAT GGGAGGACAT CCGGCAGGTC TCCCTGGTGC AGATCCGGGA  
 3051 CCTGTACGTT GATGCGGAGG CGGCCGACAC AATATCCCTG GACATTTCCG  
 3101 AGGTGGACCC GCGCTACCTC AACCTCTCAG ACCTGTACGA TATCAAGTAC  
 3151 CTCCTCATCG AGTTTATGAT CTTAGGAAA GTCCCAAGT CCGCTCAGCC  
 3201 AGAGCCGCCC TCCCCATGG CTGAGGAGGA GCTGGCCGAG TTCCCGGAGC  
 3251 CCACGTGGCC CTGGCCAGGT GAACTGGGCC CCCACGCAGG CCTGGAGATC  
 3301 ACAGAGGAGT CAGAGGATGT GGACGCGCTG CTGGCAGAGG CTGCCGTGGG  
 3351 CAGGAAGCGC AAGTGGTCTT CGCCGTACAG CAGCCTCTTC CACTTCCCTG  
 3401 GGAGGCACCT GCGCTGGAT GAGCCTGCAG AGCTGGGGCT GCGTGAGAGA  
 3451 GTGAAGGCTT CCGTGGAGCA CATCTCCCG ATCCTGAAG GCAGGCCGGA  
 3501 AGGTCTGGAG AAGGAGGGGC CCCCAGGAA GAAGCCAGGC CTTGCTTCTC  
 3551 TCCGGCTCTC AGGTCTGAAG AGCTGGGACC GAGCGCCGAC ATTCTAAGG  
 3601 GAGCTCTCAG ATGAGACTGT GGTCTGGGC CAGTCAGTGA CACTGGCCTG

FIGURE 1A

SUBSTITUTE SHEET (RULE 26)

2/17

```

3651 CCAGGTGTCA GCCCAGCCAG CTGCCCAGGC CACCTGGAGC AAAGACGGAG
3701 CCCCCTTGGG GAGCAGCAGC CGTGTCTTCA TCTCTGCCAC CCTCAAGAAC
3751 TTCCAGCTTC TGACCATCCT GGTGGTGGTG GCTGAGGACC TGGGTGTGTA
3801 CACCTGCAGC GTGAGCAATG CGCTGGGGAC AGTGACCACC ACGGGCGTCC
3851 TCCGGAAGGC AGAGCGCCCC TCATCTTCGC CATGCCCGGA TATCGGGGAG
3901 GTGTACGCGG ATGGGGTGCT GCTGGTCTGG AAGCCCGTGG AATCCTACGG
3951 CCCTGTGACC TACATTGTGC AGTGCAGCCT AGAAGGCGGC AGCTGGACCA
4001 CACTGGCCTC CGACATCTTT GACTGCTGCT ACCTGACCAG CAAGCTCTCC
4051 CGGGGTGGCA CCTACACCTT CCGCACGGCA TGTGTACGCA AGGCAGGAAT
4101 GGGTCCCTAC AGCAGCCCTT CGGAGCAAGT CCTCCTGGGA GGGCCAGCC
4151 ACCTGGCCCTC TGAGGAGGAG AGCCAGGGGC GGTACAGCCA ACCCTTGCCC
4201 AGCACAAGA CCTTCGCATT CCAGACACAG ATCCAGAGGG GCCGCTTCAG
4251 CGTGGTGGCG CAATGCTGGG AGAAGGCCAG CGGGCGGGCG CTGGCCGCCA
4301 AGATCATCCC CTACCACCCC AAGGACAAGA CAGCAGTGCT GCGCGAATAC
4351 GAGGCCCTCA AGGGCCTGCG CCACCCGCAC CTGGCCAGC TGCACGCAGC
4401 CTACCTCAGC CCCCAGCACC TGGTGTCTAT CTTGGAGCTG TGCTCTGGGC
4451 CCGAGCTGCT CCCCTGCTG GCGGAGAGGG CCTCCTACTC AGAATCTGAG
4501 GTGAAGGACT ACCTGTGGCA GATGTTGAGT GCCACCCAGT ACCTGCACAA
4551 CCAGCACATC CTGCACCTGG ACCTGAGGTC CGAGAACATG ATCATCACCG
4601 AATACAACCT GCTCAAGGTC GTGGACCTGG GCAATGCACA GAGCCTCAGC
4651 CAGGAGAAGG TGCTGCCCTC AGACAAGTTC AAGGACTACC TAGAGACCAT
4701 GGCTCCAGAG CTCTTGGAGG GCCAGGGGGC TGTTCACAG ACAGACATCT
4751 GGGCCATCGG TGTGACAGCC TTCATCATGC TGAGCGCCGA GTACCCGGTG
4801 AGCAGCGAGG GTGCACGCGA CCTGCAGAGA GGAAGTGCAG AGGGGCTGGT
4851 CCGGCTGAGC CGCTGCTACG CGGGGCTGTC CGGGGGCGCC GTGGCCTTCC
4901 TGGCGAGCAC TCTGTGCGCC CAGCCCTGGG CCGGGCCCTG CGCGTCCAGC
4951 TGCCTGCAGT GCCCGTGGCT AACAGAGGAG GGCCCGGCTT GTTCGCGGCC
5001 CGCGCCCGTG ACCTTCCCTA CCGCGCGGCT GCGCGTCTTC GTGCGCAATC
5051 GCGAGAAGAG ACGCGCGCTG CTGTACAAGA GGCACAACCT GGCCAGGTG
5101 CGCTGAGGGT CGCCCGGGCC ACACCTTGG TCTCCCGCTG GGGGGTCTCT
5151 GCAGACGCGC CAATAAAAC GCACAGCCGG GCGAGAAAAA AAAAAAA
5201 AAAAAA . (SEQ ID NO:1)

```

#### FEATURES:

Start: 109

Stop: 5104

#### Homologous proteins:

##### Top BLAST Hits:

	Score	E
gi 7242949 dbj BAA92535.1  (AB037718) KIAA1297 protein [Homo sa...	425	e-117
gi 8928460 sp O75962 TRIO HUMAN TRIPLE FUNCTIONAL DOMAIN PROTEI...	229	1e-58
gi 6005922 ref NP_009049.1  triple functional domain (PTPRF int...	229	1e-58
gi 3024081 sp Q15746 KMLS_HUMAN MYOSIN LIGHT CHAIN KINASE, SMOO...	206	2e-51
gi 90103 pir A41674 myosin-light-chain kinase (EC 2.7.1.117), ...	205	4e-51
gi 7239696 gb AAC18423.2  (U48959) myosin light chain kinase [H...	204	6e-51
gi 7239698 gb AAD15921.2  (AF069601) myosin light chain kinase ...	204	6e-51
gi 1103677 emb CAA62378.1  (X90870) myosin-light-chain kinase [...	204	6e-51
gi 3024085 sp Q28824 KMLS_BOVIN MYOSIN LIGHT CHAIN KINASE, SMOO...	203	1e-50
gi 2851405 sp P29294 KMLS_RABBIT MYOSIN LIGHT CHAIN KINASE, SMOO...	203	1e-50
gi 3982821 gb AAC83683.1  (AF081663) myosin light chain kinase ...	198	3e-49
gi 3982823 gb AAC83684.1  (AF081664) myosin light chain kinase ...	198	3e-49
gi 3982827 gb AAC83686.1  (AF081666) myosin light chain kinase ...	198	3e-49
gi 3982807 gb AAC83676.1  (AF081656) myosin light chain kinase ...	198	3e-49

##### BLAST dbEST hit:

gi|7958129 /dataset=dbest /taxon=960... 1283 0.0

#### EXPRESSION INFORMATION FOR MODULATORY USE:

##### From BLAST dbEST hit:

gi|7958129 Human Colon carcinoma

##### From PCR-based tissue screening panels:

Human Placenta  
Human Kidney  
Human Lung  
Human skeletal muscle  
Human heart  
Human fetal whole brain

FIGURE 1B

SUBSTITUTE SHEET (RULE 26)

```

1 MGCCRLGCGG CSVAHSVSQG LTNHPSMVGC GWHPGLCGWG GGLHSSLPAL
51 PGPPSMQVTI EDVQAQTGGT AQFEAIIEGD PQPSVTWYKD SVQLVDSTRL
101 SQQEGTYS LVLRHVASKD AGVYTCLAQN TGGQVLCKAE LLVLGGDNEP
51 DSEKQSHRRK LHSFYEVKEE IGRGVFGFVK RVQHKGNKIL CAAKFIPLRS
101 RTRAQAYRER DILAALSHPL VTGLLDQFET RKTILILILEL CSSEELDLRL
251 YRKGVVTEAE VKVYIQQLVE GLHYLHSHGV LHLDIKPSNI LMVHPAREDI
301 KICDFGFAQN ITPAELQFSQ YGSPEFVSPE IIQONPVSEA SDIWAMGVIS
351 YLSLTCSSPF AGESDRATLL NVLEGRVSW SPMAAHLS ED AKDFIKATLQ
401 RAPQARPSAA QCLSHPWFLK SMPAEAAHFI NTKQLKFLLA RSRWQSLMS
451 YKSILVMRSI PELLRGPPDS PSLGVARHLC RDTGSSSSSS SSSDNELAPF
501 ARAKSLPPSP VTHSPLLHPR GFLRPSASLP EEAEASERST EAPAPPASPE
551 GAGPPAAQC VPRHSVIRSL FYHQAGESPE HGALAPGSRR HPARRRHLLK
601 GGYIAGALPG LREPLMEHRV LEEEAAREEQ ATLLAKAPSF ETALRLPASG
651 THLAPGSHS LEHDSPTPR PSSEACGEAQ RLPSAPSGGA PIRDMGHPOG
701 SKQLPSTGGH PGTAQPERPS PDSPWGQFAP FCHPKQSAP QEGCSPHFAV
751 APCPFGSFP GSCKEAPLVP SSPFLGQFQA PPAPAKASPP LDSKMGPGDI
801 SLPGRPKPGP CSSPGSASQA SSSQVSSLRV GSSQVGTPEG PSLDAEGWTQ
851 EAEDLSDSTP TLQRPOEQVT MRKPSLGGRG GYAGVAGYGT FAFGGDAGGM
901 LGQGPMWARI AWAVSQSEEE EQEARAESQ SEEQQEARAE SPLPQVSARP
951 VPEVGRAPTR SSPEPTPWED IGQVSLVQIR DLSGDAEAD TISLDISEVD
1001 PAYLNLSLY DIKYLPEFEM IFRKVPKSAQ PEPPSPMAEE ELAEFFPEPTW
1051 PWPGEGLGPHA GLEITEESD VDALLAEAAV GRKRKWSPPS RSLFHFPGRH
1101 LPLDEPABLG LRERVKASVE HISRILKGRP EGLEKEGPPR KKPGLASFRL
1151 SGLKSWDRAP TFLRELSDET VVLGQSVTLA CQVSAQPAQ ATWSKDGAPL
1201 ESSSRVLISA TLKNFQLTI LVVVAEDLV YTCVSNAIG TVTTTGVLRK
1251 AERPSSPCP DIGEYADGV LLVWKPVESY GPVTYIVQCS LEGGSWTTLA
1301 SDIFDCCYLT SKLSRGTYT FRTACVSKAG MGPYSSPSEQ VLLGGPSHLA
1351 SEESQGRSA QPLPSTKFA FQTQIQGRF SVVRQCWEKA SGRALAAKII
1401 PYHPKDKTAV LREYEALKGL RHPHLAQLHA AYLSPRHLVL ILELCSGPEL
1451 LPCLAERASY SESEVKDYLV QMLSATQYLH NQHILHLDLR SENMIITEYN
1501 LLKVVDLGNA QSLSQEKVLP SDRFKDYLET MAPELLEGQG AVPQTDIWI
1551 GVTAFIMLSA EYPVSSEGAR DLQRGLRKGL VRLSRCYAGL SGGAVAFRLS
1601 TLCAQPWGRP CASSCLQCPW LTEGPACSR PAPVTFPTAR LRVFVRNREK
1651 RRALLYKRHN LAQVR (SEQ ID NO:2)

```

**FEATURES:**

Functional domains and key regions:

Prosite results:

[1] PDOC00001 PS00001 ASN\_GLYCOSYLATION  
N-glycosylation site

1005-1008 NLS

---

[2] PDOC00004 PS00004 CAMP\_PHOSPHO\_SITE

cAMP- and cGMP-dependent protein kinase phosphorylation site

Number of matches: 2

1 872-875 RKFS  
2 1084-1087 RKWS

---

[3] PDOC00005 PS00005 PKC\_PHOSPHO\_SITE

Protein kinase C phosphorylation site

Number of matches: 23

1 97-99 STR  
2 152-154 SEK  
3 156-158 SHR  
4 230-232 TRK  
5 364-366 SDR  
6 450-452 SYK  
7 536-538 SER  
8 588-590 SRR  
9 668-670 TPR  
10 762-764 SCK  
11 827-829 SLR  
12 870-872 TMR  
13 947-949 SAR  
14 1147-1149 SFR  
15 1203-1205 SSR  
16 1211-1213 TLK  
17 1310-1312 TSK  
18 1320-1322 TFR

**FIGURE 2A**

19 1365-1367 STK  
 20 1391-1393 SGR  
 21 1434-1436 SPR  
 22 1521-1523 SDK  
 23 1638-1640 TAR

---

[4] PDOC00006 PS00006 CK2 PHOSPHO SITE  
 Casein kinase II phosphorylation site

Number of matches: 21

1 59-62 TIED  
 2 163-166 SFYE  
 3 242-245 SSEE  
 4 257-260 TEAE  
 5 312-315 TPAE  
 6 459-462 SIPE  
 7 491-494 SSSD  
 8 493-496 SDNE  
 9 528-531 SLPE  
 10 762-765 SCKE  
 11 915-918 SQSE  
 12 929-932 SQSE  
 13 917-920 SEEE  
 14 1351-1354 SEEE  
 15 915-918 SQSE  
 16 929-932 SQSE  
 17 961-964 SSPE  
 18 966-969 TPWE  
 19 997-1000 SEVD  
 20 1336-1339 SPSE  
 21 917-920 SEEE

---

[5] PDOC00008 PS00008 MYRISTYL  
 N-myristoylation site

Number of matches: 27

1 7-12 GCGGCS  
 2 10-15 GCSVAH  
 3 41-46 GGLHSS  
 4 42-47 GLHSSL  
 5 106-111 GTTYSL  
 6 122-127 GVTCL  
 7 133-138 GQVLCK  
 8 484-489 GGSSSS  
 9 485-490 GSSSSS  
 10 601-606 GGYIAG  
 11 606-611 GALPGL  
 12 708-713 GGHPGT  
 13 877-882 GGRGGY  
 14 880-885 GGYAGV  
 15 894-899 GGDAGG  
 16 898-903 GGMLGQ  
 17 1061-1066 GLEITE  
 18 1174-1179 GQSVTL  
 19 1229-1234 GVTCS  
 20 1240-1245 GTVTTT  
 21 1293-1298 GGSWTT  
 22 1294-1299 GSWTTL  
 23 1316-1321 GGTYYF  
 24 1508-1513 GNAQSL  
 25 1575-1580 GLRKGL  
 26 1589-1594 GLSGGA  
 27 1592-1597 GGAVAF

---

FIGURE 2B

[6] PDOC00009 PS00009 AMIDATION  
Amidation site

1080-1083 VGRK

---

[7] PDOC00373 PS00343 GRAM\_POS\_ANCHORING  
Gram-positive cocci surface proteins 'anchoring' hexapeptide

704-709 LPSTGG

---

[8] PDOC00100 PS00107 PROTEIN\_KINASE\_ATP  
Protein kinases ATP-binding region signature

171-194 IGRGVFGFVKRVQHKGNKILCAAK

---

[9] PDOC00100 PS00108 PROTEIN\_KINASE\_ST  
Serine/Threonine protein kinases active-site signature

280-292 VLHLDIKPSNIML

---

[10] PDOC00100 PS00109 PROTEIN\_KINASE\_TYR  
Tyrosine protein kinases specific active-site signature

1484-1496 ILHLDLRSENMI

---

[11] PDOC00565 PS00659 GLYCOSYL\_HYDROL\_F5  
Glycosyl hydrolases family 5 signature

142-151 LVLGGDNEPD

FIGURE 2C

SUBSTITUTE SHEET (RULE 26)

## BLAST Alignment to Top Hits:

>gi|7242949|dbj|BAA92535.1| (AB037718) KIAA1297 protein [Homo sapiens]  
Length = 2242

Score = 425 bits (1081), Expect = e-117

Identities = 305/876 (34%), Positives = 423/876 (47%), Gaps = 106/876 (12%)

Query: 54 PSMQVTIEDVQAQTGGTAQFEAIIEGDPQPSVTWYKDSVQLVDSTRLSQQQEGTYSILVL 113  
P + +EDV+ G TA+ F ++EG P P + WYKD V L +S+ +S E . SLV+  
Sbjct: 504 PRFESIMEDVEVGAGETARFAVVVEGKPLPDIMWYKDEVLLTESSHVSFVYEENECSLTV 563

Query: 114 RHVASKDAGVYTCLAQNTGGQVLCKAELLVLGGDN----EPDSEKQSHR-RKLHSFYEVK 168  
++D GVYTC AQN G+V CKAEL V E E + HR R+L FY++  
Sbjct: 564 LSTGAQDGGVYTCTAQNLAGEVSCAELAVHSAQTAMEVEGVGEDEHRRRLSDFYDIH 623

Query: 169 EEIGRGVFGFVKRVQHKGNKILCAAKFIPLRSRTAQAYRERDILAAALSHPLVTGLLDQF 228  
+EIGRG F +++R+ + + + AAKFIP +++ +A A RE +LA L H V + F  
Sbjct: 624 QEIGRGAFSYLRIVERSSGLEFAAKFIPSAKPKASARREARLLARLQHDCLVYFHEAF 683

Query: 229 ETRKTLLILILELCSSEELLDRLYRKGVTAEVKKVYIQQLVEGLHYLHSHGVHLHLDIKPS 288  
E R+ L+++ ELC+ EELL+R+ RK V E+E++ Y++Q++EG+HYLH VLHLD+KP  
Sbjct: 684 ERRRGLVIVITELCT-EELLRIARKPTVCESEIRAYMRQVLEGIHYLHSHVHLHLDVKPE 742

Query: 289 NILMVHPA--REDIKICDFGAQNITPAELQFSQYGSPEFVSPEIIQONPVSEASDIWAM 346  
N+L+ A + ++ICDFG AQ +TP E Q+ QYG+PEFV+PEI+ Q+PVS +DIW +  
Sbjct: 743 NLLVWDGAAGEQQVRICDFGNAQELTPGEPQYQYGTPEFVAPEIVNQSPVSGVTDIWPV 802

Query: 347 GVISYLSLTCSPPFAGESDRATLLNVLEGRVSWSSPMAAHLSEDAKDF-IKATLQRAPOA 405  
GV+++L LT SPF GE+DR TL+N+ V++ LS +A+ F IK +Q +  
Sbjct: 803 GVVAFCLCTGISPFVGENDRITLMNIRNYNVAFEETTFLSLSREARGFLIKVLVQ--DRL 860

Query: 406 RPSAAQCLSHFWFLKSMPEAEAHFINTKQLKELLARSRWQSRMSYKSILVMRSIPELLR 465  
RP+A + L HPWF E ++T LK L+R RWQRS +SYK LV+R IPELLR  
Sbjct: 861 RPTAETLEHPWFKTQAKAE---VSTDHLKLFSLRRRWQRSQISYKCHLVLRPIPELLR 917

Query: 466 GPPDSPSLGVARHLCRDTGGSSSSSSSDNELAPFARAK-----SLPPSPVTH 513  
PP+ + + R +GG SSSS S + EL SL P  
Sbjct: 918 APPERVWVTMPRR-PPPSGGLSSSSDSEEEELPSVPRPLQPEFSGSRVSLTDIPTED 976

Query: 514 SPLLHPRGFLRPSASLPEEAEEASERSTEAPAPPASPEGAGPPAAQGCVPRHSVI----- 567  
L P E+ A + EAP+P A P PAA G PR +  
Sbjct: 977 EALGTPETGAATPMDWQEQGRAPSQDQEAPEALPSPGQEPAA-GASPRRGELRRGSSA 1035

Query: 568 -----RSLFYHQAGESPENHALAPG-----SRRHPPARRRHLK 600  
R L + E P+ + PG ++R A R+ LL+  
Sbjct: 1036 ESALPRAGPRELGRGLHKAASVELPQRRSPGPGATRLARGGLGEGEYAQRLQALRQLLR 1095

Query: 601 GGYIAGALPLGLREPLMEH-----RVLEEEAAREEQATL----LAKAPSFETALR 645  
GG G + GLR PL+E R EAA Q L L K+ SF  
Sbjct: 1096 GGPEDGKVSGLRGFLLESGLGRARDPRMARAASSEAAPHHPPLENRLQKSSSFSGGEA 1155

Query: 646 LPASGTHLAPGHSHSLEHDSPTPR----PSSEACGEAQRLPSAPSGGAPIRDMGHFQGS 701  
P G H G + R PS A EAQ PS+P+ P  
Sbjct: 1156 EP-RGRHRRAGAPLEIPVARLGARRLOESPSLSALSEAQ--PSSPA-----RPSAP 1203

Query: 702 KQLPSTGGHPGTAQPERPSPDPSWGPAPFCHPKQGSAPQEGCSPHFAVAPCPP---GS 757  
K PST P +A+P +P PAP P Q AP+ P A P PP +  
Sbjct: 1204 K--PST---PKSAEPSATTPTSDAPQPPAP--QPAQDKAPEPRPEPVRASKPAPPPQALQT 1256

Query: 758 FPPGSCKEAPLVSSPFLGQPQAPPAPAKASPLDSKMGPGDISLPGRPKPGPCSSPGSA 817  
A ++ S G Q P+ A+PP + K + P PG +  
Sbjct: 1257 LALPLTPYAQIIQSLQLSGHAQG-PSQGPAAPPSEPKPHAADVAVASPPPG--APEKRV 1313

Query: 818 SQASSSQVSSLRVGSSQVGTPEGPSLDAEGWTQAE 853  
A V + + V PG SL + E+E  
Sbjct: 1314 PSAGGPPVLAERARVPTVPPRPGSSLSSTIENLESE 1349 (SEQ ID NO:4)

FIGURE 2D

SUBSTITUTE SHEET (RULE 26)



Score = 210 bits (529), Expect = 1e-52  
 Identities = 111/281 (39%), Positives = 156/281 (55%), Gaps = 2/281 (0%)

Query: 1336 SPSEQVLLGGPSHLASEEESQGRSAQPLPSTKTFAFQTQIQGRFVSVVRQCWEKASGRAL 1395  
 SP+++V+ S S +G + + P K + F + RGRF VVR C E A+GR  
 Sbjct: 1952 SPAKEVVSPPGSSPRSSPRPEGTTLRQGPQKPYTFLEEKARGRFGVVRACRENATGRTF 2011

Query: 1396 AAKIIPYHPKDKTAVLREYEALKGLRHPHLAQLHAAYLSRHLVLILELCSGPPELLPCLA 1455  
 AKI+PY + K VL+EYE L+ L H + LH AY++PR+LVLI E C ELL L+  
 Sbjct: 2012 VAKIVPYAAEGKPRVLQYEVVLRTHHERIMSLHEAYITPRYLVLIAESCGNRELLCGLS 2071

Query: 1456 ERASYSESEVKDYLVQMLSATQYLHNQHLHLDLRSENMIITEYNLLKVVDLGNQSLSQ 1515  
 +R YSE +V Y+ Q+L YLH H+LHLD++ +N+++ N LK+VD G+AQ +  
 Sbjct: 2072 DRFRYSEDDVATYVMVQLQLGLDYLHGHVHLHLDIKPDNLLAPDNALKIVDFGSAQPYNP 2131

Query: 1516 EKVLPSDKFKDYLETMAPELLEGQGAVPQTDIWAIGVTAFIMLSAEYPVSSEGARDLQRG 1575  
 + + P LE MAPE+++G+ TDIW GV +IMLS P ++ +  
 Sbjct: 2132 QALRPLGHRGTGLEFMAPEMVKGEPIGSATDIWGAGVLTYYIMLSGRSPFYEPDPQETEAR 2191

Query: 1576 LRKGLVRLSRCYAGLSGGAVAFRLSTLCAQPWGRPCASSCL 1616  
 + G + Y S A FLR L PW RP SSCL  
 Sbjct: 2192 IVGGRFDAFQLYPNTSQSATLFLRKVLSVHPWSRP--SSCL 2230 (SEQ ID NO:5)

Score = 170 bits (426), Expect = 1e-40  
 Identities = 168/574 (29%), Positives = 256/574 (44%), Gaps = 42/574 (7%)

Query: 1103 LDEP--AELGLRERVKASVEHISRIKGRPEGLEKEGPPRKKPGLASFRLSGLKSWDRAP 1160  
 L EP A GLR+ V+HI R+L + K PP + L L + + AP  
 Sbjct: 358 LREFGWAATGLRK----GVQHIFRVLSTTVKSSSKPSPSEFVQL----LEHGPTLEEAP 409

Query: 1161 TFLRELSDETIVLQSVTLACQVSAQPAQATW-SKDGAFLSSSRVL-ISATLKNFQLL 1218  
 L + VV GQ ++ + AQ W S GA LE+ + V +S + L  
 Sbjct: 410 AMLDKPDIVYVVEGQPASVTVTFN-HVEAQVVRWRCRGALLEARAGVYELSQPDDQYCL 468

Query: 1219 TILVVAEDLGVYTCSVSNALGTVTITGVLRKAERPS-SSPCPD----GEVIADGVLLV 1273  
 I V D+G TC+ N GT T + L AE P S D+ GE V++  
 Sbjct: 469 RICVSRDGMALTCARNRHGTQTCSVTLELAEPFESIMEDVEVGAGETARFAVVVE 528

Query: 1274 WKPVESYGPVTYIVQCSLEGGSWTTLASDIFDCCY--LTSKLSRGGTYTFRACVSKAGM 1331  
 KP+ + Y + L S + + +C L++ GG YT C ++  
 Sbjct: 529 GKPLPDI--MWYKDEVLLTESSHVSFVYEENECSLVVLSTGAQDGGVYT----CTAQNLA 582

Query: 1332 GPYSSPSEQVLLGGPSHLASEEESQGRSAQPLPSTKTFAFQTQIQGRFVSVVRQCWEKAS 1391  
 G S +E + + + E + + + + +I RG FS +R+ E++S  
 Sbjct: 583 GEVSCKAELAVHSAQTAMEVEGVGEDEHRRRLSDFYDIHQEIGRGAFSYLRRRIVERSS 642

Query: 1392 GRALAAKIIPYHPKDKTAVLREYEALKGLRHPHLAQLHAAYLSRHLVLILELCSGPPELL 1451  
 G AAK IP K K + RE L L+H + H A+ R LV++ ELC+ ELL  
 Sbjct: 643 GLEFAAKFIPSAKPKASARREARLLARLQHDVLYFHEAFERRRGLVIVTELCT-EELL 701

Query: 1452 PCLAERASYSESEVKDYLVQMLSATQYLHNQHLHLDLRSENMIITE----YNLLKVVDL 1507  
 +A + + ESE++ Y+ Q+L YLH H+LHLD++ EN+++ + +++ D  
 Sbjct: 702 ERIARKPTVCESEIRAYMRQVLEGIHYLHQSHVHLHLDVKNLLVWDGAGEQQVRICDF 761

Query: 1508 GNAQSLSQEKVLP SDKFKDYLETMAPELLEGQGAVPQTDIWAIGVTAFIMLSAEYPVSSE 1567  
 GNAQ L+ + P E +APE++ TDIW +GV AF+ L+ P E  
 Sbjct: 762 GNAQELTPGE--PQYCYGTPEFVAPEIVNQSPVSGVTDIWPVGVAFLCLTGISPFVGE 819

Query: 1568 GARDLQRLRKGLVRLSR-CYAGLSGGAVAFRLSTLCAQPWGRPCASSCLQCPWLTEEGP 1626  
 R +R V + LS A FL L Q RP A L+ PW +  
 Sbjct: 820 NDRITLMNIRNYNVAFEETTFLSLSREARGFLIKVL-VQDRLRPTAETLEHPWFKTQ-- 876

Query: 1627 ACSRPAPVTFPTARLRVFN-RNREKRALLYKRH 1659  
 ++ A V+ T L++F+ R R +R + YK H  
 Sbjct: 877 --AKGAEV--TDHLKFLSRRRWQRSQISYKCH 906 (SEQ ID NO:6)

## FIGURE 2E

SUBSTITUTE SHEET (RULE 26)

Score = 145 bits (362), Expect = 4e-33  
Identities = 85/253 (33%), Positives = 135/253 (52%), Gaps = 5/253 (1%)

Query: 165 YEVKKEIGRGVFGFVKRVQHKGNKILCAAKFIPLRSRTAQAYRERDILAALSHPLVTGL 224  
Y EE RG FG V+ + AK +P + + + +E ++L L H + L  
Sbjct: 1985 YTFLEEKARGRFGVVRACRENATGRTEFVAKIVPYAAEGKPRVLQYEYVLRTLHHERIMSL 2044

Query: 225 LDQFETRKTILILELCSSEELLDRLYRKGVVTEAEVKVYIQQLVEGLHYLHSHGVHLHD 284  
+ + T + L+LI E C + ELL L + +E +V Y+ QL++GL YLH H VLHLD  
Sbjct: 2045 HEAYITPRYLVLIAESCGNRELLCGLSDRFYSEDVATYMVQLLQGLDYLHGHVHLHD 2104

Query: 285 IKPSNILMVHPAREDIKICDFGFAQNITPAELQ--FSQYGSPEFVSPEIIQQNPVSEASD 342  
IKP N+L+ +KI DFG AQ P L+ + G+ EF++PE+++ P+ A+D  
Sbjct: 2105 IKPDNLLA--PDNALKIVDFGSAQPYNPQALRPLGHRTGTLEFMAPEMVKGEPIGSATD 2162

Query: 343 IWAMGVISYLSLTCSSPFAGESDRATLLNVLEGRVSWSSPMAAHLSEDAKDFIKATLQRA 402  
IW GV++Y+ L+ SPF + T ++ GR + + + S+ A F++ L  
Sbjct: 2163 IWGAGVLTYYIMLSGRSFFYEPDPQETEARIVGGRFD-AFQLYPNTSQSATLFLRKVLSVH 2221

Query: 403 PQARPSAAQCLSH 415  
P +RPS+ + H  
Sbjct: 2222 PWSRPSSCLSVCH 2234 (SEQ ID NO:7)

Score = 128 bits (319), Expect = 4e-28  
Identities = 81/245 (33%), Positives = 120/245 (48%), Gaps = 19/245 (7%)

Query: 1139 PRKKPGLASFRLSGL-----KSWDRAPTFLRELSDETIVVLGQSVTLACQVSAQP 1187  
PRK GL+ LS D P F +L D+ ++ G++ TL C +A P  
Sbjct: 1571 PRKDGLSPNLSASVQEEELGHQYVRSEDFPPVFIKLDQVLEGEAATLLCLPAACP 1630

Query: 1188 AAQATWSKDGAPLESSRVLISATLKNFQLLTILVVVAEDLGVTCSVSNALGTVTTTGV 1247  
A +W KD L S V+I + QLL+I G+Y CS +N LG++T++  
Sbjct: 1631 APHISWMKDKKSLRSEPSVIVSCKDGRQLLSIPRAGKRHAGLYECSATNVLGSITSST 1690

Query: 1248 LRKAERPSSSPCPDIGEVYADGVLLVWKPVESYGPVTYIVQCSLEGG-SWTTLASDIFDC 1306  
+ A P P++ + Y D L++WKP +S P TY ++ ++G S W ++S I DC  
Sbjct: 1691 VAVARVPGLAPPEVTQTYQDTALVLWKPGDSRAPCTYTLERRVDGESVWHVSSGIPDC 1750

Query: 1307 CYLTSKLSRGGTYTFRACVSKAGMGPYSSPSEQVLLGG-----PSHLASEEESQGRS 1359  
Y + L G T FR AC ++AG GP+S+ SE+V + G PS E R  
Sbjct: 1751 YYNVTHLPVGVTVRFRVACANRAGQGPFNSSEKVFVRGTQDSSAVPSAAHQEAPVTSRP 1810

Query: 1360 AQPLP 1364  
A+ P  
Sbjct: 1811 ARARP 1815 (SEQ ID NO:8)

Score = 71.0 bits (171), Expect = 9e-11  
Identities = 41/115 (35%), Positives = 57/115 (48%), Gaps = 4/115 (3%)

Query: 60 IEDVQAQTGGTAQFEAIIEGDPQPSVTWYKDSVOLVDSTRLSQOQEGTYSVLVRHVASK 119  
+EDV+ G A+F+ I G P P VTW + +S L +Q+G +SL + HV S+  
Sbjct: 89 LEDVEVLEGRAARFDCKISGTPPPVVTWTHFGCPMEESENLRRLQDGGGLSHLHIAHVGE 148

Query: 120 DAGVYTCLAQNTGGQVLCKAELLVLGGDNEPDSEKQSHRRKLHSFYEVKKEIGRG 174  
D G+Y A NT GQ C A+L V EP + KL + EE +G  
Sbjct: 149 DEGLYAVSAVNTHGQAHCSAQLYV---EEPRTAASGPPSSKLEKMPISIEPEQ 199 (SEQ ID NO:9)

Score = 60.1 bits (143), Expect = 2e-07  
Identities = 54/199 (27%), Positives = 81/199 (40%), Gaps = 12/199 (6%)

Query: 1160 PTFLELSDETIVVLGQSVTLACQVSAQPAAQATWSKDGAPLESSRVLISATLKNFQLLT 1219  
P FLR L D V L + L CQV+ P +W +G ++SS ++ ++ L  
Sbjct: 207 PDFLRPLQDLEVLAKAEMLEQCQVTGLPYPTISWFHNGHRIQSSDDRRMT-QYRDVHRLV 265

Query: 1220 ILVVVAEDLGVTCSVSNALGTVTTTGVLLRAERPSSSP--CPDIGEVYADGVLLVWKPV 1277  
V + GYV ++N LG L + P P + V V L W P  
Sbjct: 266 FPAVGPOHAGVYKSVIANKLGAACYAHLYVTDVVPGPPDGAQVAVTGRMVTLTWNPP 325

Query: 1278 ESY-----GPVTYIVQCSLEGG-SWTTLASDIFDCCYLTSKLSRGGTYTFRACVSKAG 1330

## FIGURE 2F

SUBSTITUTE SHEET (RULE 26)

9/17

S +TY VQ + G WT L + + + + L +G + FR +  
 Sbjct: 326 RSLDMAIDPDSLTYTVQHQLGSDQWTALVTGLREPGWAATGLRKGVQHIFRVLSTTVKS 385  
 Query: 1331 MGPYSSPSE--QVLLGGPS 1347  
 S PSE Q+L GP+  
 Sbjct: 386 SSKPSPSEPVLLEHGPT 404 (SEQ ID NO:10)

Score = 45.7 bits (106), Expect = 0.004  
 Identities = 30/102 (29%), Positives = 45/102 (43%), Gaps = 1/102 (0%)

Query: 1159 APTFLRELSDETIVVLGQSVTLACQVSAQPAQAATWSKDGALESSSRVLISATLKNFQLL 1218  
 AP F R L D V+ G++ C++S P TW+ G P+E S + + L  
 Sbjct: 82 APLFTRLLEDVEVLEGRAARFDCRISGTPPPVVTWTHFGCPMEESENLRRLRQD-GGLHSL 140  
 Query: 1219 TILVVVAEDLGVYTCSVSNALGTVTTTGVLRKAERPSSSPCP 1260  
 I V +ED G+Y S N G + L E +++ P  
 Sbjct: 141 HIAHVGSEDEGLYAVSAVNTHQAHCSAQLYVEEPRTAASGP 182 (SEQ ID NO:11)

Score = 43.8 bits (101), Expect = 0.015  
 Identities = 58/217 (26%), Positives = 84/217 (37%), Gaps = 23/217 (10%)

Query: 619 RVLEEEAAREEQATLLAKAPSFETALRLPASGTHLAPGHSLSLEHDSPTPRPSSEACGE 678  
 R ++ +A A A S R P S T LAP + + T PSS  
 Sbjct: 1788 RGTQDSSAVPSAAHQEAPVTSRPARARPPDSPTSLAPPLAPAAPTPPSVTVPSSPPTPP 1847  
 Query: 679 AQRLPSAPSGGAPIRDMGHPOGSKQLPSTGGHPGTAQPERSPDPSFWGQAPFCHPKQGS 738  
 +Q L S + G P + P+ + L + A+P PS +P PF  
 Sbjct: 1848 SQALSSLKAVGPPPPQTP--PRRHRLQAAR----PAEPTLPSTHVTPEPKPFVLD---- 1897  
 Query: 739 APQEGCSPHAPAVAPCPGSPFPGSCKEAPLVPSSPFLGQOQAPPAPAKASPPLDKMGPG 798  
 + P A P G P S P+ + F+ P AP PA PP +K+  
 Sbjct: 1898 -----TGTPIPASTPQGVKPVSS--STPVYVVTSEFVSAPPAPPEPPAPEPPPEPTKVTVQ 1949  
 Query: 799 DISLPGRPKPGPCSSPGSASQAS--SSQVSSLRVGSSQ 834  
 +S P SSPGS+ ++S + ++LR G Q  
 Sbjct: 1950 SLS----PAKEVVSSPGSSPRSSPRPEGTTLRQGPQ 1982 (SEQ ID NO:12)

Score = 43.0 bits (99), Expect = 0.026  
 Identities = 25/92 (27%), Positives = 44/92 (47%), Gaps = 4/92 (4%)

Query: 54 PSMQVTIEDVQAQTGGTAQFEAIIIEGDPQPSVTWYKDS--VQLVDSTRLSQQQEGTTYSL 111  
 P ++D++ A E + G P P+++W+ + +Q D R++Q ++ + L  
 Sbjct: 207 PDFLRPLQDLEVLAKAMELCQVTGLPYPTISWFHNGHRIQSSDDRRMTQYRD--VHRL 264  
 Query: 112 VLRHVASKDAGVYTCLAQNTGGQVLCKAELLV 143  
 V V + AGVY + N G+ C A L V  
 Sbjct: 265 VFFAVGPQHAGVYKSVIANKLGAACYAHLV 296 (SEQ ID NO:13)

>gi|8928460|sp|O75962|TRIO HUMAN TRIPLE FUNCTIONAL DOMAIN PROTEIN  
 (PTRF INTERACTING PROTEIN) >gi|3644048|gb|AAC43042.1|  
 (AF091395) Trio isoform [Homo sapiens]  
 Length = 3038

Score = 229 bits (579), Expect = 1e-58  
 Identities = 143/418 (34%), Positives = 215/418 (51%), Gaps = 11/418 (2%)

Query: 53 PPSMQVTIEDVQAQTGGTAQFEAIIIEGDPQPSVTWYKDSVQLVDST---RLSQQQEGTTY 109  
 PP + + +V +TG T + G P+ S+TW +++ +S G  
 Sbjct: 2625 PPEFVIPLSEVTCETGETVVLRCRVCGRPKASITWKGPEHNTLNNDGHYSISYSDLGEA- 2683  
 Query: 110 SLVLRHVASKDAGVYTCLAQNTGGQVLCKAELLVLGGDNEPDSEKQSHRRKLHSFYEVKE 169  
 +L + V ++D G+YTC+A N G A L VLG D + + SFY  
 Sbjct: 2684 TLKIVGVTTEDDGIYTCIAVNDMGSAASSASLRVLGPGM---DGMVTVKDNFDSFYSEVA 2741

## FIGURE 2G

SUBSTITUTE SHEET (RULE 26)

10/17

Query: 170 EIGRGVFGFVKRVQHKGNKILCAAKFIPLRSRTRAQAYRERDILAALSHPLVTGLLDQFE 229  
E+GRG F VK+ KG K A KF+ + R Q E IL +L HPL+ GLLD FE  
Sbjct: 2742 ELGRGRFSVVKCDQKGTKRAVATK FVNKKLMKRDQVTHELGILQSLQHPLLVLGLDTFE 2801

Query: 230 TRKTLILILELCSSEELLDRLYRKGVVTEAEVKVYIQQLVEGLHYLHSHGVLHLDIKPSN 289  
T + IL+LE+ LLD + R G +TE +++ ++ +++E + YLH+ + HLD+KP N  
Sbjct: 2802 TPTSILVLEMAHQGRLLDCVVRWGSLETKIRAHLEGEVLEAVRYLHNCRIAHLDLKPEN 2861

Query: 290 ILMVHP-AREDIKIDFGFAQNITPAELQFSQYGSPEFVSPEIIQQNPVSEASDIWANGV 348  
IL+ A+ IK+ DFG A + G+PEF +PEII NPVS SD W++GV  
Sbjct: 2862 ILVDESLAKETIKLADFGDAVQLNTTYIHQLLGNPEFAAPEIILGNPVSLTSDTWSVG 2921

Query: 349 ISYLSLTCSSPFAGESDRATLLNVLEGRVSWSSPMAAHLSEDAKDFIKATLQRAPOARPS 408  
++Y+ L+ SPF +S T LN+ S+ +S+ AK+F+ LQ P RPS  
Sbjct: 2922 LTYVLLSGVSPFLDDSVETCLNICRLDFSFPDDYFKGVSQKAKEFVCFLLOEDPAKRPS 2981

Query: 409 AAQCLSHPWFLKSMPEAEAHFINTKQLKFLIARSRWQ---RSLMSYKSILVMRSIPEL 463  
AA L W L++ ++T +L + R + Q R + S K+ L R +P +  
Sbjct: 2982 AALALQEQW-LQAGNGRSTGVLDTSRLTSFIERRKHQNDVRPIRSIKNFLQSRLLPRV 3038  
(SEQ ID NO:14)

Score = 121 bits (300), Expect = 7e-26  
Identities = 82/280 (29%), Positives = 137/280 (48%), Gaps = 10/280 (3%)

Query: 1374 QIQGRGRFSVVRQCEKASGRALAAKIIYPHPKDKTAVLREYEALKGLRHPHLAQLHAAYL 1433  
++ RGRFSV++C +K + RA+A K + + V E L+ L+HP L L +  
Sbjct: 2742 ELGRGRFSVVKCDQKGTKRAVATK FVNKKLMKRDQVTHELGILQSLQHPLLVLGLDTFE 2801

Query: 1434 SPRHLVLILELCSGPELLPCLAERASYSESEVKDYLMQMSATQYLNHILHLDLRSN 1493  
+P +L+LE+ LL C+ S +E +++ +L ++L A +YLHN I HLDL+ EN  
Sbjct: 2802 TPTSILVLEMAHQGRLLDCVVRWGSLETKIRAHLEGEVLEAVRYLHNCRIAHLDLKPEN 2861

Query: 1494 MIITE---YNLLKVVDLGNASLSQEKVLPDKFKDYLETMAPELLEGQGVAPQTDIWI 1550  
+++ E +K+ D G+A L+ + + E APE++ G +D W++  
Sbjct: 2862 ILVDESLAKETIKLADFGDAVQLNTTYI--HQLLGNPEFAAPEIILGNPVSLTSDTWSV 2919

Query: 1551 GVTAFIMLSAEYVPVSESGARDLQRL-RKGLVRLSRCYAGLSGGAVAFRLSTLCAQPWGR 1609  
GV +++LS P + + + R + G+S A F+ L P R  
Sbjct: 2920 GVLTYVLLSGVSPFLDDSVETCLNICRLDFSFPDDYFKGVSQKAKEFVCFLLOEDPAKR 2979

Query: 1610 PCASSCLQCPWLTEEGPACSRPAPVTFPTARLRFVVRNRE 1649  
P A+ LQ WL A + + T+RL F+ R+  
Sbjct: 2980 PSAALALQEQWL---QAGNGRSTGVLDTSRLTSFIERRK 3015 (SEQ ID NO:15)

Score = 55.4 bits (131), Expect = 5e-06  
Identities = 42/153 (27%), Positives = 70/153 (45%), Gaps = 17/153 (11%)

Query: 1128 GRPEGLEKEGPPRKKPGLASFRLSGLKS----WDRAPTFLRELSDETIVVLGQSVTLACQV 1183  
G+ EG + G + + GL++ L + +D P F+ LS+ T G++V L C+V  
Sbjct: 2590 GKREGKLENGYRKSREGLSNKVSVKLLNPNIYDVPPEFVIPLEVTCTETGETVVLRCRV 2649

Query: 1184 SAQPAQAQW-SKDGAPLESSSRVLISATLKNFQLLTILVVVAEDLGVTCTSVSNALGTV 1242  
+P A TW + L + IS + L I+ V ED G+YTC N +G+  
Sbjct: 2650 CGRPKASITWKGPEHNTLNNDGHYSISYSDLGEATLKIVGVTTEDDGIYTCIAVNDMGSA 2709

Query: 1243 TTTGVLKKAERPSSSPCPDIGEVYADGVLLVWK 1275  
+++ LR + DG+++ WK  
Sbjct: 2710 SSSASLR-----VLGPGMDGIMVTWK 2730 (SEQ ID NO:16)

Score = 39.1 bits (89), Expect = 0.39  
Identities = 61/208 (29%), Positives = 76/208 (36%), Gaps = 65/208 (31%)

Query: 688 GGAPIRDMGHPOGSKQLPSTGGHPGTA-----QPERPSPD-----S 723  
GGAP GH G S GG P T+ QP R P S  
Sbjct: 2252 GGAPSGSGHSGGPS---SCGGAPSTSRSPSRIPQPVRRHPPVLVSSAASSQAEADKMS 2308

FIGURE 2H

SUBSTITUTE SHEET (RULE 26)

11/17

Query: 724 PWGQPAFFCHPKQGSAPQEGCSPHPAVAPCPPGSFPPGSCKEAPLVSSPFLGQPQ---- 779  
P P P G+AP+ G S A + PPG+ GS +EA +P L P+  
Sbjct: 2309 GTSTPGPSL-PPPGAAPEAGPS---APSRPPGADAEGSEREAEPIPKMKVLESFPRKGAA 2364

Query: 780 -----APPAPAK-----ASPLDSKMGPGDISLPGRPKPGPCSSPGSA 817  
+P APAK A+ PL+S + SL P P P S  
Sbjct: 2365 NASGSSPDAPAKDARASLGTPLGKPRAGAASPLNSPLSSAVPSLGKEPPF-----PSSP 2419

Query: 818 SQASSSQVSSLRVG-SSQVG--TEPGPS 842  
Q S SS+ +S+ G T PG S  
Sbjct: 2420 LQKGSFWSSIPASPASRPGSFTFPGDS 2447 (SEQ ID NO:17)

>gi|3024081|sp|Q15746|KMLS\_HUMAN MYOSIN LIGHT CHAIN KINASE, SMOOTH  
MUSCLE AND NON-MUSCLE ISOZYMES (MLCK) [CONTAINS: TELOKIN]  
Length = 1913

Score = 206 bits (518), Expect = 2e-51  
Identities = 104/298 (34%), Positives = 173/298 (57%), Gaps = 2/298 (0%)

Query: 159 RKLHSFYEVKEEIGRGVFGFKRVQHKGNKILCAAKFIPLRSRTRAQYRER-DILAALS 217  
+K+ FY+++E +G G FG V R+ K + + A KF S + R+ I+ L  
Sbjct: 1458 QKVSDFYDIEERLGSQKFGQVFRLEVEKTRKRVWAGKFFKAYSAREKENIRQEISIMNCLH 1517

Query: 218 HPLVTGLLDQFETRKTLILILELCSSEELDLRYRKGV-VTEAEVKVYIQQLEGLHYLH 276  
HP + +D FE + +++LE+ S EL +R+ + +TE E Y++Q+ EG+ Y+H  
Sbjct: 1518 HPKLVQCVDAFEKANIVMVLEIVSGGELFERIIDEDFELTERECIKYMRQISEGVEYIH 1577

Query: 277 SHGVLHLDIKPSNILMVHPAREDIKICDFGAQNTPAELQFSQYGSPEFVSPEIIQQNP 336  
G++HLD+KP NI+ V+ IK+ DFG A+ + A +G+PEFV+PE+I P  
Sbjct: 1578 KQGVHLDLKPENIMCVNKTGTRIKLIDFGLARLENAGSLKVLFGTPEFVAPEVINYEP 1637

Query: 337 VSEASDIWAMGVISYLSITCSPFAGESDRATLLNVLEGRVSWSSPMAHLSEDARDFIK 396  
+S A+D+W++GVI Y+ ++ SPF G++D TL NV + +S+DAKDFI  
Sbjct: 1638 ISYATDMWSIGVICIYLVSGLSPFMGDNDNETLANVTSATWDFDDEAFDEISDDAKDFIS 1697

Query: 397 ATLQAPQARPSAAQCLSHPWFLKSMPAEAHFINTKQLKFLARSRWQSRSLMSYKSI 454  
L++ + R QCL HPW +K EA ++ ++K +AR +WQ++ + ++I  
Sbjct: 1698 NLLKKDMKNRLDCTQCLQHPWLMKDTKNMEAKLSKDRMKYMARKWQKTGNVRAI 1755  
(SEQ ID NO:18)

Score = 127 bits (315), Expect = 1e-27  
Identities = 134/528 (25%), Positives = 219/528 (41%), Gaps = 55/528 (10%)

Query: 1132 GLEKEGPPRRKKPGLASFRLSGLKSWDRAPTFLRELSDETIVVLGQSVTLACQVSAQPAQA 1191  
G E + +KKP + + + P ++ D+ V G+SV L +V+  
Sbjct: 1215 GTESDATVKKKPAKPTPPKAAMP-----PQIIQFPEDQKVRAGESVELFGKVTGTQPTIC 1269

Query: 1192 TWSKDGAPLESSSRVLISATLKNFQLLTILVVVAEDLGVYTCSVSNALGT----VTTTGV 1247  
TW K ++ S + + + +N LTIL E G YT V N LG+ V T V  
Sbjct: 1270 TWMKFRKQIQDSEHIKVEN-ENGSKLTILAAQEHCGCYTLLVENKLSRQAQVNLTV 1327

Query: 1248 LRKAERPSSSPCPDIGEVYADGVLLVWKPVESYGPVTYIVQCSLE-----GGSWTTLASD 1302  
+ K + P+ +PC ++ + + L W SY + + S+E +W LA+  
Sbjct: 1328 VDKPDPPAGTPCAS--DIRSSSLTLSWYG-SSYDGGSAVQSYSIEIWD SANKTWKELAT- 1383

Query: 1303 IFDCCYLTS----KLSRGGTYTFRACVSKAGMPYSSPSEQVLLGGPSHLAS----- 1351  
C TS L Y FR ++ G S SE +G  
Sbjct: 1384 ----CRSTSFNVQDLLPDHEYKFRVRAINVYGTSEPSQSELTTVGEKPEEPKMKWRCQT 1439

Query: 1352 ----EESQGRSAQPLPSTKTFAF---QTQIQGRFSVVRQCWEKASGRALAAKIIP-YH 1403  
E E. R+ K F + ++ G+F V + EK + + A K Y  
Sbjct: 1440 DDEKEPEVDYRTVTINTEQKVSDFYDIEERLGSQKFGQVFRLEVEKTRKRVWAGKFFKAYS 1499

Query: 1404 PKDKTAVLREYEALGLRHPHLAQLHAAYLSRHLVLILELCSGPELLP-CLAERASYSE 1462  
K+K + +E + L HP L Q A+ ++V++LE+ SG EL + E +E  
Sbjct: 1500 AREKENIRQEISIMNCLHHPKLVQCVDAFEKANIVMVLEIVSGGELFERIIDEDFELTE 1559

Query: 1463 SEVKDYLMQMSATQYLHNQHIHLHLDRSENMIITEY--NLLKVVDLGNQSLSQE---K 1517  
E Y+ Q+ +Y+H Q I+HLDL+ EN++ +K++D G A+ L K  
Sbjct: 1560 RECIKYMRQISEGVEYIHKQGIHLDLKPENIMCVNKTGTRIKLIDFGLARLENAGSLK 1619

Query: 1518 VLPSPDKFKDYLETMAPELLEGQGAVPQTDIWAIGVTAIFMLSAEYPVSSEGARDLQRLR 1577

## FIGURE 2I

SUBSTITUTE SHEET (RULE 26)

12/17

VL E +APE++ + TD+W+IGV +I++S P + + +  
 Sbjct: 1620 VLFGRP-----EFVAPEVINYEPISYATDMWSIGVICYILVSGLSPFMGDNDNETLANVT 1674  
 Query: 1578 KGLVRL-SRCYAGLSGGAVAFRLSTLCAQPWGRPCASSCLQCPWLTEE 1624  
 + +S A F+ + L R + CLQ PWL ++  
 Sbjct: 1675 SATWDFDDEAFDEISDDAKDFISNLLKKDMKNRLDCTQCLQHPWLMKD 1722 (SEQ ID NO:19)

Score = 64.4 bits (154), Expect = 9e-09  
 Identities = 36/106 (33%), Positives = 52/106 (48%), Gaps = 4/106 (3%)

Query: 54 PSMQVTIEDVQAQTGGTAQFEAIIEGDPQPSVTWYKDSVOLVDSTRLS-QQEGTTYSLV 112  
 P TI D++ G A+F+ IEG P P V W+KD + +S E SL+  
 Sbjct: 1808 PYFSKTIRDLEVVESGAARFDCKIEGYPDEVVWFKDDQSIRESRHFQIDYDEGNCSLI 1867  
 Query: 113 LRHVASKDAGVYTCLAQNTGGQVLCKAELLV---LGGDNEPDSEKQ 155  
 + V D YTC A N+ G+ C AEL+V G+ E + E++  
 Sbjct: 1868 ISDVCDDDAKYTCRAVNSLGEATCTAELIVETMEEGEGEGE EEE 1913 (SEQ ID NO:20)

Score = 64.0 bits (153), Expect = 1e-08  
 Identities = 35/96 (36%), Positives = 46/96 (47%)

Query: 53 PPSMQVTIEDVQAQTGGTAQFEAIIEGDPQPSVTWYKDSVOLVDSTRLSQQEGTTYSLV 112  
 PP + V + G +F I G POP VTW K +V L S R+S ++ L  
 Sbjct: 160 PKKFATKLGRVVVKEGQMGFRFSCKITGRPPQVTLKGNVPLQPSARVSVSEKNGMQVLE 219  
 Query: 113 LRHVASKDAGVYTCLAQNTGGQVLCKAELLVGGDN 148  
 + V D GVTCL N G+ AEL + G D+  
 Sbjct: 220 IHGVNQDDVGVTCLVVGSGKASMSAELSIQGLDS 255 (SEQ ID NO:21)

Score = 59.3 bits (141), Expect = 3e-07  
 Identities = 30/100 (30%), Positives = 50/100 (50%), Gaps = 3/100 (3%)

Query: 47 LPALPGPPSMQVTIE---DVQAQTGGTAQFEAIIEGDPQPSVTWYKDSVOLVDSTRLSQQ 103  
 LP P P+ + ++ D++ G + G+P P V W + ++ +S +  
 Sbjct: 613 LPVAPSKPTAFIFLQGLSDLKVMDSQVTMTVQVSGNPPPEVIWLHNGNEIQESEDHFHE 672  
 Query: 104 QEGTTYSLVLRHVASKDAGVYTCLAQNTGGQVLCKAELLV 143  
 Q GT +SL ++ V +D G YTC A N+ G+V +A L V  
 Sbjct: 673 QRGTQHSWLQIEVFPEDTGTYTCEAWNSAGEVRTQAVLTV 712 (SEQ ID NO:22)

Score = 57.4 bits (136), Expect = 1e-06  
 Identities = 32/89 (35%), Positives = 46/89 (50%), Gaps = 1/89 (1%)

Query: 1160 PTFRLSDETIVVLGQSVTLACQVSAQPAQAATWSKDGAPLESSSRVLISATLKNFOLLT 1219  
 P F +L V GQ +C+++ +P Q TW K PL+ S+RV +S Q+L  
 Sbjct: 161 PKFATKLGRVVVKEGQMGFRFSCKITGRPPQVTLKGNVPLQPSARVSVSEK-NGMQVLE 219  
 Query: 1220 ILVVVAEDLGVTCSVSNALGTVTGVL 1248  
 I V +D+GVYTC V N G + + L  
 Sbjct: 220 IHGVNQDDVGVTCLVVGSGKASMSAEL 248 (SEQ ID NO:23)

Score = 53.5 bits (126), Expect = 2e-05  
 Identities = 32/98 (32%), Positives = 46/98 (46%), Gaps = 4/98 (4%)

Query: 1159 APTFLRELSDETIVVLGQSVTLACQVSAQPAQAATWSKDGAPLESSSRVLISATLKNFOLL 1218  
 AP+F L D V+ GQ L C V P + TW +G P++ + + L  
 Sbjct: 513 APSFSSVLKDCAVIEGQDFVLQCSVRGTPVPRITWLLNGQPIQYARSTCEAGVAE----L 568  
 Query: 1219 TILVVVAEDLGVTCSVSNALGTVTGVLKRAERPSS 1256  
 I + ED G YTC NALG V+ + + E+ SS  
 Sbjct: 569 HIQDALPEDHGTYTCLAENALGQVSCSAWVTVHEKKSS 606 (SEQ ID NO:24)

## FIGURE 2J

SUBSTITUTE SHEET (RULE 26)

Score = 53.1 bits (125), Expect = 2e-05  
Identities = 37/113 (32%), Positives = 48/113 (41%), Gaps = 1/113 (0%)

Query: 1140 RKKPGLASFRSLGLKSWDRAPTFRLRELSDETIVLQSVTLACQVSAQPAQAQATWSKDGAP 1199  
+K ++L S AP FL+ LSD V+ G VT+ QVS P + W +G  
Sbjct: 603 KSSRKSEYLLLEVAPSKPTAPIFLQGLSDLKVMDSQVMTVQVSGNPPPEVIWLHNGNE 662

Query: 1200 LESSSRVLISATLKNFOLLTILVVVAEDLGVYTCVSNSALGTVTTTGVLRKAE 1252  
++ S L I V ED G YTC N+ G V T VL E  
Sbjct: 663 IQESEDHFHEQRGTQHS-LWIEVFPEDTGTYTCEAWNSAGEVRTQAVLTVQE 714 (SEQ ID NO:25)

Score = 51.9 bits (122), Expect = 5e-05  
Identities = 34/101 (33%), Positives = 50/101 (48%), Gaps = 2/101 (1%)

Query: 46 SLPALPGPPSMQVTIEDVQAQTGGTAQFEAIIEGDPQPSVTWYKDSVQLVDSTR-LSQQQ 104  
S+P L P+ + ++ + G TA+FE + G P+P VTW+++ + R L  
Sbjct: 26 SMP-LTEAPAFILPPRNLCIKEGATAKEGVRGYPEPQVTHRNQGPITSGGRFLDCG 84

Query: 105 EGTYSVLVRHVASKDAGVYTCLAQNTGGQVLCKAELLVLG 145  
T+SLV+ V +D G YTC A N G EL V G  
Sbjct: 85 IRGTFSLVIHAVHEEDRGKYTCATNGSGARQVTVELTVEG 125 (SEQ ID NO:26)

Score = 50.8 bits (119), Expect = 1e-04  
Identities = 41/182 (22%), Positives = 65/182 (35%), Gaps = 26/182 (14%)

Query: 1130 PEGLEKEGPPRKKPGLASFRSLGLKSWDRA-----PTFLRELSDETV 1171  
P G E++ P +P R GL S D P F + + V  
Sbjct: 366 PSGEERKRPAPPRPATFPTRQPLGSDVVSKAANRRIPMEGQRDSAFPKEKPSQSEV 425

Query: 1172 VLGQSVTLACQVSAQPAQAQATWSKDGAPLESSSRVLISATLKNFOLLTILVVVAEDLGVY 1231  
Q+V C+VS P + W +G P+ + L +L D G Y  
Sbjct: 426 KENQTVKRCFCEVSGIPKEVAWFLEGTFPVRRQEGSIEVYEDAGSHYLCCLKARTRDSGT 485

Query: 1232 TCSVSNSALGTVTTTGVLRKAERPSSSPCPDIGEVYADGVLLVWKPVESYGPVTYIVQCSL 1291  
+C+ SNA G V+ + L+ P V D ++ + +++QCS+  
Sbjct: 486 SCTASNAQGQVSCSWTLQVERLAVMEVAPSFSSVLKDCAVIEGQ-----DFVLQCSV 537

Query: 1292 EG 1293  
G  
Sbjct: 538 RG 539 (SEQ ID NO:27)

Score = 50.4 bits (118), Expect = 2e-04  
Identities = 26/100 (26%), Positives = 47/100 (47%), Gaps = 3/100 (3%)

Query: 54 PSMQVTIEDVQAQTGGTAQFEAIIEGDPQPSVTWYKDSVQLVDSTRLSQQQEGTYSVLV 113  
P+ + ++DV G + + DP ++ W + L + + QEG+ S+ +  
Sbjct: 1098 PAFKQKLQDVHVAEGKLLQLQCQVSSDPPATIIWTLNGKTLKTKFIILSQEGSLCSVSI 1157

Query: 114 RHVASKDAGVYTCLAQNTGGQVLCKAELLVLGGDNEPDSE 153  
+D G+Y C+A+N GQ C ++ V D+ P SE  
Sbjct: 1158 EKALLEDRLGYKCAKNDAGQAECSCQVTV---DDAPASE 1194 (SEQ ID NO:28)

Score = 50.0 bits (117), Expect = 2e-04  
Identities = 35/125 (28%), Positives = 59/125 (47%), Gaps = 16/125 (12%)

Query: 1154 KSWDRAPTFRLRELSDETIVLQSVTLACQVSAQPAQAQATWSKDGAPLESSSRVLISATLK 1213  
+S AP F ++L D V G+ + L CQVS+ P A W+ +G L+++ ++S  
Sbjct: 1092 ESQGTAPAFKQKLQDVHVAEGKLLQLQCQVSSDPPATIIWTLNGKTLKTKFIILSQE-G 1150

Query: 1214 NFQLLTILVVVAEDLGVYTC-----SVNSALGTVTTTGVLRKAERPSSSP 1258  
+ ++I + ED G+Y C +V +A + T K+ RP SS  
Sbjct: 1151 SLCSVSIKALLEDRLGYKCAKNDAGQAECSCQVTVDDAPASENTPAKENKSRRPKSSL 1210

Query: 1259 CPDIG 1263  
P +G  
Sbjct: 1211 PPVLG 1215 (SEQ ID NO:29)

## FIGURE 2K

SUBSTITUTE SHEET (RULE 26)

14/17

Score = 48.0 bits (112), Expect = 8e-04  
Identities = 26/87 (29%), Positives = 38/87 (42%)

Query: 1159 APTFLRELSDET VVLGQSVTLACQVSAQPAAQATWSKDGPLESSSRVLISATLKNFQLL 1218  
AP F+ + + G + +V P Q TW ++G P+ S R L+ ++ L  
Sbjct: 32 APAFILPPRNLCIKEGATAKFEGRVRGYPEPQVTWHRNGQPITSGGRFLLDCGIRGTFSL 91

Query: 1219 TILVVVAEDLGVYTCSVSNALGTVTTT 1245  
I V ED G YTC +N G T  
Sbjct: 92 VIHAVHEEDRGKYTCEATNGSGARQVT 118 (SEQ ID NO:30)

Score = 45.3 bits (105), Expect = 0.005  
Identities = 37/140 (26%), Positives = 54/140 (38%), Gaps = 23/140 (16%)

Query: 22 TNHPSMVGCGWHPGLCGWGGGLHSSLPALPGPPSMQVTIEDVQAQTGGTAQFEAIEGDP 81  
+N V C W + L + PS ++D G + + G P  
Sbjct: 490 SNAQQQVSCSWTLQV-----ERLAVMEVAPSFSSVLKDCAVIEGQDFVLQCSVRGTP 541

Query: 82 QPSVTWYKDS--VQLVDSTRLSQQOEGTTYSLVLRHVASKDAGVYTCLAQNTGGQVLCKA 139  
P +TW + +Q ST E L ++ +D G YTCLA+N GOV C A  
Sbjct: 542 VPRITWLLNGQPIQYARSTC-----EAGVAELHIQDALPEDHGTYTCLAENALGOVSCSA 596

Query: 140 ELLVLGGDNEPDSEKQSHRR 159  
+ V EK+S R+  
Sbjct: 597 WTV-----HEKKSSRK 608 (SEQ ID NO:31)

Score = 44.5 bits (103), Expect = 0.009  
Identities = 26/104 (25%), Positives = 44/104 (42%), Gaps = 7/104 (6%)

Query: 41 GGLHSSLPALPGPPSMQVTIEDVQAQTGGTAQFEAIEGDPQPSVTWYKDSVQLV-DSTR 99  
G S+ P P Q + + T +F + G P+P V W+ + +  
Sbjct: 407 GQRDSAFPKEFKPQSQ-----EVKENQTVKFRCEVSGIPKPEVAWFLEGT FVRRQEGS 460

Query: 100 LSQQOEGTTYSLVLRHVASKDAGVYTCLAQNTGGQVLCKAELLV 143  
+ ++ ++ L L ++D+G Y+C A N GOV C L V  
Sbjct: 461 IEVYEDAGSHYLCLLKARTRDSGTYSCTASNAQQQVSCSWTLQV 504 (SEQ ID NO:32)

Score = 44.1 bits (102), Expect = 0.012  
Identities = 26/82 (31%), Positives = 38/82 (45%), Gaps = 1/82 (1%)

Query: 63 VQAQTGGTAQFEAIEGDPQPSVTWYKDSVQLV-DSTRLSQQOEGTTYSLVLRHVASKDA 121  
V A G + I GDP P+V W +D L D+ Q ++LVL+ V A  
Sbjct: 730 VTASLGQSVLISCAIAGDPFPTVHWLRDGLCKDTGHFEVLQNEVDVFTLVLLKKVQFWHA 789

Query: 122 GVYTCLAQNTGGQVLCKAELLV 143  
G Y L +N G+ C+ L++  
Sbjct: 790 GQYEILLKNRVGECSCQVSLML 811 (SEQ ID NO:33)

Score = 43.8 bits (101), Expect = 0.015  
Identities = 26/89 (29%), Positives = 35/89 (39%)

Query: 1160 PTFLELSDET VVLGQSVTLACQVSAQPAAQATWSKDGPLESSSRVLISATLKNFQLLT 1219  
P F + + D VV G + C++ P + W KD + S I L  
Sbjct: 1808 PYFSKTIRDLEVVEGSAARFDCKIEGYPDPEVVWFKDDQSIRESRHFQIDYDEDEGNCSLI 1867

Query: 1220 ILVVVAEDLGVYTCSVSNALGTVTTTGVL 1248  
I V +D YTC N+LG T T L  
Sbjct: 1868 ISDVCGDDDAKYTCKAVNSLGEATCTAEL 1896 (SEQ ID NO:34)

FIGURE 2L

SUBSTITUTE SHEET (RULE 26)



1 CAGCACGAGG AACTCCTTCT GATCACCTGG CCAGCTGAGG TCAGAGTGGG  
 51 AGAGGCAGTG GTTCCATTGA AGGAGTACTC CTAAGTGTCA GAAGCCTGGG  
 101 CGGTACAGAT GGGGTGCTGT CGCTTGGGCT GCGGGGGGTG TTCAGTTGCC  
 151 CACAGTGTAT CTCAGGGTCT CACCAACCAT CCAAGCATGG TAGGCTGTGG  
 201 CTGGCACCCA GGGTTGTGTG GCTGGGGAGG TGGTCTCCAC AGTTCCTCC  
 251 CTGCCCTCCC AGGGCCCCCA TCCATGCAGG TAACCATCGA GGATGTGCAG  
 301 GCACAGACAG GCGGAACGGC CCAATTCCAG GCTATCATTG AGGGCGACCC  
 351 ACAGCCCTCG GTGACCTGGT ACAAGGACAG CGTCCAGCTG GTGGACAGCA  
 401 CCCGGCTTAG CAGCAGACAA GAAGGCACCA CATACTCCCT GGTGCTGAGG  
 451 CATGTGGCCT CGAAGGATGC CGGCGTTTAC ACCTGCCTGG CCCAAAACAC  
 501 TGGTGGCCAG GTGCTCTGCA AGGCAGAGCT GCTGGTGTCT GGGGGGGACA  
 551 ATGAGCCGGA CTCAGAGAAG CAAAGCCACC GGAGGAAGCT GCATCTCTTC  
 601 TATGAGGTCA AGGAGGAGAT TGAAGGGGCT GTGTTTGGCT TCGTAAAAAG  
 651 AGTGCAGCAC AAAGGAAACA AGATCTTGTG CGTGCCTAAG TTCATCCCCC  
 701 TACGGAGCAG AACTCGGGCC CAGGCATACA GGGAGCGAGA CATCTGGCC  
 751 GCGCTGAGCC ACCCGCTGGT CACGGGGCTG CTGGACCACT TTGAGACCCG  
 801 CAAGACCTCT ATCTCATCC TGGAGCTGTG CTCATCCGAG GAGTGTCTGG  
 851 ACCGCTGTGA CAGGAAGGGC GTGGTGACGG AGGCCGAGGT CAAGGTCTAC  
 901 ATCCAGCAGC TGGTGGAGGG GCTGCATAC CTGCACAGCC ATGGCGTTCT  
 951 CCACCTGGAC ATAAAGCCCT CTAACATCCT GATGGTGCAT CCTGCCCGGG  
 1001 AAGACATTAA AATCTGCGAC TTTGGCTTTG CCCAGAACAT CACCCAGCA  
 1051 GAGCTGCAGT TCAGCCAGTA CGGCTCCCTT GAGTTCGTCT CCCCCGAGT  
 1101 CATCCAGCAG AACCTGTGA GCGAAGCCTC CGACATTGG GCCATGGGTG  
 1151 TCATCTCCTA CCTCAGCCTG ACCTGCTCAT CCCCATTGTC CGGCGAGAGT  
 1201 GACCGTGCCA CCTCTCTGAA CGTCTGGAG GGGCGCGTGT CATGGAGCAG  
 1251 CCCCATGGCT GCCACCTCA GCGAAGACGC CAAAGACTTC ATCAAGGCTA  
 1301 CGCTGCAGAG AGCCCCCTAG GCCCGGCCCTA GTGCGGCCCA GTGCTCTCC  
 1351 CACCCCTGGT TCCTGAAATC CATGCCTGCG GAGGAGGCCC ACTTCATCAA  
 1401 CACCAAGCAG CTCAGTTCC TCCTGGCCCG AAGTCGCTGG CAGCGTTCCC  
 1451 TGATGAGCTA CAAGTCCATC CTGGTGATGC GCTCCATCCC TGAGCTGCTG  
 1501 CGGGGGCCAC CCGACAGCCC CTCCTCTGGC GTAGCCCGGC ACCTCTGCAG  
 1551 GGACACTGGT GGCTCTCCA GTTCTCTCTC CTCCTCTGAC AACGAGCTCG  
 1601 CCCCATTGTC CCGGGCTAAG TCACTGCCAC CCTCCCGGT GACACACTCA  
 1651 CCACTGTGTC ACCCCCGGGG CTTCTCTGCG CCCTCGGCCA GCCTGCCTGA  
 1701 GGAAGCCGAG GCCAGTGAGC GCTCCACCGA GGCCCCAGCT CCGCTGCAT  
 1751 CTCGCCAGGG TGCCGGGCCA CCGGCCGCCC AGGGCTGCGT GCCCGGCAC  
 1801 AGCGTCATCC GCAGCCTGTT CTACCACAG GCGGGTGAGA GCCCTGAGCA  
 1851 CGGGGCCCTG GCCCGGGGGA GCAGGCGGCA CCGGGCCCGG CGGCGGCACC  
 1901 TGCTGAAGGG CGGCTACATT GCGGGGGCGC TGCCAGGCCT GCGCGAGCCA  
 1951 CTGATGAGAG ACCGCGTGTG GGAGGAGGAG GCCGCCAGGG AGGAGCAGGC  
 2001 CACCCCTCTG GCCAAAGCCC CCTCATTGCA GACTGCCCTC CGGCTGCTG  
 2051 CCTCTGGCAC CCACTTGGCC CTTGGCCACA GCCACTCCCT GGAACATGAC  
 2101 TCTCCGAGCA CCCCCCGCCC CTCCTCGGAG GCCTGCGGTG AGGCACAGCG  
 2151 ACTGCCCTCA GCCCCCTCCG GGGGGGCCCC TATCAGGGAC ATGGGGCACC  
 2201 CTCAGGGCTC CAAGCAGCTT CCATCCACTG GTGGCCACCC AGGCACTGCT  
 2251 CAGCCAGAGA GGCCATCCCC GGACAGCCCT TGGGGGAGC CAGCCCTTT  
 2301 CTGCCACCCC AAGCAGGGTT CTGCCCCCA GGAGGGCTGC AGCCCCCACC  
 2351 CAGCAGTTGC CCAATGCCCT CTTGGCTCCT TCCCTCCAGG ATCTTGCAAA  
 2401 GAGGCCCTCT TAGTACCCTC AAGCCCTTC TTGGGACAGC CCCAGGCACC  
 2451 CCTTGGCCCT GCCAAAGCAA GCGCCCATTT GGACTCTAAG ATGGGGCTG  
 2501 GAGACATCTC TCTTCTGGG AGGCCAAAAC CCGGCCCTG CAGTTCCCA  
 2551 GGGTCAGCCT CCCAGGCGAG CTCCTCCCAA GTGAGCTCCC TCAGGGTGGG  
 2601 CTCCTCCAG GTGGGCACAG AGCCTGGCCC CTCCTTGAT GCGGAGGGCT  
 2651 GGACCCAGGA GGCTGAGGAT CTGTCCGACT CCACACCCAC CTTGCAGCGG  
 2701 CCTCAGGAAC AGGTGACCAT GCGCAAGTTC TCCCTGGGTG GTCGCGGGG  
 2751 CTACGAGGCG GTGGCTGGCT ATGGCACCTT TGCTTTGGT GGAGATGCAG  
 2801 GGGGCATGCT GGGGAGGGG CCCATGTGGG CCAGGATAGC TGGGCTGTG  
 2851 TCCCAGTCGG AGGAGGAGGA GCAGGAGGAG GCCAGGGCTG AGTCCAGTC  
 2901 GGAGGAGCAG CAGGAGGCCA GGGCTGAGAG CCCACTGCCC CAGGTCACTG  
 2951 CAAGGCCCTG CCTGAGGTC GGCAGGGCTC CCACCAGAG CTCCTCAGAG  
 3001 CCCACCCCAT GGGAGGACAT CGGGAGGTC TCCCTGGTGC AGATCCGGGA  
 3051 CCTGTAGGT GATGCGGAGG CCGCCGACAC AATATCCCTG GACATTTCCG  
 3101 AGGTGGACCT CGCTACCTC AACCTCTCAG ACCTGTACGA TATCAAGTAC  
 3151 CTCCTCATTC AGTTTATGAT CTTAGGAAA GTCCCCAAGT CCGCTCAGCC  
 3201 AGAGCCGCCC TCCCCATGG CTGAGGAGGA GCTGGCCGAG TTCCCGGAGC  
 3251 CCACGTGGCC CTGGCCAGGT GAACTGGGCC CCCACGACAG CCTGGAGATC  
 3301 ACAGAGGAGT CAGAGGATGT GGACGCGCTG CTGGCAGAGG CTGCCGTGGG  
 3351 CAGGAAGGCG AAGTGGTCTT CGCCGTACAG CAGCTCTTC CACTTCCCTG  
 3401 GGAGGCACCT GCCGCTGGAT GAGCCTGCAG AGCTGGGGCT GCGTGAGAGA  
 3451 GTGAAGGCTT CCGTGGAGCA CATCTCCCG ATCTGAAGG GCAGGCGGGA  
 3501 AGGTCTGAG AAGGAGGGG CCCCAGGAA GAAGCCAGGC CTTGCTTCTT  
 3551 TCCGGCTCTC AGGTCTGAAG AGCTGGGACC GAGCGCCGAC ATTCTAAGG  
 3601 GAGCTCTCAG ATGAGACTGT GGTCTTGGC CAGTCAGTGA CACTGGCCTG

FIGURE 3A

SUBSTITUTE SHEET (RULE 26)

3651 CCAGGTGTCA GCCCAGCCAG CTGCCCAGGC CACCTGGAGC AAAGACGGAG  
 3701 CCCCCCTGGA GAGCAGCAGC CGTGTCTCTCA TCTCTGCCAC CCTCAAGAAC  
 3751 TTCCAGCTTC TGACCATCCT GGTGGTGGTG GCTGAGGACC TGGGTGTGTA  
 3801 CACCTGCAGC GTGAGCAATG CGCTGGGGAC AGTGACCACC ACGGGCGTCC  
 3851 TCCGGAAGGC AGAGCGCCCC TCATCTTCGC CATGCCCGGA TATCGGGGAG  
 3901 GTGTACGCGG ATGGGGTGCT GCTGGTCTGG AAGCCCGTGG AATCCTACGG  
 3951 CCCTGTGACC TACATTGTGC AGTGCAGCCT AGAAGGCGGC AGCTGGACCA  
 4001 CACTGGCCTC CGACATCTTT GACTGCTGCT ACCTGACCAG CAAGCTCTCC  
 4051 CGGGGTGGCA CCTACACCTT CCGCAGCGCA TGTGTAGCA AGGCAGGAAT  
 4101 GGGTCCCTAC AGCAGCCCTT CGGAGCAAGT CCTCTGGGA GGGCCAGCC  
 4151 ACCTGGCCTC TGAGGAGGAG AGCCAGGGGC GGTGAGCCCA ACCCCTGCCC  
 4201 AGCACAAAGA CCTTCGCATT CCAGACACAG ATCCAGAGGG GCCGCTTCAG  
 4251 CGTGGTGGCG CAATGCTGGG AGAAGGCCAG CGGGCGGGCG CTGGCCGCCA  
 4301 AGATCATCCC CTACCACCCC AAGGACAAGA CAGCAGTGCT GCGCGAATAC  
 4351 GAGGCCCTCA AGGGCCTGCG CCACCCGCAC CTGGCCAGC TGCACGCAGC  
 4401 CTACCTCAGC CCCCAGCACC TGGTGTCTAT CTGGAGCTG TGCTCTGGGC  
 4451 CCGAGCTGCT CCCCTGCCTG GCCGAGAGGG CCTCCTACT AGAATCTGAG  
 4501 GTGAAGGACT ACCTGTGGCA GATGTTGAGT GCCACCCAGT ACCTGCACAA  
 4551 CCAGCACATC CTGCACCTGG ACCTGAGGTC CGAGAATATG ATCATCACCG  
 4601 AATACAACTT GCTCAAGGTC GTGGACCTGG GCAATGCACA GAGCCTCAGC  
 4651 CAGGAGAAGG TGCTGCCCTC AGACAAGTTC AAGGACTACC TAGAGACCAT  
 4701 GGCTCCAGAG CTCCTGGAGG GCCAGGGGGC TGTTCACAG ACAGACATCT  
 4751 GGGCCATCGG TGTGACAGCC TTCATCATGC TGAGCGCCGA GTACCCGGTG  
 4801 AGCAGCGAGG GTGCACGCGA CCTGCAGAGA GGAAGTCCGA AGGGGCTGGT  
 4851 CCGGCTGAGC CGCTGCTACG CGGGGCTGTC CGGGGGCGCC GTGGCCTTCC  
 4901 TGCGCAGCAC TCTGTGCGCC CAGCCCTGGG GCCGGCCCTG CGCGTCCAGC  
 4951 TGCTTGCAGT GCCCGTGGCT AACAGAGGAG GGCCCGGCCT GTTCGCGGCC  
 5001 CCGCGCCGTG ACCTTCCCTA CCGCGCGGCT GCGCGTCTTC GTGCGCAATC  
 5051 GCGAGAAGAG ACGCGCGCTG CTGTACAAGA GGCACAACCT GGCCAGGTG  
 5101 CGCTGAGGGT CGCCCCGGCC ACACCCCTGG TCTCCCCGCT GGGGGTCTGCT  
 5151 GCAGACGCGC CAATAAAAC GCACAGCCGG GCGAGAAAAA AAAAAAAAAA  
 5201 AAAAAA (SEQ ID NO:3)

## FEATURES:

Start: 109  
 Exon: 109-5103  
 Stop: 5104

## SNPs:

DNA				Protein		
Position	Major	Minor	Domain	Position	Major	Minor
311	T	C G	Exon	68	V	A G
1741	C	T	Exon	545	P	S
2714	T	C	Exon	869	V	A
2745	C	T	Exon	879	R	R
2859	A	G	Exon	917	S	S
3420	T	C	Exon	1104	D	D

## Context:

## DNA

## Position

311 AACTCCTTCTGATCACCTGGCCAGCTGAGGTCAGAGTGGGAGAGGCAGTGGTTCCATTGA  
 AGGAGTACTCCTAACTGTGCAAGCCTGGGCGGTCAGGATGGGGTGCTGTCGCTTGGGCT  
 GCGGGGGGTGTTCAAGTTGCCACAGTGTATCTCAGGGTCTACCAACCATCCAAGCATGG  
 TAGGGTGTGGCTGGCAGCCAGGGTTGTGTGGCTGGGGAGGTGGTCTCCACAGTTCCTCC  
 CTGCCCTCCAGGGCCCCATCCATGCAGGTAACCATCGAGGATGTGCAGGCACAGACAG  
 [T, C, G]  
 CGGAACGGCCCCAATTCGAGGCTATCATTGAGGGCGACCCACAGCCCTCGGTGACCTGGTA  
 CAAGGACAGCGTCCAGCTGGTGGACAGCACCCGGCTTAGCCAGCAGCAAGAAGGCACAC  
 ATACTCCCTGGTGTGAGGCATGTGGCCTCGAAGGATGCCGGCGTTACACCTGCCTGGC  
 CCAAACACTGGTGGCCAGGTGCTCTGCAAGGCAGAGTGTGGTGTGGGGGGGACAA  
 TGAGCCGGACTCAGAGAAGCAAGCCACCGAGGAAGCTGCACTCCTTCTATGAGGTCAA

FIGURE 3B

SUBSTITUTE SHEET (RULE 26)

1741 CAGCGTTCCCTGATGAGCTACAAGTCCATCCTGGTGATGCGCTCCATCCCTGAGCTGCTG  
CGGGGGCCACCCGACAGCCCTCCCTCGGCGTAGCCCGGCACCTCTGACGGGACACTGGT  
GGCTCCTCCAGTTCCCTCCTCCTCCTCTGACAACGAGCTCGCCCCATTGCGCCGGGCTAAG  
TCACTGCCACCCCTCCCGGTGACACACTCACCCTGCTGCACCCCGGGGCTTCTGCGG  
CCCTCGGCCAGCCTGCCTGAGGAAGCCGAGGCCAGTGAGCGCTCCACCGAGGCCCCAGCT  
[C, T]  
CGCTGCATCTCCCGAGGGTGCCGGGCCACCGGCCGCCAGGGCTGCGTGCCCCGGCACA  
GCGTCATCCGAGCCTGTCTACCACAGCGGGTGAGAGCCCTGAGCACGGGGCCCTGG  
CCCCGGGGAGCAGGGCGCACCCGGCCCGGCGGCCACCTGCTGAAGGGCGGCTACATTG  
CGGGGGCGCTGCCAGGCCTGCGCGAGCCACTGATGGAGCACCGCGTCTGGAGGAGGAGG  
CGCGCAGGGAGGAGCAGGCCACCCTCCTGGCCAAAGCCCCCTCATTGAGACTGCCCTCC

2714 TACCCTCAAGCCCTTCTTGGGACAGCCCCAGGCACCCCTGCCCTGCCAAAGCAAGCC  
CCCCATTGGACTCTAAGATGGGGCTGGAGACATCTCTTCTTGGGAGGCCAAACCCG  
GCCCTGCACTTCCCCAGGCTCAGCCTCCAGGCCAGCTCTTCCCAAGTGAGCTCCCTCA  
GGGTGGGCTCCTCCAGGTGGGCACAGAGCCTGGCCCTCCTTGGATGCGGAGGGCTGGA  
CCCAGGAGGCTGAGGATCTGTCCGACTCCACACCCACCTTGACGCGGCTCAGGAACAGG  
[T, C]  
GACCATGCGCAAGTTCTCCTTGGGTGGTTCGCGGGGGTACGCAGCGTGGCTGGCTATGG  
CACCTTTGCCTTGGTGGAGATGCAGGGGGCATGCTGGGGCAGGGGCCCATGTGGGCCAG  
GATAGCCTGGGCTGTGTCCAGTCGGAGGAGGAGGAGCAGGAGGAGGCCAGGGCTGAGTC  
CCAGTCGGAGGAGCAGCAGGAGGCCAGGGCTGAGAGCCCACTGCCCAAGTCAAGTCAAG  
GCCTGTGCCTGAGGTGCGCAGGGCTCCACAGGAGCTCTCCAGAGCCACCCCATGGGA

2745 GGCACCCCTGCCCCCTGCCAAAGCAAGCCCCCATTTGACTCTAAGATGGGGCTGGAGA  
CATCTCTTCTTGGGAGGCCAAAACCCGGCCCTGCAAGTTCCCCAGGGTCAGCCTCCCA  
GGCGAGCTCTTCCCAAGTGAGCTCCCTCAGGGTGGGCTCCTCCCAAGTGGGCACAGAGCC  
TGGCCCCCTCCTGGATGCGGAGGGCTGGACCCAGGAGGCTGAGGATCTGTCCGACTCCAC  
ACCCACCTTGACGCGGCTCAGGAACAGGTGACCATGCGCAAGTTCTCCTGGGTGGTGC  
[C, T]  
GGGGGCTACGCAGGCGTGGCTGGCTATGGCACCTTTGCCTTTGGTGGAGATGCAGGGGGC  
ATGCTGGGGCAGGGGCCCATGTGGGCCAGGATAGCCTGGGCTGTGTCCCAAGTCGGAGGAG  
GAGGAGCAGGAGGAGGCCAGGGCTGAGTCCAGTCGGAGGAGCAGCAGGAGGCCAGGGCT  
GAGAGCCCACTGCCCCAGGTGAGTGAAGGCCTGTGCTGAGGTGCGCAGGGCTCCACCC  
AGGAGCTCTCCAGAGCCCAACCCATGGGAGGACATCGGGCAGGTCTCCTTGGTGCAGATC

2859 CTCCAGGCGAGCTCTTCCCAAGTGAGCTCCCTCAGGGTGGGCTCCTCCCAAGTGGGCAC  
AGAGCCTGGCCCTCCTTGGATGCGGAGGGCTGGACCCAGGAGGCTGAGGATCTGTCCGA  
CTCCACACCCACCTTGACGCGGCTCAGGAACAGGTGACCATGCGCAAGTTCTCCTTGGG  
TGGTGGCGGGGCTACGCAGGCGTGGCTGGCTATGGCACCTTTGCCTTTGGTGGAGATGC  
AGGGGGCATGTGGGCGAGGGGCCATGTGGGCCAGGATAGCCTGGGCTGTGTCCAGTC  
[A, G]  
GAGGAGGAGGAGCAGGAGGAGGCCAGGGCTGAGTCCCAAGTCGGAGGAGCAGCAGGAGGCC  
AGGGCTGAGAGCCCACTGCCCAAGTCAAGTGAAGGCCTGTGCTGAGGTGCGCAGGGCT  
CCCACAGGAGCTCTCCAGAGCCACCCCATGGGAGGACATCGGGCAGGTCTCCTTGGTG  
CAGATCCGGGACCTGTCAAGTGATGCGGAGGCGGCCGACACAATATCCTGGACATTTCC  
GAGGTGGACCCCGCTACCTCAACCTCTCAGACCTGTACGATATCAAGTACCTCCCATTC

3420 CAACCTCTCAGACCTGTACGATATCAAGTACCTCCCATTCGAGTTTATGATCTTCAGGAA  
AGTCCCCAAGTCCGCTCAGCCAGAGCCGCCCTCCCCATGGCTGAGGAGGAGCTGGCCGA  
GTTCCCGGAGCCACGTGGCCCTGGCCAGGTGAAGTGGGCCCCACGAGGCTTGAGAT  
CACAGAGGAGTCAGAGGATGTGACCGCTGCTGCGAGAGGCTGCCGTGGGAGGAAGCG  
CAAGTGGTCTCGCCGTACGCGAGCCTCTTCCACTTCCCTGGGAGGCACCTGCCGTGGA  
[T, C]  
GAGCCTGCAGAGCTGGGGCTGCGTGAGAGAGTGAAGGCCTCCGTGGAGCACATCTCCCGG  
ATCTGAAGGGCAGGCCGGAAGGTCTGGAGAAGGAGGGGCCCCCAGGAAGAAGCCAGGC  
CTTGCTTCTTCCGGCTCTCAGGTCTGAAGAGCTGGGACCGAGCGCCGACATTCTTAAGG  
GAGCTCTCAGATGAGACTGTGTCTGGGCCAGTCACTGACACTGGCCTGCCAGGTGTCA  
GCCAGCCAGCTGCCAGGCCACCTGGAGCAAAGACGGAGCCCCCTGGAGAGCAGCAGC

Chromosome map position: 1

Bac accession number: AC023889

## FIGURE 3C

SUBSTITUTE SHEET (RULE 26)

# SEQUENCE LISTING

<110> PE CORPORATION (NY)

<120> ISOLATED HUMAN KINASE PROTEINS, NUCLEIC  
ACID MOLECULES ENCODING HUMAN KINASE PROTEINS, AND USES  
THEREOF

<130> CL000927PCT

<140> TO BE ASSIGNED

<141> 2001-22-10

<150> 09/858,664

<151> 2001-17-05

<150> 09/711,134

<151> 2000-14-11

<160> 34

<170> FastSEQ for Windows Version 4.0

<210> 1

<211> 5207

<212> DNA

<213> Human

<400> 1

```

cagcacgagg aactccttct gatcacctgg ccagctgagg tcagagtggg agaggcagtg 60
gttccattga aggagtactc ctaactgtca gaagcctggg cggtcaggat ggggtgctgt 120
cgcttgggct gcgggggggtg ttccagttgcc cacagtgtat ctcagggtct caccaacccat 180
ccaagcatgg taggtgtggt ctggcaccca ggggtgtgtg gctggggagg tggctccac 240
agttccctcc ctgccctccc agggcccca tccatgcagg taaccatcga ggatgtgcag 300
gcacagacag gcggaacggc ccaattcgag gctatcattg agggcgaccc acagccctcg 360
gtgacctggt acaaggacag cgtccagctg gtggacagca cccggcttag ccagcagcaa 420
gaaggacca catactccct ggtgctgagg catgtggcct cgaaggatgc cggcgtttac 480
acctgcctgg cccaaaacac tgggtggcag gtgctctgca aggcagagct gctggtgctt 540
gggggggaca atgagccgga ctcagagaag caaagccacc ggaggaagct gcactccttc 600
tataggttca aggagagat tggaaagggc gtgtttggct tcgtaaaaag agtgcagcac 660
aaaggaaaca agatcttgtg cgctgccaag ttcattcccc tacggagcag aactcgggcc 720
caggcataca gggagcgaga cactctggcc gcgctgagcc acccgctggt cacggggctg 780
ctggaccagt ttgagaccgg caagaccctc atcctcatcc tggagctgtg ctcattccgag 840
gagctgctgg accgcctgta caggaagggc gtggtgacgg aggccgaggt caaggtctac 900
atccagcagc tgggtggaggg gctgcactac ctgcacagcc atggcgttct ccacctggac 960
ataaagccct ctaacatcct gatggtgcat cctgcccggg aagacattaa aatctgcgac 1020
tttggttttg ccagaaacat caccacagca gagctgcagt tcagccagta cggctccctt 1080
gagttcgtct ccccgagat catccagcag aacctgtgta gcgaagcctc cgacatttgg 1140
gccatgggtg tcattcctta cctcagcctg acctgctcat ccccatttgc cggcgagagt 1200
gaccgtgcca cctcctgaa cgtcctggag gggcgctgt catggagcag ccccatggct 1260
gcccacctca gcgaagacgc caaagacttc atcaaggcta cgctgcagag agccctcag 1320
gcccggccta gtgcggccca gtgcctctcc caccctggt tcctgaaatc catgcctgcg 1380
gaggaggccc attcatcaa caccaagcag ctcaagttcc tcctggcccc aagtcgctgg 1440
cagcgttccc tgatgagcta caagtccatc ctggtgatgc gctccatccc tgagctgctg 1500
cggggccccg ccgacagccc ctccctcggc gtagccgggc acctctgag ggacactggt 1560
ggctcctcca gttcctctc ctctctgac aacgagctcg ccccatttgc ccgggctaag 1620
tactgccac cctccccggg gacacactca cactgctgc accccggggg ctctctgcg 1680
cctcggcca gcctgctga ggaagccgag gccagtgagc gctccaccga ggccccagct 1740
ccgctgcat ctcccgaggg tgccgggcca ccggccggcc agggctgctg gccccggcac 1800

```

agcgtcatcc	gcagcctgtt	ctaccaccag	gcgggtgaga	gccctgagca	cggggccctg	1860
gccccgggga	gcaggcgga	cccggccccg	cggcggcacc	tgctgaagg	cggctacatt	1920
gcgggggagc	tgccaggcct	gcgcgagcca	ctgatggagc	accgcgtgct	ggaggaggag	1980
gccgccaggg	aggagcaggc	caccctcctg	gccaaagccc	cctcattcga	gactgccctc	2040
cggtgcctg	cctctggcac	ccacttggcc	cctggccaca	gccactccct	ggaacatgac	2100
tctccgagca	ccccccgccc	ctcctcggag	gcctgcggtg	aggcacagcg	actgccttca	2160
gccccctccg	ggggggcccc	tatcagggac	atggggcacc	ctcagggctc	caagcagctt	2220
ccatccactg	gtggccaccc	aggcactgct	cagccagaga	ggccatcccc	ggacagccct	2280
tggggggcagc	cagccccttt	ctgccacccc	aagcagggtt	ctgcccccca	ggagggctgc	2340
agccccacc	cagcagttgc	cccatgccct	cctggctcct	tccctccagg	atcttgcaaa	2400
gaggccccct	tagtaccctc	aagccccctc	ttgggacagc	cccaggcacc	ccctgccctc	2460
gccaaagcaa	gccccccatt	ggactctaag	atggggcctg	gagacatctc	tcttctctggg	2520
aggccaaaac	ccggcccctg	cagttcccca	gggtcagcct	cccaggcgag	ctcttcccaa	2580
gtgagctccc	tcagggtggg	ctcctcccag	gtgggcacag	agcctggccc	ctccctggat	2640
gcggagggt	ggaccagga	ggctgaggat	ctgtccgact	ccacaccac	cttgacgcgg	2700
cctcagggaac	aggtgacat	gcgcaagttc	tccctgggtg	gtcgcggggg	ctacgcaggc	2760
gtggctggct	atggcacctt	tgccttttgt	ggagatgcag	ggggcatgct	ggggcagggg	2820
cccatgtggg	ccaggatagc	ctgggctgtg	tcccagtcgg	aggaggagga	gcaggaggag	2880
gccagggctg	agtcccagtc	ggaggagcag	caggaggcca	gggctgagag	ccactgccc	2940
caggtcagtg	caaggcctgt	gcctgaggtc	ggcagggtc	ccaccaggag	ctctccagag	3000
cccaccccat	gggaggacat	cgggcaggtc	tccctggtgc	agatccggga	cctgtcaggt	3060
gatgcggagg	cggccgacac	aatatccctg	gacatttccg	aggtggaccc	cgcctacctc	3120
aacctctcag	acctgtacga	tatcaagtac	ctcccattcg	agtttatgat	cttcaggaaa	3180
gtccccaagt	ccgctcagcc	agagccgccc	tccccatgg	ctgaggagga	gctggccgag	3240
ttcccgagc	ccacgtggcc	ctggccaggt	gaactgggcc	cccacgcagg	cctggagatc	3300
acagaggagt	cagaggatgt	ggacgcgctg	ctggcagagg	ctgccgtggg	caggaagcgc	3360
aagtgtcct	cgccgtcacg	cagcctcttc	cacttccctg	ggaggcacct	gccgctggat	3420
gagcctgcag	agctggggct	gcgtgagaga	gtgaaggcct	ccgtggagca	catctcccgg	3480
atcctgaagg	gcagcccgga	aggtctggag	aaggaggggc	ccccaggaa	gaagccagge	3540
cttgcttcct	tccggctctc	aggtctgaag	agctgggacc	gagcgccgac	attcctaagg	3600
gagctctcag	atgagactgt	ggtcctgggc	cagtcagtga	cactggcctg	ccaggtgtca	3660
gcccagccag	ctgccaggc	cacctggagc	aaagacggag	cccctctgga	gagcagcagc	3720
cgtgtcctca	tctctgcac	cctcaagaac	ttccagcttc	tgaccatcct	ggtggtggtg	3780
gctgaggacc	tgggtgtgta	cacctgcagc	gtgagcaatg	cgctggggac	agtgaccacc	3840
acggggtcc	tccggaaggc	agagcgcccc	tcactctgc	catgcccgga	tatcggggag	3900
gtgtacgcgg	atgggtgct	gctggtctgg	aagccgtgg	aatctacgg	ccctgtgacc	3960
tacattgtgc	agtgcagcct	agaaggcggc	agctggacca	cactggcctc	cgacatcttt	4020
gactgctgct	acctgaccag	caagctctcc	cggggtggca	cctacacctt	ccgcacggca	4080
tgtgtcagca	aggcaggaat	gggtccctac	agcagcccct	cggagcaagt	cctcctggga	4140
gggcccagcc	acctggcctc	tgaggaggag	agccaggggc	ggtcagccca	acccctgccc	4200
agcacaaga	ccttcgcatt	ccagacacag	atccagaggg	gccgcttcag	cgtggtgcgg	4260
caatgctggg	agaaggccag	cgggcgggcg	ctggccgcca	agatcatccc	ctaccacccc	4320
aaggacaaga	cagcagtgct	gcgcgaatac	gaggccctca	agggcctgcg	ccaccgcac	4380
ctggcccagc	tgcacgcagc	ctacctcagc	ccccggcacc	tgggtgctcat	cttgagctg	4440
tgtctggtgc	ccgagctgct	cccctgcctg	gccgagaggg	cctcctactc	agaatctgag	4500
gtgaaggact	acctgtggca	gatgttgagt	gccaccagct	acctgcacaa	ccagcacatc	4560
ctgcacctgg	acctgaggtc	cgagaacatg	atcatcaccg	aatacaacct	gctcaaggct	4620
gtggacctgg	gcaatgcaca	gagcctcagc	caggagaagg	tgctgccctc	agacaagttc	4680
aaggactacc	tagagaccat	ggctccagag	ctcctggagg	gccagggggc	tgttccacag	4740
acagacatct	gggccatcgg	tgtgacagcc	ttcatcatgc	tgagcgccga	gtaccgggtg	4800
agcagcgagg	gtgcacgcga	cctgcagaga	ggactgcgca	aggggctggt	ccggctgagc	4860
cgctgtctacg	cggggctgtc	cgggggccc	gtggccttcc	tgcgcagcac	tctgtgcgcc	4920
cagccctggg	gccggccctg	cgcgtccagc	tgcctgcagt	gcccgtggct	aacagaggag	4980
ggcccggcct	gttcgcggcc	cgcgcccgtg	accttcccta	ccgcgcggct	gcgcgtcttc	5040
gtgcgcaatc	gcgagaagag	acgcgcgctg	ctgtacaaga	ggcacaacct	ggcccagggtg	5100
cgctgagggt	cgcgccggcc	acacccttgg	tctcccgcgt	gggggtcgct	gcagacgcgc	5160
caataaaaaac	gcacagccgg	gcgagaaaaa	aaaaaaaaaa	aaaaaaa		5207

<211> 1665  
 <212> PRT  
 <213> Human

<400> 2 -

Met	Gly	Cys	Cys	Arg	Leu	Gly	Cys	Gly	Gly	Cys	Ser	Val	Ala	His	Ser
1				5					10					15	
Val	Ser	Gln	Gly	Leu	Thr	Asn	His	Pro	Ser	Met	Val	Gly	Cys	Gly	Trp
		20						25					30		
His	Pro	Gly	Leu	Cys	Gly	Trp	Gly	Gly	Gly	Leu	His	Ser	Ser	Leu	Pro
		35					40					45			
Ala	Leu	Pro	Gly	Pro	Pro	Ser	Met	Gln	Val	Thr	Ile	Glu	Asp	Val	Gln
		50				55					60				
Ala	Gln	Thr	Gly	Gly	Thr	Ala	Gln	Phe	Glu	Ala	Ile	Ile	Glu	Gly	Asp
65					70				75						80
Pro	Gln	Pro	Ser	Val	Thr	Trp	Tyr	Lys	Asp	Ser	Val	Gln	Leu	Val	Asp
				85					90					95	
Ser	Thr	Arg	Leu	Ser	Gln	Gln	Gln	Glu	Gly	Thr	Thr	Tyr	Ser	Leu	Val
			100					105					110		
Leu	Arg	His	Val	Ala	Ser	Lys	Asp	Ala	Gly	Val	Tyr	Thr	Cys	Leu	Ala
		115					120					125			
Gln	Asn	Thr	Gly	Gly	Gln	Val	Leu	Cys	Lys	Ala	Glu	Leu	Leu	Val	Leu
		130				135						140			
Gly	Gly	Asp	Asn	Glu	Pro	Asp	Ser	Glu	Lys	Gln	Ser	His	Arg	Arg	Lys
145					150					155					160
Leu	His	Ser	Phe	Tyr	Glu	Val	Lys	Glu	Glu	Ile	Gly	Arg	Gly	Val	Phe
			165					170						175	
Gly	Phe	Val	Lys	Arg	Val	Gln	His	Lys	Gly	Asn	Lys	Ile	Leu	Cys	Ala
			180					185					190		
Ala	Lys	Phe	Ile	Pro	Leu	Arg	Ser	Arg	Thr	Arg	Ala	Gln	Ala	Tyr	Arg
		195					200					205			
Glu	Arg	Asp	Ile	Leu	Ala	Ala	Leu	Ser	His	Pro	Leu	Val	Thr	Gly	Leu
		210				215					220				
Leu	Asp	Gln	Phe	Glu	Thr	Arg	Lys	Thr	Leu	Ile	Leu	Ile	Leu	Glu	Leu
225					230					235					240
Cys	Ser	Ser	Glu	Glu	Leu	Leu	Asp	Arg	Leu	Tyr	Arg	Lys	Gly	Val	Val
				245					250					255	
Thr	Glu	Ala	Glu	Val	Lys	Val	Tyr	Ile	Gln	Gln	Leu	Val	Glu	Gly	Leu
			260					265					270		
His	Tyr	Leu	His	Ser	His	Gly	Val	Leu	His	Leu	Asp	Ile	Lys	Pro	Ser
		275					280					285			
Asn	Ile	Leu	Met	Val	His	Pro	Ala	Arg	Glu	Asp	Ile	Lys	Ile	Cys	Asp
		290				295					300				
Phe	Gly	Phe	Ala	Gln	Asn	Ile	Thr	Pro	Ala	Glu	Leu	Gln	Phe	Ser	Gln
305					310					315					320
Tyr	Gly	Ser	Pro	Glu	Phe	Val	Ser	Pro	Glu	Ile	Ile	Gln	Gln	Asn	Pro
				325					330					335	
Val	Ser	Glu	Ala	Ser	Asp	Ile	Trp	Ala	Met	Gly	Val	Ile	Ser	Tyr	Leu
			340					345					350		
Ser	Leu	Thr	Cys	Ser	Ser	Pro	Phe	Ala	Gly	Glu	Ser	Asp	Arg	Ala	Thr
		355					360					365			
Leu	Leu	Asn	Val	Leu	Glu	Gly	Arg	Val	Ser	Trp	Ser	Ser	Pro	Met	Ala
		370				375					380				
Ala	His	Leu	Ser	Glu	Asp	Ala	Lys	Asp	Phe	Ile	Lys	Ala	Thr	Leu	Gln
385					390					395					400
Arg	Ala	Pro	Gln	Ala	Arg	Pro	Ser	Ala	Ala	Gln	Cys	Leu	Ser	His	Pro
				405					410					415	
Trp	Phe	Leu	Lys	Ser	Met	Pro	Ala	Glu	Glu	Ala	His	Phe	Ile	Asn	Thr
			420					425				430			
Lys	Gln	Leu	Lys	Phe	Leu	Leu	Ala	Arg	Ser	Arg	Trp	Gln	Arg	Ser	Leu
		435					440					445			

Met	Ser	Tyr	Lys	Ser	Ile	Leu	Val	Met	Arg	Ser	Ile	Pro	Glu	Leu	Leu
450						455					460				
Arg	Gly	Pro	Pro	Asp	Ser	Pro	Ser	Leu	Gly	Val	Ala	Arg	His	Leu	Cys
465					470					475					480
Arg	Asp	Thr	Gly	Gly	Ser	Ser	Ser	Ser	Ser	Ser	Ser	Ser	Asp	Asn	Glu
				485					490					495	
Leu	Ala	Pro	Phe	Ala	Arg	Ala	Lys	Ser	Leu	Pro	Pro	Ser	Pro	Val	Thr
			500					505					510		
His	Ser	Pro	Leu	Leu	His	Pro	Arg	Gly	Phe	Leu	Arg	Pro	Ser	Ala	Ser
		515					520					525			
Leu	Pro	Glu	Glu	Ala	Glu	Ala	Ser	Glu	Arg	Ser	Thr	Glu	Ala	Pro	Ala
	530					535					540				
Pro	Pro	Ala	Ser	Pro	Glu	Gly	Ala	Gly	Pro	Pro	Ala	Ala	Gln	Gly	Cys
545					550					555					560
Val	Pro	Arg	His	Ser	Val	Ile	Arg	Ser	Leu	Phe	Tyr	His	Gln	Ala	Gly
				565					570					575	
Glu	Ser	Pro	Glu	His	Gly	Ala	Leu	Ala	Pro	Gly	Ser	Arg	Arg	His	Pro
			580					585					590		
Ala	Arg	Arg	Arg	His	Leu	Leu	Lys	Gly	Gly	Tyr	Ile	Ala	Gly	Ala	Leu
			595				600					605			
Pro	Gly	Leu	Arg	Glu	Pro	Leu	Met	Glu	His	Arg	Val	Leu	Glu	Glu	Glu
	610					615					620				
Ala	Ala	Arg	Glu	Glu	Gln	Ala	Thr	Leu	Leu	Ala	Lys	Ala	Pro	Ser	Phe
625					630					635					640
Glu	Thr	Ala	Leu	Arg	Leu	Pro	Ala	Ser	Gly	Thr	His	Leu	Ala	Pro	Gly
				645					650					655	
His	Ser	His	Ser	Leu	Glu	His	Asp	Ser	Pro	Ser	Thr	Pro	Arg	Pro	Ser
			660					665					670		
Ser	Glu	Ala	Cys	Gly	Glu	Ala	Gln	Arg	Leu	Pro	Ser	Ala	Pro	Ser	Gly
		675					680					685			
Gly	Ala	Pro	Ile	Arg	Asp	Met	Gly	His	Pro	Gln	Gly	Ser	Lys	Gln	Leu
	690					695					700				
Pro	Ser	Thr	Gly	Gly	His	Pro	Gly	Thr	Ala	Gln	Pro	Glu	Arg	Pro	Ser
705					710					715					720
Pro	Asp	Ser	Pro	Trp	Gly	Gln	Pro	Ala	Pro	Phe	Cys	His	Pro	Lys	Gln
				725					730					735	
Gly	Ser	Ala	Pro	Gln	Glu	Gly	Cys	Ser	Pro	His	Pro	Ala	Val	Ala	Pro
			740					745					750		
Cys	Pro	Pro	Gly	Ser	Phe	Pro	Pro	Gly	Ser	Cys	Lys	Glu	Ala	Pro	Leu
			755				760					765			
Val	Pro	Ser	Ser	Pro	Phe	Leu	Gly	Gln	Pro	Gln	Ala	Pro	Pro	Ala	Pro
	770					775					780				
Ala	Lys	Ala	Ser	Pro	Pro	Leu	Asp	Ser	Lys	Met	Gly	Pro	Gly	Asp	Ile
785					790					795					800
Ser	Leu	Pro	Gly	Arg	Pro	Lys	Pro	Gly	Pro	Cys	Ser	Ser	Pro	Gly	Ser
				805					810					815	
Ala	Ser	Gln	Ala	Ser	Ser	Ser	Gln	Val	Ser	Ser	Leu	Arg	Val	Gly	Ser
			820					825					830		
Ser	Gln	Val	Gly	Thr	Glu	Pro	Gly	Pro	Ser	Leu	Asp	Ala	Glu	Gly	Trp
			835				840						845		
Thr	Gln	Glu	Ala	Glu	Asp	Leu	Ser	Asp	Ser	Thr	Pro	Thr	Leu	Gln	Arg
	850					855					860				
Pro	Gln	Glu	Gln	Val	Thr	Met	Arg	Lys	Phe	Ser	Leu	Gly	Gly	Arg	Gly
865					870					875					880
Gly	Tyr	Ala	Gly	Val	Ala	Gly	Tyr	Gly	Thr	Phe	Ala	Phe	Gly	Gly	Asp
				885					890					895	
Ala	Gly	Gly	Met	Leu	Gly	Gln	Gly	Pro	Met	Trp	Ala	Arg	Ile	Ala	Trp
			900					905					910		
Ala	Val	Ser	Gln	Ser	Glu	Glu	Glu	Glu	Gln	Glu	Glu	Ala	Arg	Ala	Glu
			915				920					925			
Ser	Gln	Ser	Glu	Glu	Gln	Gln	Glu	Ala	Arg	Ala	Glu	Ser	Pro	Leu	Pro

930		935		940
Gln Val Ser Ala Arg Pro Val Pro Glu Val Gly Arg Ala Pro Thr Arg				
945	950	955		960
Ser Ser Pro Glu Pro Thr Pro Trp Glu Asp Ile Gly Gln Val Ser Leu				
	965	970		975
Val Gln Ile Arg Asp Leu Ser Gly Asp Ala Glu Ala Ala Asp Thr Ile				
	980	985		990
Ser Leu Asp Ile Ser Glu Val Asp Pro Ala Tyr Leu Asn Leu Ser Asp				
	995	1000		1005
Leu Tyr Asp Ile Lys Tyr Leu Pro Phe Glu Phe Met Ile Phe Arg Lys				
	1010	1015		1020
Val Pro Lys Ser Ala Gln Pro Glu Pro Pro Ser Pro Met Ala Glu Glu				
	1025	1030		1035
Glu Leu Ala Glu Phe Pro Glu Pro Thr Trp Pro Trp Pro Gly Glu Leu				
	1045	1050		1055
Gly Pro His Ala Gly Leu Glu Ile Thr Glu Glu Ser Glu Asp Val Asp				
	1060	1065		1070
Ala Leu Leu Ala Glu Ala Ala Val Gly Arg Lys Arg Lys Trp Ser Ser				
	1075	1080		1085
Pro Ser Arg Ser Leu Phe His Phe Pro Gly Arg His Leu Pro Leu Asp				
	1090	1095		1100
Glu Pro Ala Glu Leu Gly Leu Arg Glu Arg Val Lys Ala Ser Val Glu				
	1105	1110		1115
His Ile Ser Arg Ile Leu Lys Gly Arg Pro Glu Gly Leu Glu Lys Glu				
	1125	1130		1135
Gly Pro Pro Arg Lys Lys Pro Gly Leu Ala Ser Phe Arg Leu Ser Gly				
	1140	1145		1150
Leu Lys Ser Trp Asp Arg Ala Pro Thr Phe Leu Arg Glu Leu Ser Asp				
	1155	1160		1165
Glu Thr Val Val Leu Gly Gln Ser Val Thr Leu Ala Cys Gln Val Ser				
	1170	1175		1180
Ala Gln Pro Ala Ala Gln Ala Thr Trp Ser Lys Asp Gly Ala Pro Leu				
	1185	1190		1195
Glu Ser Ser Ser Arg Val Leu Ile Ser Ala Thr Leu Lys Asn Phe Gln				
	1205	1210		1215
Leu Leu Thr Ile Leu Val Val Val Ala Glu Asp Leu Gly Val Tyr Thr				
	1220	1225		1230
Cys Ser Val Ser Asn Ala Leu Gly Thr Val Thr Thr Thr Gly Val Leu				
	1235	1240		1245
Arg Lys Ala Glu Arg Pro Ser Ser Ser Pro Cys Pro Asp Ile Gly Glu				
	1250	1255		1260
Val Tyr Ala Asp Gly Val Leu Leu Val Trp Lys Pro Val Glu Ser Tyr				
	1265	1270		1275
Gly Pro Val Thr Tyr Ile Val Gln Cys Ser Leu Glu Gly Gly Ser Trp				
	1285	1290		1295
Thr Thr Leu Ala Ser Asp Ile Phe Asp Cys Cys Tyr Leu Thr Ser Lys				
	1300	1305		1310
Leu Ser Arg Gly Gly Thr Tyr Thr Phe Arg Thr Ala Cys Val Ser Lys				
	1315	1320		1325
Ala Gly Met Gly Pro Tyr Ser Ser Pro Ser Glu Gln Val Leu Leu Gly				
	1330	1335		1340
Gly Pro Ser His Leu Ala Ser Glu Glu Glu Ser Gln Gly Arg Ser Ala				
	1345	1350		1355
Gln Pro Leu Pro Ser Thr Lys Thr Phe Ala Phe Gln Thr Gln Ile Gln				
	1365	1370		1375
Arg Gly Arg Phe Ser Val Val Arg Gln Cys Trp Glu Lys Ala Ser Gly				
	1380	1385		1390
Arg Ala Leu Ala Ala Lys Ile Ile Pro Tyr His Pro Lys Asp Lys Thr				
	1395	1400		1405
Ala Val Leu Arg Glu Tyr Glu Ala Leu Lys Gly Leu Arg His Pro His				
	1410	1415		1420



Leu Ala Gln Leu His Ala Ala Tyr Leu Ser Pro Arg His Leu Val Leu  
 1425 1430 1435 1440  
 Ile Leu Glu Leu Cys Ser Gly Pro Glu Leu Leu Pro Cys Leu Ala Glu  
 1445 1450 1455  
 Arg Ala Ser Tyr Ser Glu Ser Glu Val Lys Asp Tyr Leu Trp Gln Met  
 1460 1465 1470  
 Leu Ser Ala Thr Gln Tyr Leu His Asn Gln His Ile Leu His Leu Asp  
 1475 1480 1485  
 Leu Arg Ser Glu Asn Met Ile Ile Thr Glu Tyr Asn Leu Leu Lys Val  
 1490 1495 1500  
 Val Asp Leu Gly Asn Ala Gln Ser Leu Ser Gln Glu Lys Val Leu Pro  
 1505 1510 1515 1520  
 Ser Asp Lys Phe Lys Asp Tyr Leu Glu Thr Met Ala Pro Glu Leu Leu  
 1525 1530 1535  
 Glu Gly Gln Gly Ala Val Pro Gln Thr Asp Ile Trp Ala Ile Gly Val  
 1540 1545 1550  
 Thr Ala Phe Ile Met Leu Ser Ala Glu Tyr Pro Val Ser Ser Glu Gly  
 1555 1560 1565  
 Ala Arg Asp Leu Gln Arg Gly Leu Arg Lys Gly Leu Val Arg Leu Ser  
 1570 1575 1580  
 Arg Cys Tyr Ala Gly Leu Ser Gly Gly Ala Val Ala Phe Leu Arg Ser  
 1585 1590 1595 1600  
 Thr Leu Cys Ala Gln Pro Trp Gly Arg Pro Cys Ala Ser Ser Cys Leu  
 1605 1610 1615  
 Gln Cys Pro Trp Leu Thr Glu Glu Gly Pro Ala Cys Ser Arg Pro Ala  
 1620 1625 1630  
 Pro Val Thr Phe Pro Thr Ala Arg Leu Arg Val Phe Val Arg Asn Arg  
 1635 1640 1645  
 Glu Lys Arg Arg Ala Leu Leu Tyr Lys Arg His Asn Leu Ala Gln Val  
 1650 1655 1660  
 Arg  
 1665

<210> 3  
 <211> 5207  
 <212> DNA  
 <213> Human

<400> 3  
 cagcacgagg aactccttct gatcacctgg ccagctgagg tcagagtggg agaggcagtg 60  
 gttccattga aggagtactc ctaactgtca gaagcctggg cggtcaggat ggggtgctgt 120  
 cgcttgggct gcggggggtg ttcagttgcc cacagtgtat ctcagggtct caccaaccat 180  
 ccaagcatgg taggctgtgg ctggcaccca gggttgtgtg gctggggagg tggctctccac 240  
 agttccctcc ctgcccctcc agggccccca tccatgcagg taaccatcga ggatgtgcag 300  
 gcacagacag gcggaacggc ccaattcgag gctatcattg agggcgaccc acagccctcg 360  
 gtgacctggt acaaggacag cgtccagctg gtggacagca cccggcttag ccagcagcaa 420  
 gaaggcacca catactccct ggtgctgagg catgtggcct cgaaggatgc cggcgtttac 480  
 acctgcctgg cccaaaacac tgggtggccag gtgctctgca aggagagct gctggtgctt 540  
 ggggggggaca atgagccgga ctacagagaag caaagccacc ggaggaagct gcactccttc 600  
 tatgaggtca aggaggagat tgggaaggggc gtgtttggct tcgtaaaaag agtcagcac 660  
 aaaggaaaca agatcttgtg cgctgccaaag ttcaccccc tacggagcag aactcgggcc 720  
 caggcataca gggagcgaga catcctggcc gcgctgagcc acccgctggt cagggggctg 780  
 ctggaccagt ttgagaccgg caagaccctc atcctcatcc tggagctgtg ctcatccgag 840  
 gagctgctgg accgcctgta caggaagggc gtggtgacgg agggcgaggt caaggtctac 900  
 atccagcagc tgggtgaggg gctgcactac ctgcacagcc atggcgttct ccacctggac 960  
 ataaagccct ctaacatcct gatggtgcat cctgcccggg aagacattaa aatctgcgac 1020  
 tttggctttg cccagaacat caccacagca gagctgcagt tcagccagta cggctcccct 1080  
 gagttcgtct cccccgagat catccagcag aaccctgtga gcgaagcctc cgacatttgg 1140  
 gccatgggtg tcattctcta cctcagcctg acctgctcat cccatttgc cggcgagagt 1200  
 cagcgtgccca ccctcctgaa cgtcctggag gggcgctgt catggagcag ccccatggct 1260

gcccaccta	ggaagacgc	caaagacttc	atcaaggcta	cgctgcagag	agccccctcag	1320
gccccgccta	gtgcggccca	gtgcctctcc	cacccctggt	tcctgaaatc	catgcctgcg	1380
gaggaggccc	acttcatcaa	caccaagcag	ctcaagttcc	tcctggcccg	aagtcgctgg	1440
cagcgttccc	tgatgagcta	caagtccatc	ctggtgatgc	gctccatccc	tgagctgctg	1500
cggggcccac	ccgacagccc	ctccctcggc	gtagcccggc	acctctgcag	ggacactggt	1560
ggctcctcca	gttctctctc	ctcctctgac	aacgagctcg	ccccatttgc	ccgggctaag	1620
tcactgccac	cctccccggt	gacacactca	ccactgctgc	acccccgggg	cttcctgcgg	1680
ccctcggcc	gcctgcctga	ggaagccgag	gccagtgcgc	gctccaccga	ggccccagct	1740
ccgcctgcat	ctcccgaggg	tgccgggcc	ccggccgccc	agggctgcgt	gccccggcac	1800
agcgtcatcc	gcagcctgtt	ctaccaccag	gcgggtgaga	gccctgagca	cggggccctg	1860
gccccgggga	gcaggcgcca	cccggcccg	cgccggcacc	tgctgaagg	cggctacatt	1920
gcgggggggc	tgccaggcct	gcgcgagcca	ctgatggagc	accgcgtgct	ggaggaggag	1980
gccgccaggg	aggagcaggc	caccctcctg	gccaaagccc	cctcattcga	gactgccctc	2040
cggctgcctg	cctctggcac	ccacttgccc	cctggccaca	gccactccct	ggaacatgac	2100
tctccgagca	cccccgccc	ctcctcgagg	gcctgcgggt	aggcacagcg	actgccttca	2160
gccccctccg	ggggggcccc	tatcagggac	atggggcacc	ctcagggctc	caagcagctt	2220
ccatccactg	gtggccaccc	aggcactgct	cagccagaga	ggccatcccc	ggacagccct	2280
tgggggcagc	cagcccttct	ctgccacccc	aagcagggtt	ctgcccccca	ggagggctgc	2340
agccccccacc	cagcagttgc	cccatgccct	cctggctcct	tccctccagg	atcttgcaaa	2400
gaggccccct	tagtaccctc	aagccccctc	ttgggacagc	cccaggcacc	ccctgccccct	2460
gccaaagcaa	gccccccatt	ggactctaag	atggggcctg	gagacatctc	tcttctggg	2520
aggccaaaac	ccggcccctg	cagttcccca	gggtcagcct	cccaggcgag	ctcttcccaa	2580
gtgagctccc	tcagggtggg	ctcctccag	gtgggcacag	agcctggccc	ctccctggat	2640
gcggagggct	ggaccagga	ggctgaggat	ctgtccgact	ccacaccac	cttgacggg	2700
cctcaggaac	aggtgaccat	gcgcaagttc	tccctgggtg	gtcgcggggg	ctacgcaggc	2760
gtggctggct	atggcacctt	tgctttgggt	ggagatgcag	ggggcatgct	ggggcagggg	2820
cccattgtgg	ccaggatagc	ctgggctgtg	tcccagtcgg	aggaggagg	gcaggaggag	2880
gccagggctg	agtcccagtc	ggaggagcag	caggaggcca	gggctgagag	cccactgccc	2940
caggctcagt	caaggcctgt	gcctgaggtc	ggcagggctc	ccaccaggag	ctctccagag	3000
cccaccccat	gggaggacat	cgggcaggtc	tccttggtgc	agatccggga	cctgtcaggt	3060
gatgcggagg	ggccgcagac	aatatccctg	gacatttccg	aggtggacc	cgccctacct	3120
aacctctcag	acctgtacga	tatcaagtac	ctcccattcg	agtttatgat	cttcaggaaa	3180
gtccccaagt	ccgctcagcc	agagccgccc	tcccccatgg	ctgaggagg	gctggccgag	3240
ttcccgagc	ccacgtggcc	ctggccagg	gaactggg	cccacgcagg	cctggagatc	3300
acagaggagt	cagaggatgt	ggacgcgctg	ctggcagagg	ctgccgtggg	cagggaagcg	3360
aagtggctct	cgccgtcacg	cagcctcttc	cacttccctg	ggaggcacct	gccgtggat	3420
gagcctgcag	agctggggct	gcgtgagaga	gtgaaggcct	ccgtggagca	catctcccgg	3480
atcctgaagg	gcaggccgga	aggtctggag	aaggaggggc	ccccaggaa	gaagccaggc	3540
cttgcttctc	tccggctctc	aggtctgaag	agctgggacc	gagcgccgac	attcctaagg	3600
gagctctcag	atgagactgt	ggtcctgggc	cagtcagtga	cactggcctg	ccagggtgtc	3660
gcccagccag	ctgcccaggc	cacctggagc	aaagacggag	ccccctgga	gagcagcagc	3720
cgtgtcctca	tctctgccac	cctcaagaac	ttccagcttc	tgaccatcct	ggtgggtggg	3780
gctgaggacc	tggtgtgtga	cacctgcagc	gtgagcaatg	cgctggggac	agtgaccacc	3840
acgggcgtcc	tccggaaggc	agagcgcctc	tcactcttcg	catgcccgga	tatcggggag	3900
gtgtacgcgg	atggggtgct	gctggctggt	aagcccgtgg	aatcctacgg	ccctgtgacc	3960
tacattgtgc	agtgcagcct	agaaggcggc	agctggacca	cactggcctc	cgacatcttt	4020
gactgctgct	acctgaccag	caagctctcc	cggggtggca	cctacacctt	ccgcacggca	4080
tgtgtcagca	aggcaggaat	gggtccctac	agcagcccct	cggagcaagt	cctcctggga	4140
gggcccagcc	acctggcctc	tgaggaggag	agccaggggc	ggtcagccca	acccctgccc	4200
agcacaaaaga	ccttcgcatt	ccagacacag	atccagaggg	gccgcttcag	cgtggtgcgg	4260
caatgctggg	agaaggccag	cgggcggg	ctggccgcca	agatcatccc	ctaccacccc	4320
aaggacaaga	cagcagtgct	gcgcgaatac	gaggccctca	agggcctgcg	ccaccgcac	4380
ctggcccagc	tgacgcagc	ctacctcagc	ccccggcacc	tggtgctcat	cttgagctg	4440
tgctctgggc	ccgagctgct	cccctgectg	gccgagaggg	cctcctaact	agaatctgag	4500
gtgaaggact	acctgtggca	gatgttgagt	gccacccagt	acctgcacaa	ccagcacatc	4560
ctgcacctgg	acctgaggtc	cgagaacatg	atcatcaccg	aatacaacct	gctcaaggtc	4620
gtggacctgg	gcaatgcaca	gagcctcagc	caggagaagg	tgctgccctc	agacaagttc	4680
aaggactacc	tagagaccat	ggctccagag	ctcctggagg	gccagggggc	tggtccacag	4740
acagacatct	gggccatcgg	tgtgacagcc	ttcatcatgc	tgagcgccga	gtaccgggtg	4800
agcagcgagg	gtgcacgcga	cctgcagaga	ggactgcgca	aggggctggt	ccggctgagc	4860
cgctgctacg	cggggctgtc	cgggggcgcc	gtggccttcc	tgccgcagc	tctgtgcgcc	4920

cagccctggg gccggccctg cgcgtccagc tgcctgcagt gcccgtaggt aacagaggag 4980  
ggcccgccct gttcgcggcc cgcgcccgtg accttcccta ccgcgcggct gcgcgtcttc 5040  
gtgcgcaatc gcgagaagag acgcgcgctg ctgtacaaga ggcacaacct ggcccagggt 5100  
cgctgagggt cgcgccggcc acacccttgg tctccccgct gggggtcgct gcagacgcgc 5160  
caataaaaac gcacagccgg gcgagaaaaa aaaaaaaaaa aaaaaaa 5207

<210> 4  
<211> 846  
<212> PRT  
<213> Human

<400> 4  
Pro Arg Phe Glu Ser Ile Met Glu Asp Val Glu Val Gly Ala Gly Glu  
1 5 10 15  
Thr Ala Arg Phe Ala Val Val Val Glu Gly Lys Pro Leu Pro Asp Ile  
20 25 30  
Met Trp Tyr Lys Asp Glu Val Leu Leu Thr Glu Ser Ser His Val Ser  
35 40 45  
Phe Val Tyr Glu Glu Asn Glu Cys Ser Leu Val Val Leu Ser Thr Gly  
50 55 60  
Ala Gln Asp Gly Gly Val Tyr Thr Cys Thr Ala Gln Asn Leu Ala Gly  
65 70 75 80  
Glu Val Ser Cys Lys Ala Glu Leu Ala Val His Ser Ala Gln Thr Ala  
85 90 95  
Met Glu Val Glu Gly Val Gly Glu Asp Glu Asp His Arg Gly Arg Arg  
100 105 110  
Leu Ser Asp Phe Tyr Asp Ile His Gln Glu Ile Gly Arg Gly Ala Phe  
115 120 125  
Ser Tyr Leu Arg Arg Ile Val Glu Arg Ser Ser Gly Leu Glu Phe Ala  
130 135 140  
Ala Lys Phe Ile Pro Ser Gln Ala Lys Pro Lys Ala Ser Ala Arg Arg  
145 150 155 160  
Glu Ala Arg Leu Leu Ala Arg Leu Gln His Asp Cys Val Leu Tyr Phe  
165 170 175  
His Glu Ala Phe Glu Arg Arg Arg Gly Leu Val Ile Val Thr Glu Leu  
180 185 190  
Cys Thr Glu Glu Leu Leu Glu Arg Ile Ala Arg Lys Pro Thr Val Cys  
195 200 205  
Glu Ser Glu Ile Arg Ala Tyr Met Arg Gln Val Leu Glu Gly Ile His  
210 215 220  
Tyr Leu His Gln Ser His Val Leu His Leu Asp Val Lys Pro Glu Asn  
225 230 235 240  
Leu Leu Val Trp Asp Gly Ala Ala Gly Glu Gln Gln Val Arg Ile Cys  
245 250 255  
Asp Phe Gly Asn Ala Gln Glu Leu Thr Pro Gly Glu Pro Gln Tyr Cys  
260 265 270  
Gln Tyr Gly Thr Pro Glu Phe Val Ala Pro Glu Ile Val Asn Gln Ser  
275 280 285  
Pro Val Ser Gly Val Thr Asp Ile Trp Pro Val Gly Val Val Ala Phe  
290 295 300  
Leu Cys Leu Thr Gly Ile Ser Pro Phe Val Gly Glu Asn Asp Arg Thr  
305 310 315 320  
Thr Leu Met Asn Ile Arg Asn Tyr Asn Val Ala Phe Glu Glu Thr Thr  
325 330 335  
Phe Leu Ser Leu Ser Arg Glu Ala Arg Gly Phe Leu Ile Lys Val Leu  
340 345 350  
Val Gln Asp Arg Leu Arg Pro Thr Ala Glu Glu Thr Leu Glu His Pro  
355 360 365  
Trp Phe Lys Thr Gln Ala Lys Gly Ala Glu Val Ser Thr Asp His Leu

370		375		380
Lys Leu Phe Leu Ser Arg Arg Arg Trp Gln Arg Ser Gln Ile Ser Tyr				
385		390		395
Lys Cys His Leu Val Leu Arg Pro Ile Pro Glu Leu Leu Arg Ala Pro				400
	405		410	415
Pro Glu Arg Val Trp Val Thr Met Pro Arg Arg Pro Pro Pro Ser Gly				
	420		425	430
Gly Leu Ser Ser Ser Asp Ser Glu Glu Glu Glu Leu Glu Glu Leu				
	435		440	445
Pro Ser Val Pro Arg Pro Leu Gln Pro Glu Phe Ser Gly Ser Arg Val				
	450		455	460
Ser Leu Thr Asp Ile Pro Thr Glu Asp Glu Ala Leu Gly Thr Pro Glu				
465		470		475
Thr Gly Ala Ala Thr Pro Met Asp Trp Gln Glu Gln Gly Arg Ala Pro				
	485		490	495
Ser Gln Asp Gln Glu Ala Pro Ser Pro Glu Ala Leu Pro Ser Pro Gly				
	500		505	510
Gln Glu Pro Ala Ala Gly Ala Ser Pro Arg Arg Gly Glu Leu Arg Arg				
	515		520	525
Gly Ser Ser Ala Glu Ser Ala Leu Pro Arg Ala Gly Pro Arg Glu Leu				
	530		535	540
Gly Arg Gly Leu His Lys Ala Ala Ser Val Glu Leu Pro Gln Arg Arg				
545		550		555
Ser Pro Gly Pro Gly Ala Thr Arg Leu Ala Arg Gly Gly Leu Gly Glu				
	565		570	575
Gly Glu Tyr Ala Gln Arg Leu Gln Ala Leu Arg Gln Arg Leu Leu Arg				
	580		585	590
Gly Gly Pro Glu Asp Gly Lys Val Ser Gly Leu Arg Gly Pro Leu Leu				
	595		600	605
Glu Ser Leu Gly Gly Arg Ala Arg Asp Pro Arg Met Ala Arg Ala Ala				
	610		615	620
Ser Ser Glu Ala Ala Pro His His Gln Pro Pro Leu Glu Asn Arg Gly				
625		630		635
Leu Gln Lys Ser Ser Ser Phe Ser Gln Gly Glu Ala Glu Pro Arg Gly				
	645		650	655
Arg His Arg Arg Ala Gly Ala Pro Leu Glu Ile Pro Val Ala Arg Leu				
	660		665	670
Gly Ala Arg Arg Leu Gln Glu Ser Pro Ser Leu Ser Ala Leu Ser Glu				
	675		680	685
Ala Gln Pro Ser Ser Pro Ala Arg Pro Ser Ala Pro Lys Pro Ser Thr				
	690		695	700
Pro Lys Ser Ala Glu Pro Ser Ala Thr Thr Pro Ser Asp Ala Pro Gln				
705		710		715
Pro Pro Ala Pro Gln Pro Ala Gln Asp Lys Ala Pro Glu Pro Arg Pro				
	725		730	735
Glu Pro Val Arg Ala Ser Lys Pro Ala Pro Pro Pro Gln Ala Leu Gln				
	740		745	750
Thr Leu Ala Leu Pro Leu Thr Pro Tyr Ala Gln Ile Ile Gln Ser Leu				
	755		760	765
Gln Leu Ser Gly His Ala Gln Gly Pro Ser Gln Gly Pro Ala Ala Pro				
	770		775	780
Pro Ser Glu Pro Lys Pro His Ala Ala Val Phe Ala Arg Val Ala Ser				
785		790		795
Pro Pro Pro Gly Ala Pro Glu Lys Arg Val Pro Ser Ala Gly Gly Pro				
	805		810	815
Pro Val Leu Ala Glu Lys Ala Arg Val Pro Thr Val Pro Pro Arg Pro				
	820		825	830
Gly Ser Ser Leu Ser Ser Ser Ile Glu Asn Leu Glu Ser Glu				
	835		840	845

<210> 5  
 <211> 279  
 <212> PRT  
 <213> Human

<400> 5  
 Ser Pro Ala Lys Glu Val Val Ser Ser Pro Gly Ser Ser Pro Arg Ser  
 1 5 10 15  
 Ser Pro Arg Pro Glu Gly Thr Thr Leu Arg Gln Gly Pro Pro Gln Lys  
 20 25 30  
 Pro Tyr Thr Phe Leu Glu Glu Lys Ala Arg Gly Arg Phe Gly Val Val  
 35 40 45  
 Arg Ala Cys Arg Glu Asn Ala Thr Gly Arg Thr Phe Val Ala Lys Ile  
 50 55 60  
 Val Pro Tyr Ala Ala Glu Gly Lys Pro Arg Val Leu Gln Glu Tyr Glu  
 65 70 75 80  
 Val Leu Arg Thr Leu His His Glu Arg Ile Met Ser Leu His Glu Ala  
 85 90 95  
 Tyr Ile Thr Pro Arg Tyr Leu Val Leu Ile Ala Glu Ser Cys Gly Asn  
 100 105 110  
 Arg Glu Leu Leu Cys Gly Leu Ser Asp Arg Phe Arg Tyr Ser Glu Asp  
 115 120 125  
 Asp Val Ala Thr Tyr Met Val Gln Leu Leu Gln Gly Leu Asp Tyr Leu  
 130 135 140  
 His Gly His His Val Leu His Leu Asp Ile Lys Pro Asp Asn Leu Leu  
 145 150 155 160  
 Leu Ala Pro Asp Asn Ala Leu Lys Ile Val Asp Phe Gly Ser Ala Gln  
 165 170 175  
 Pro Tyr Asn Pro Gln Ala Leu Arg Pro Leu Gly His Arg Thr Gly Thr  
 180 185 190  
 Leu Glu Phe Met Ala Pro Glu Met Val Lys Gly Glu Pro Ile Gly Ser  
 195 200 205  
 Ala Thr Asp Ile Trp Gly Ala Gly Val Leu Thr Tyr Ile Met Leu Ser  
 210 215 220  
 Gly Arg Ser Pro Phe Tyr Glu Pro Asp Pro Gln Glu Thr Glu Ala Arg  
 225 230 235 240  
 Ile Val Gly Gly Arg Phe Asp Ala Phe Gln Leu Tyr Pro Asn Thr Ser  
 245 250 255  
 Gln Ser Ala Thr Leu Phe Leu Arg Lys Val Leu Ser Val His Pro Trp  
 260 265 270  
 Ser Arg Pro Ser Ser Cys Leu  
 275

<210> 6  
 <211> 549  
 <212> PRT  
 <213> Human

<400> 6  
 Leu Arg Glu Pro Gly Trp Ala Ala Thr Gly Leu Arg Lys Gly Val Gln  
 1 5 10 15  
 His Ile Phe Arg Val Leu Ser Thr Thr Val Lys Ser Ser Ser Lys Pro  
 20 25 30  
 Ser Pro Pro Ser Glu Pro Val Gln Leu Leu Glu His Gly Pro Thr Leu  
 35 40 45  
 Glu Glu Ala Pro Ala Met Leu Asp Lys Pro Asp Ile Val Tyr Val Val  
 50 55 60  
 Glu Gly Gln Pro Ala Ser Val Thr Val Thr Phe Asn His Val Glu Ala  
 65 70 75 80  
 Gln Val Val Trp Arg Ser Cys Arg Gly Ala Leu Leu Glu Ala Arg Ala



<210> 7  
 <211> 250  
 <212> PRT  
 <213> Human

<400> 7

Tyr	Thr	Phe	Leu	Glu	Glu	Lys	Ala	Arg	Gly	Arg	Phe	Gly	Val	Val	Arg
1				5					10					15	
Ala	Cys	Arg	Glu	Asn	Ala	Thr	Gly	Arg	Thr	Phe	Val	Ala	Lys	Ile	Val
			20					25					30		
Pro	Tyr	Ala	Ala	Glu	Gly	Lys	Pro	Arg	Val	Leu	Gln	Glu	Tyr	Glu	Val
		35					40					45			
Leu	Arg	Thr	Leu	His	His	Glu	Arg	Ile	Met	Ser	Leu	His	Glu	Ala	Tyr
	50					55					60				
Ile	Thr	Pro	Arg	Tyr	Leu	Val	Leu	Ile	Ala	Glu	Ser	Cys	Gly	Asn	Arg
65					70					75					80
Glu	Leu	Leu	Cys	Gly	Leu	Ser	Asp	Arg	Phe	Arg	Tyr	Ser	Glu	Asp	Asp
				85					90					95	
Val	Ala	Thr	Tyr	Met	Val	Gln	Leu	Leu	Gln	Gly	Leu	Asp	Tyr	Leu	His
			100						105					110	
Gly	His	His	Val	Leu	His	Leu	Asp	Ile	Lys	Pro	Asp	Asn	Leu	Leu	Leu
		115					120					125			
Ala	Pro	Asp	Asn	Ala	Leu	Lys	Ile	Val	Asp	Phe	Gly	Ser	Ala	Gln	Pro
	130					135					140				
Tyr	Asn	Pro	Gln	Ala	Leu	Arg	Pro	Leu	Gly	His	Arg	Thr	Gly	Thr	Leu
145					150					155					160
Glu	Phe	Met	Ala	Pro	Glu	Met	Val	Lys	Gly	Glu	Pro	Ile	Gly	Ser	Ala
				165					170					175	
Thr	Asp	Ile	Trp	Gly	Ala	Gly	Val	Leu	Thr	Tyr	Ile	Met	Leu	Ser	Gly
			180					185					190		
Arg	Ser	Pro	Phe	Tyr	Glu	Pro	Asp	Pro	Gln	Glu	Thr	Glu	Ala	Arg	Ile
		195					200					205			
Val	Gly	Gly	Arg	Phe	Asp	Ala	Phe	Gln	Leu	Tyr	Pro	Asn	Thr	Ser	Gln
	210					215					220				
Ser	Ala	Thr	Leu	Phe	Leu	Arg	Lys	Val	Leu	Ser	Val	His	Pro	Trp	Ser
225					230					235					240
Arg	Pro	Ser	Ser	Cys	Leu	Ser	Val	Cys	His						
				245					250						

<210> 8  
 <211> 245  
 <212> PRT  
 <213> Human

<400> 8

Pro	Arg	Lys	Asp	Lys	Gly	Leu	Ser	Pro	Pro	Asn	Leu	Ser	Ala	Ser	Val
1				5					10					15	
Gln	Glu	Glu	Leu	Gly	His	Gln	Tyr	Val	Arg	Ser	Glu	Ser	Asp	Phe	Pro
			20					25					30		
Pro	Val	Phe	His	Ile	Lys	Leu	Lys	Asp	Gln	Val	Leu	Leu	Glu	Gly	Glu
		35					40					45			
Ala	Ala	Thr	Leu	Leu	Cys	Leu	Pro	Ala	Ala	Cys	Pro	Ala	Pro	His	Ile
	50					55					60				
Ser	Trp	Met	Lys	Asp	Lys	Lys	Ser	Leu	Arg	Ser	Glu	Pro	Ser	Val	Ile
65					70					75					80
Ile	Val	Ser	Cys	Lys	Asp	Gly	Arg	Gln	Leu	Leu	Ser	Ile	Pro	Arg	Ala
				85					90					95	
Gly	Lys	Arg	His	Ala	Gly	Leu	Tyr	Glu	Cys	Ser	Ala	Thr	Asn	Val	Leu
			100					105					110		
Gly	Ser	Ile	Thr	Ser	Ser	Cys	Thr	Val	Ala	Val	Ala	Arg	Val	Pro	Gly





Thr Leu Thr Trp Asn Pro Pro Arg Ser Leu Asp Met Ala Ile Asp Pro  
 115 120 125  
 Asp Ser Leu Thr Tyr Thr Val Gln His Gln Val Leu Gly Ser Asp Gln  
 130 135 140  
 Trp Thr Ala Leu Val Thr Gly Leu Arg Glu Pro Gly Trp Ala Ala Thr  
 145 150 155 160  
 Gly Leu Arg Lys Gly Val Gln His Ile Phe Arg Val Leu Ser Thr Thr  
 165 170 175  
 Val Lys Ser Ser Lys Pro Ser Pro Ser Glu Pro Val Gln Leu  
 180 185 190  
 Leu Glu His Gly Pro Thr  
 195

<210> 11  
 <211> 101  
 <212> PRT  
 <213> Human

<400> 11  
 Ala Pro Leu Phe Thr Arg Leu Leu Glu Asp Val Glu Val Leu Glu Gly  
 1 5 10 15  
 Arg Ala Ala Arg Phe Asp Cys Lys Ile Ser Gly Thr Pro Pro Val  
 20 25 30  
 Val Thr Trp Thr His Phe Gly Cys Pro Met Glu Glu Ser Glu Asn Leu  
 35 40 45  
 Arg Leu Arg Gln Asp Gly Gly Leu His Ser Leu His Ile Ala His Val  
 50 55 60  
 Gly Ser Glu Asp Glu Gly Leu Tyr Ala Val Ser Ala Val Asn Thr His  
 65 70 75 80  
 Gly Gln Ala His Cys Ser Ala Gln Leu Tyr Val Glu Glu Pro Arg Thr  
 85 90 95  
 Ala Ala Ser Gly Pro  
 100

<210> 12  
 <211> 195  
 <212> PRT  
 <213> Human

<400> 12  
 Arg Gly Thr Gln Asp Ser Ser Ala Val Pro Ser Ala Ala His Gln Glu  
 1 5 10 15  
 Ala Pro Val Thr Ser Arg Pro Ala Arg Ala Arg Pro Pro Asp Ser Pro  
 20 25 30  
 Thr Ser Leu Ala Pro Pro Leu Ala Pro Ala Ala Pro Thr Pro Pro Ser  
 35 40 45  
 Val Thr Val Ser Pro Ser Ser Pro Pro Thr Pro Pro Ser Gln Ala Leu  
 50 55 60  
 Ser Ser Leu Lys Ala Val Gly Pro Pro Pro Gln Thr Pro Pro Arg Arg  
 65 70 75 80  
 His Arg Gly Leu Gln Ala Ala Arg Pro Ala Glu Pro Thr Leu Pro Ser  
 85 90 95  
 Thr His Val Thr Pro Ser Glu Pro Lys Pro Phe Val Leu Asp Thr Gly  
 100 105 110  
 Thr Pro Ile Pro Ala Ser Thr Pro Gln Gly Val Lys Pro Val Ser Ser  
 115 120 125  
 Ser Thr Pro Val Tyr Val Val Thr Ser Phe Val Ser Ala Pro Pro Ala  
 130 135 140  
 Pro Glu Pro Pro Ala Pro Glu Pro Pro Pro Glu Pro Thr Lys Val Thr

145                      150                      155                      160  
 Val Gln Ser Leu Ser Pro Ala Lys Glu Val Val Ser Ser Pro Gly Ser  
                                  165                      170                      175  
 Ser Pro Arg Ser Ser Pro Arg Pro Glu Gly Thr Thr Leu Arg Gln Gly  
                                  180                      185                      190  
 Pro Pro Gln  
                                  195

<210> 13  
 <211> 90  
 <212> PRT  
 <213> Human

<400> 13  
 Pro Asp Phe Leu Arg Pro Leu Gln Asp Leu Glu Val Gly Leu Ala Lys  
   1                                 5                                 10                                 15  
 Glu Ala Met Leu Glu Cys Gln Val Thr Gly Leu Pro Tyr Pro Thr Ile  
                                  20                                 25                                 30  
 Ser Trp Phe His Asn Gly His Arg Ile Gln Ser Ser Asp Asp Arg Arg  
                                  35                                 40                                 45  
 Met Thr Gln Tyr Arg Asp Val His Arg Leu Val Phe Pro Ala Val Gly  
                                  50                                 55                                 60  
 Pro Gln His Ala Gly Val Tyr Lys Ser Val Ile Ala Asn Lys Leu Gly  
   65                                 70                                 75                                 80  
 Lys Ala Ala Cys Tyr Ala His Leu Tyr Val  
                                  85                                 90

<210> 14  
 <211> 414  
 <212> PRT  
 <213> Human

<400> 14  
 Pro Pro Glu Phe Val Ile Pro Leu Ser Glu Val Thr Cys Glu Thr Gly  
   1                                 5                                 10                                 15  
 Glu Thr Val Val Leu Arg Cys Arg Val Cys Gly Arg Pro Lys Ala Ser  
                                  20                                 25                                 30  
 Ile Thr Trp Lys Gly Pro Glu His Asn Thr Leu Asn Asn Asp Gly His  
                                  35                                 40                                 45  
 Tyr Ser Ile Ser Tyr Ser Asp Leu Gly Glu Ala Thr Leu Lys Ile Val  
                                  50                                 55                                 60  
 Gly Val Thr Thr Glu Asp Asp Gly Ile Tyr Thr Cys Ile Ala Val Asn  
   65                                 70                                 75                                 80  
 Asp Met Gly Ser Ala Ser Ser Ser Ala Ser Leu Arg Val Leu Gly Pro  
                                  85                                 90                                 95  
 Gly Met Asp Gly Ile Met Val Thr Trp Lys Asp Asn Phe Asp Ser Phe  
                                  100                                 105                                 110  
 Tyr Ser Glu Val Ala Glu Leu Gly Arg Gly Arg Phe Ser Val Val Lys  
                                  115                                 120                                 125  
 Lys Cys Asp Gln Lys Gly Thr Lys Arg Ala Val Ala Thr Lys Phe Val  
                                  130                                 135                                 140  
 Asn Lys Lys Leu Met Lys Arg Asp Gln Val Thr His Glu Leu Gly Ile  
   145                                 150                                 155                                 160  
 Leu Gln Ser Leu Gln His Pro Leu Leu Val Gly Leu Leu Asp Thr Phe  
                                  165                                 170                                 175  
 Glu Thr Pro Thr Ser Tyr Ile Leu Val Leu Glu Met Ala Asp Gln Gly  
                                  180                                 185                                 190  
 Arg Leu Leu Asp Cys Val Val Arg Trp Gly Ser Leu Thr Glu Gly Lys  
                                  195                                 200                                 205

Ile	Arg	Ala	His	Leu	Gly	Glu	Val	Leu	Glu	Ala	Val	Arg	Tyr	Leu	His
210						215					220				
Asn	Cys	Arg	Ile	Ala	His	Leu	Asp	Leu	Lys	Pro	Glu	Asn	Ile	Leu	Val
225					230					235					240
Asp	Glu	Ser	Leu	Ala	Lys	Pro	Thr	Ile	Lys	Leu	Ala	Asp	Phe	Gly	Asp
				245					250					255	
Ala	Val	Gln	Leu	Asn	Thr	Thr	Tyr	Tyr	Ile	His	Gln	Leu	Leu	Gly	Asn
			260					265					270		
Pro	Glu	Phe	Ala	Ala	Pro	Glu	Ile	Ile	Leu	Gly	Asn	Pro	Val	Ser	Leu
	275						280					285			
Thr	Ser	Asp	Thr	Trp	Ser	Val	Gly	Val	Leu	Thr	Tyr	Val	Leu	Leu	Ser
	290					295					300				
Gly	Val	Ser	Pro	Phe	Leu	Asp	Asp	Ser	Val	Glu	Glu	Thr	Cys	Leu	Asn
305					310					315					320
Ile	Cys	Arg	Leu	Asp	Phe	Ser	Phe	Pro	Asp	Asp	Tyr	Phe	Lys	Gly	Val
				325					330					335	
Ser	Gln	Lys	Ala	Lys	Glu	Phe	Val	Cys	Phe	Leu	Leu	Gln	Glu	Asp	Pro
			340					345					350		
Ala	Lys	Arg	Pro	Ser	Ala	Ala	Leu	Ala	Leu	Gln	Glu	Gln	Trp	Leu	Gln
	355						360					365			
Ala	Gly	Asn	Gly	Arg	Ser	Thr	Gly	Val	Leu	Asp	Thr	Ser	Arg	Leu	Thr
	370					375					380				
Ser	Phe	Ile	Glu	Arg	Arg	Lys	His	Gln	Asn	Asp	Val	Arg	Pro	Ile	Arg
385					390					395					400
Ser	Ile	Lys	Asn	Phe	Leu	Gln	Ser	Arg	Leu	Leu	Pro	Arg	Val		
			405						410						

<210> 15  
 <211> 274  
 <212> PRT  
 <213> Human

<400> 15															
Glu	Leu	Gly	Arg	Gly	Arg	Phe	Ser	Val	Val	Lys	Lys	Cys	Asp	Gln	Lys
1				5					10					15	
Gly	Thr	Lys	Arg	Ala	Val	Ala	Thr	Lys	Phe	Val	Asn	Lys	Lys	Leu	Met
		20						25					30		
Lys	Arg	Asp	Gln	Val	Thr	His	Glu	Leu	Gly	Ile	Leu	Gln	Ser	Leu	Gln
		35					40					45			
His	Pro	Leu	Leu	Val	Gly	Leu	Leu	Asp	Thr	Phe	Glu	Thr	Pro	Thr	Ser
	50					55					60				
Tyr	Ile	Leu	Val	Leu	Glu	Met	Ala	Asp	Gln	Gly	Arg	Leu	Leu	Asp	Cys
65					70					75					80
Val	Val	Arg	Trp	Gly	Ser	Leu	Thr	Glu	Gly	Lys	Ile	Arg	Ala	His	Leu
				85					90					95	
Gly	Glu	Val	Leu	Glu	Ala	Val	Arg	Tyr	Leu	His	Asn	Cys	Arg	Ile	Ala
		100						105					110		
His	Leu	Asp	Leu	Lys	Pro	Glu	Asn	Ile	Leu	Val	Asp	Glu	Ser	Leu	Ala
	115						120					125			
Lys	Pro	Thr	Ile	Lys	Leu	Ala	Asp	Phe	Gly	Asp	Ala	Val	Gln	Leu	Asn
	130					135					140				
Thr	Thr	Tyr	Tyr	Ile	His	Gln	Leu	Leu	Gly	Asn	Pro	Glu	Phe	Ala	Ala
145					150					155					160
Pro	Glu	Ile	Ile	Leu	Gly	Asn	Pro	Val	Ser	Leu	Thr	Ser	Asp	Thr	Trp
				165					170					175	
Ser	Val	Gly	Val	Leu	Thr	Tyr	Val	Leu	Leu	Ser	Gly	Val	Ser	Pro	Phe
		180						185					190		
Leu	Asp	Asp	Ser	Val	Glu	Glu	Thr	Cys	Leu	Asn	Ile	Cys	Arg	Leu	Asp
	195						200					205			
Phe	Ser	Phe	Pro	Asp	Asp	Tyr	Phe	Lys	Gly	Val	Ser	Gln	Lys	Ala	Lys

210	215	220
Glu Phe Val Cys Phe Leu	Leu Gln Glu Asp Pro	Ala Lys Arg Pro Ser
225	230	235
Ala Ala Leu Ala Leu Gln	Glu Gln Trp Leu Gln	Ala Gly Asn Gly Arg
	245	250
Ser Thr Gly Val Leu Asp	Thr Ser Arg Leu Thr	Ser Phe Ile Glu Arg
	260	265
		270
Arg Lys		

<210> 16  
 <211> 141  
 <212> PRT  
 <213> Human

<400> 16
Gly Lys Arg Glu Gly Lys Leu Glu Asn Gly Tyr Arg Lys Ser Arg Glu
1 5 10 15
Gly Leu Ser Asn Lys Val Ser Val Lys Leu Leu Asn Pro Asn Tyr Ile
20 25 30
Tyr Asp Val Pro Pro Glu Phe Val Ile Pro Leu Ser Glu Val Thr Cys
35 40 45
Glu Thr Gly Glu Thr Val Val Leu Arg Cys Arg Val Cys Gly Arg Pro
50 55 60
Lys Ala Ser Ile Thr Trp Lys Gly Pro Glu His Asn Thr Leu Asn Asn
65 70 75 80
Asp Gly His Tyr Ser Ile Ser Tyr Ser Asp Leu Gly Glu Ala Thr Leu
85 90 95
Lys Ile Val Gly Val Thr Thr Glu Asp Asp Gly Ile Tyr Thr Cys Ile
100 105 110
Ala Val Asn Asp Met Gly Ser Ala Ser Ser Ser Ala Ser Leu Arg Val
115 120 125
Leu Gly Pro Gly Met Asp Gly Ile Met Val Thr Trp Lys
130 135 140

<210> 17  
 <211> 196  
 <212> PRT  
 <213> Human

<400> 17
Gly Gly Ala Pro Ser Gly Gly Ser Gly His Ser Gly Gly Pro Ser Ser
1 5 10 15
Cys Gly Gly Ala Pro Ser Thr Ser Arg Ser Arg Pro Ser Arg Ile Pro
20 25 30
Gln Pro Val Arg His His Pro Pro Val Leu Val Ser Ser Ala Ala Ser
35 40 45
Ser Gln Ala Glu Ala Asp Lys Met Ser Gly Thr Ser Thr Pro Gly Pro
50 55 60
Ser Leu Pro Pro Pro Gly Ala Ala Pro Glu Ala Gly Pro Ser Ala Pro
65 70 75 80
Ser Arg Arg Pro Pro Gly Ala Asp Ala Glu Gly Ser Glu Arg Glu Ala
85 90 95
Glu Pro Ile Pro Lys Met Lys Val Leu Glu Ser Pro Arg Lys Gly Ala
100 105 110
Ala Asn Ala Ser Gly Ser Ser Pro Asp Ala Pro Ala Lys Asp Ala Arg
115 120 125
Ala Ser Leu Gly Thr Leu Pro Leu Gly Lys Pro Arg Ala Gly Ala Ala
130 135 140

Ser Pro Leu Asn Ser Pro Leu Ser Ser Ala Val Pro Ser Leu Gly Lys  
 145 150 155 160  
 Glu Pro Phe Pro Pro Ser Ser Pro Leu Gln Lys Gly Gly Ser Phe Trp  
 165 170 175  
 Ser Ser Ile Pro Ala Ser Pro Ala Ser Arg Pro Gly Ser Phe Thr Phe  
 180 185 190  
 Pro Gly Asp Ser  
 195

<210> 18  
 <211> 298  
 <212> PRT  
 <213> Human

<400> 18  
 Gln Lys Val Ser Asp Phe Tyr Asp Ile Glu Glu Arg Leu Gly Ser Gly  
 1 5 10 15  
 Lys Phe Gly Gln Val Phe Arg Leu Val Glu Lys Lys Thr Arg Lys Val  
 20 25 30  
 Trp Ala Gly Lys Phe Phe Lys Ala Tyr Ser Ala Lys Glu Lys Glu Asn  
 35 40 45  
 Ile Arg Gln Glu Ile Ser Ile Met Asn Cys Leu His His Pro Lys Leu  
 50 55 60  
 Val Gln Cys Val Asp Ala Phe Glu Glu Lys Ala Asn Ile Val Met Val  
 65 70 75 80  
 Leu Glu Ile Val Ser Gly Gly Glu Leu Phe Glu Arg Ile Ile Asp Glu  
 85 90 95  
 Asp Phe Glu Leu Thr Glu Arg Glu Cys Ile Lys Tyr Met Arg Gln Ile  
 100 105 110  
 Ser Glu Gly Val Glu Tyr Ile His Lys Gln Gly Ile Val His Leu Asp  
 115 120 125  
 Leu Lys Pro Glu Asn Ile Met Cys Val Asn Lys Thr Gly Thr Arg Ile  
 130 135 140  
 Lys Leu Ile Asp Phe Gly Leu Ala Arg Arg Leu Glu Asn Ala Gly Ser  
 145 150 155 160  
 Leu Lys Val Leu Phe Gly Thr Pro Glu Phe Val Ala Pro Glu Val Ile  
 165 170 175  
 Asn Tyr Glu Pro Ile Ser Tyr Ala Thr Asp Met Trp Ser Ile Gly Val  
 180 185 190  
 Ile Cys Tyr Ile Leu Val Ser Gly Leu Ser Pro Phe Met Gly Asp Asn  
 195 200 205  
 Asp Asn Glu Thr Leu Ala Asn Val Thr Ser Ala Thr Trp Asp Phe Asp  
 210 215 220  
 Asp Glu Ala Phe Asp Glu Ile Ser Asp Asp Ala Lys Asp Phe Ile Ser  
 225 230 235 240  
 Asn Leu Leu Lys Lys Asp Met Lys Asn Arg Leu Asp Cys Thr Gln Cys  
 245 250 255  
 Leu Gln His Pro Trp Leu Met Lys Asp Thr Lys Asn Met Glu Ala Lys  
 260 265 270  
 Lys Leu Ser Lys Asp Arg Met Lys Lys Tyr Met Ala Arg Arg Lys Trp  
 275 280 285  
 Gln Lys Thr Gly Asn Ala Val Arg Ala Ile  
 290 295

<210> 19  
 <211> 508  
 <212> PRT  
 <213> Human

<400> 19

Gly	Thr	Glu	Ser	Asp	Ala	Thr	Val	Lys	Lys	Lys	Pro	Ala	Pro	Lys	Thr
1				5					10					15	
Pro	Pro	Lys	Ala	Ala	Met	Pro	Pro	Gln	Ile	Ile	Gln	Phe	Pro	Glu	Asp
			20					25					30		
Gln	Lys	Val	Arg	Ala	Gly	Glu	Ser	Val	Glu	Leu	Phe	Gly	Lys	Val	Thr
		35					40					45			
Gly	Thr	Gln	Pro	Ile	Thr	Cys	Thr	Trp	Met	Lys	Phe	Arg	Lys	Gln	Ile
	50					55					60				
Gln	Asp	Ser	Glu	His	Ile	Lys	Val	Glu	Asn	Ser	Glu	Asn	Gly	Ser	Lys
65					70					75					80
Leu	Thr	Ile	Leu	Ala	Ala	Arg	Gln	Glu	His	Cys	Gly	Cys	Tyr	Thr	Leu
				85					90					95	
Leu	Val	Glu	Asn	Lys	Leu	Gly	Ser	Arg	Gln	Ala	Gln	Val	Asn	Leu	Thr
			100					105					110		
Val	Val	Asp	Lys	Pro	Asp	Pro	Pro	Ala	Gly	Thr	Pro	Cys	Ala	Ser	Asp
		115					120					125			
Ile	Arg	Ser	Ser	Ser	Leu	Thr	Leu	Ser	Trp	Tyr	Gly	Ser	Ser	Tyr	Asp
	130					135					140				
Gly	Gly	Ser	Ala	Val	Gln	Ser	Tyr	Ser	Ile	Glu	Ile	Trp	Asp	Ser	Ala
145					150					155					160
Asn	Lys	Thr	Trp	Lys	Glu	Leu	Ala	Thr	Cys	Arg	Ser	Thr	Ser	Phe	Asn
				165					170					175	
Val	Gln	Asp	Leu	Leu	Pro	Asp	His	Glu	Tyr	Lys	Phe	Arg	Val	Arg	Ala
			180					185					190		
Ile	Asn	Val	Tyr	Gly	Thr	Ser	Glu	Pro	Ser	Gln	Glu	Ser	Glu	Leu	Thr
	195						200					205			
Thr	Val	Gly	Glu	Lys	Pro	Glu	Glu	Pro	Lys	Met	Lys	Trp	Arg	Cys	Gln
	210					215					220				
Thr	Asp	Asp	Glu	Lys	Glu	Pro	Glu	Val	Asp	Tyr	Arg	Thr	Val	Thr	Ile
225					230					235					240
Asn	Thr	Glu	Gln	Lys	Val	Ser	Asp	Phe	Tyr	Asp	Ile	Glu	Glu	Arg	Leu
				245					250					255	
Gly	Ser	Gly	Lys	Phe	Gly	Gln	Val	Phe	Arg	Leu	Val	Glu	Lys	Lys	Thr
			260					265					270		
Arg	Lys	Val	Trp	Ala	Gly	Lys	Phe	Phe	Lys	Ala	Tyr	Ser	Ala	Lys	Glu
		275					280					285			
Lys	Glu	Asn	Ile	Arg	Gln	Glu	Ile	Ser	Ile	Met	Asn	Cys	Leu	His	His
	290					295					300				
Pro	Lys	Leu	Val	Gln	Cys	Val	Asp	Ala	Phe	Glu	Glu	Lys	Ala	Asn	Ile
305					310					315					320
Val	Met	Val	Leu	Glu	Ile	Val	Ser	Gly	Gly	Glu	Leu	Phe	Glu	Arg	Ile
				325					330					335	
Ile	Asp	Glu	Asp	Phe	Glu	Leu	Thr	Glu	Arg	Glu	Cys	Ile	Lys	Tyr	Met
			340					345					350		
Arg	Gln	Ile	Ser	Glu	Gly	Val	Glu	Tyr	Ile	His	Lys	Gln	Gly	Ile	Val
		355					360					365			
His	Leu	Asp	Leu	Lys	Pro	Glu	Asn	Ile	Met	Cys	Val	Asn	Lys	Thr	Gly
	370					375					380				
Thr	Arg	Ile	Lys	Leu	Ile	Asp	Phe	Gly	Leu	Ala	Arg	Arg	Leu	Glu	Asn
385					390					395					400
Ala	Gly	Ser	Leu	Lys	Val	Leu	Phe	Gly	Thr	Pro	Glu	Phe	Val	Ala	Pro
				405					410					415	
Glu	Val	Ile	Asn	Tyr	Glu	Pro	Ile	Ser	Tyr	Ala	Thr	Asp	Met	Trp	Ser
			420					425					430		
Ile	Gly	Val	Ile	Cys	Tyr	Ile	Leu	Val	Ser	Gly	Leu	Ser	Pro	Phe	Met
		435					440					445			
Gly	Asp	Asn	Asp	Asn	Glu	Thr	Leu	Ala	Asn	Val	Thr	Ser	Ala	Thr	Trp
	450					455					460				
Asp	Phe	Asp	Asp	Glu	Ala	Phe	Asp	Glu	Ile	Ser	Asp	Asp	Ala	Lys	Asp
465					470					475					480

Phe Ile Ser Asn Leu Leu Lys Lys Asp Met Lys Asn Arg Leu Asp Cys  
485 490 495  
Thr Gln Cys Leu Gln His Pro Trp Leu Met Lys Asp  
500 505

<210> 20  
<211> 106  
<212> PRT  
<213> Human

<400> 20  
Pro Tyr Phe Ser Lys Thr Ile Arg Asp Leu Glu Val Val Glu Gly Ser  
1 5 10 15  
Ala Ala Arg Phe Asp Cys Lys Ile Glu Gly Tyr Pro Asp Pro Glu Val  
20 25 30  
Val Trp Phe Lys Asp Asp Gln Ser Ile Arg Glu Ser Arg His Phe Gln  
35 40 45  
Ile Asp Tyr Asp Glu Asp Gly Asn Cys Ser Leu Ile Ile Ser Asp Val  
50 55 60  
Cys Gly Asp Asp Asp Ala Lys Tyr Thr Cys Lys Ala Val Asn Ser Leu  
65 70 75 80  
Gly Glu Ala Thr Cys Thr Ala Glu Leu Ile Val Glu Thr Met Glu Glu  
85 90 95  
Gly Glu Gly Glu Gly Glu Glu Glu Glu  
100 105

<210> 21  
<211> 96  
<212> PRT  
<213> Human

<400> 21  
Pro Pro Lys Phe Ala Thr Lys Leu Gly Arg Val Val Val Lys Glu Gly  
1 5 10 15  
Gln Met Gly Arg Phe Ser Cys Lys Ile Thr Gly Arg Pro Gln Pro Gln  
20 25 30  
Val Thr Trp Leu Lys Gly Asn Val Pro Leu Gln Pro Ser Ala Arg Val  
35 40 45  
Ser Val Ser Glu Lys Asn Gly Met Gln Val Leu Glu Ile His Gly Val  
50 55 60  
Asn Gln Asp Asp Val Gly Val Tyr Thr Cys Leu Val Val Asn Gly Ser  
65 70 75 80  
Gly Lys Ala Ser Met Ser Ala Glu Leu Ser Ile Gln Gly Leu Asp Ser  
85 90 95

<210> 22  
<211> 96  
<212> PRT  
<213> Human

<400> 22  
Pro Pro Lys Phe Ala Thr Lys Leu Gly Arg Val Val Val Lys Glu Gly  
1 5 10 15  
Gln Met Gly Arg Phe Ser Cys Lys Ile Thr Gly Arg Pro Gln Pro Gln  
20 25 30  
Val Thr Trp Leu Lys Gly Asn Val Pro Leu Gln Pro Ser Ala Arg Val  
35 40 45  
Ser Val Ser Glu Lys Asn Gly Met Gln Val Leu Glu Ile His Gly Val

50		55		60											
Asn	Gln	Asp	Asp	Val	Gly	Val	Tyr	Thr	Cys	Leu	Val	Val	Asn	Gly	Ser
65				70						75					80
Gly	Lys	Ala	Ser	Met	Ser	Ala	Glu	Leu	Ser	Ile	Gln	Gly	Leu	Asp	Ser
				85					90					95	

<210> 23  
 <211> 88  
 <212> PRT  
 <213> Human

<400> 23

Pro	Lys	Phe	Ala	Thr	Lys	Leu	Gly	Arg	Val	Val	Val	Lys	Glu	Gly	Gln
1				5					10					15	
Met	Gly	Arg	Phe	Ser	Cys	Lys	Ile	Thr	Gly	Arg	Pro	Gln	Pro	Gln	Val
			20					25				30			
Thr	Trp	Leu	Lys	Gly	Asn	Val	Pro	Leu	Gln	Pro	Ser	Ala	Arg	Val	Ser
		35				40						45			
Val	Ser	Glu	Lys	Asn	Gly	Met	Gln	Val	Leu	Glu	Ile	His	Gly	Val	Asn
	50				55					60					
Gln	Asp	Asp	Val	Gly	Val	Tyr	Thr	Cys	Leu	Val	Val	Asn	Gly	Ser	Gly
65				70					75						80
Lys	Ala	Ser	Met	Ser	Ala	Glu	Leu								
				85											

<210> 24  
 <211> 94  
 <212> PRT  
 <213> Human

<400> 24

Ala	Pro	Ser	Phe	Ser	Ser	Val	Leu	Lys	Asp	Cys	Ala	Val	Ile	Glu	Gly
1				5					10					15	
Gln	Asp	Phe	Val	Leu	Gln	Cys	Ser	Val	Arg	Gly	Thr	Pro	Val	Pro	Arg
			20					25				30			
Ile	Thr	Trp	Leu	Leu	Asn	Gly	Gln	Pro	Ile	Gln	Tyr	Ala	Arg	Ser	Thr
		35				40						45			
Cys	Glu	Ala	Gly	Val	Ala	Glu	Leu	His	Ile	Gln	Asp	Ala	Leu	Pro	Glu
	50				55					60					
Asp	His	Gly	Thr	Tyr	Thr	Cys	Leu	Ala	Glu	Asn	Ala	Leu	Gly	Gln	Val
65				70					75						80
Ser	Cys	Ser	Ala	Trp	Val	Thr	Val	His	Glu	Lys	Lys	Ser	Ser		
				85				90							

<210> 25  
 <211> 112  
 <212> PRT  
 <213> Human

<400> 25

Lys	Lys	Ser	Ser	Arg	Lys	Ser	Glu	Tyr	Leu	Leu	Pro	Val	Ala	Pro	Ser
1				5					10					15	
Lys	Pro	Thr	Ala	Pro	Ile	Phe	Leu	Gln	Gly	Leu	Ser	Asp	Leu	Lys	Val
			20					25				30			
Met	Asp	Gly	Ser	Gln	Val	Thr	Met	Thr	Val	Gln	Val	Ser	Gly	Asn	Pro
		35				40						45			
Pro	Pro	Glu	Val	Ile	Trp	Leu	His	Asn	Gly	Asn	Glu	Ile	Gln	Glu	Ser
	50					55					60				



Glu Asp Phe His Phe Glu Gln Arg Gly Thr Gln His Ser Leu Trp Ile  
65 70 75 80  
Gln Glu Val Phe Pro Glu Asp Thr Gly Thr Tyr Thr Cys Glu Ala Trp  
85 90 95  
Asn Ser Ala Gly Glu Val Arg Thr Gln Ala Val Leu Thr Val Gln Glu  
100 105 110

<210> 26  
<211> 100  
<212> PRT  
<213> Human

<400> 26  
Ser Met Pro Leu Thr Glu Ala Pro Ala Phe Ile Leu Pro Pro Arg Asn  
1 5 10 15  
Leu Cys Ile Lys Glu Gly Ala Thr Ala Lys Phe Glu Gly Arg Val Arg  
20 25 30  
Gly Tyr Pro Glu Pro Gln Val Thr Trp His Arg Asn Gly Gln Pro Ile  
35 40 45  
Thr Ser Gly Gly Arg Phe Leu Leu Asp Cys Gly Ile Arg Gly Thr Phe  
50 55 60  
Ser Leu Val Ile His Ala Val His Glu Glu Asp Arg Gly Lys Tyr Thr  
65 70 75 80  
Cys Glu Ala Thr Asn Gly Ser Gly Ala Arg Gln Val Thr Val Glu Leu  
85 90 95  
Thr Val Glu Gly  
100

<210> 27  
<211> 174  
<212> PRT  
<213> Human

<400> 27  
Pro Ser Gly Glu Glu Arg Lys Arg Pro Ala Pro Pro Arg Pro Ala Thr  
1 5 10 15  
Phe Pro Thr Arg Gln Pro Gly Leu Gly Ser Gln Asp Val Val Ser Lys  
20 25 30  
Ala Ala Asn Arg Arg Ile Pro Met Glu Gly Gln Arg Asp Ser Ala Phe  
35 40 45  
Pro Lys Phe Glu Ser Lys Pro Gln Ser Gln Glu Val Lys Glu Asn Gln  
50 55 60  
Thr Val Lys Phe Arg Cys Glu Val Ser Gly Ile Pro Lys Pro Glu Val  
65 70 75 80  
Ala Trp Phe Leu Glu Gly Thr Pro Val Arg Arg Gln Glu Gly Ser Ile  
85 90 95  
Glu Val Tyr Glu Asp Ala Gly Ser His Tyr Leu Cys Leu Leu Lys Ala  
100 105 110  
Arg Thr Arg Asp Ser Gly Thr Tyr Ser Cys Thr Ala Ser Asn Ala Gln  
115 120 125  
Gly Gln Val Ser Cys Ser Trp Thr Leu Gln Val Glu Arg Leu Ala Val  
130 135 140  
Met Glu Val Ala Pro Ser Phe Ser Ser Val Leu Lys Asp Cys Ala Val  
145 150 155 160  
Ile Glu Gly Gln Asp Phe Val Leu Gln Cys Ser Val Arg Gly  
165 170

<210> 28

<211> 97  
 <212> PRT  
 <213> Human

<400> 28  
 Pro Ala Phe Lys Gln Lys Leu Gln Asp Val His Val Ala Glu Gly Lys  
 1 5 10 15  
 Lys Leu Leu Leu Gln Cys Gln Val Ser Ser Asp Pro Pro Ala Thr Ile  
 20 25 30  
 Ile Trp Thr Leu Asn Gly Lys Thr Leu Lys Thr Thr Lys Phe Ile Ile  
 35 40 45  
 Leu Ser Gln Glu Gly Ser Leu Cys Ser Val Ser Ile Glu Lys Ala Leu  
 50 55 60  
 Leu Glu Asp Arg Gly Leu Tyr Lys Cys Val Ala Lys Asn Asp Ala Gly  
 65 70 75 80  
 Gln Ala Glu Cys Ser Cys Gln Val Thr Val Asp Asp Ala Pro Ala Ser  
 85 90 95  
 Glu

<210> 29  
 <211> 124  
 <212> PRT  
 <213> Human

<400> 29  
 Glu Ser Gln Gly Thr Ala Pro Ala Phe Lys Gln Lys Leu Gln Asp Val  
 1 5 10 15  
 His Val Ala Glu Gly Lys Lys Leu Leu Gln Cys Gln Val Ser Ser  
 20 25 30  
 Asp Pro Pro Ala Thr Ile Ile Trp Thr Leu Asn Gly Lys Thr Leu Lys  
 35 40 45  
 Thr Thr Lys Phe Ile Ile Leu Ser Gln Glu Gly Ser Leu Cys Ser Val  
 50 55 60  
 Ser Ile Glu Lys Ala Leu Leu Glu Asp Arg Gly Leu Tyr Lys Cys Val  
 65 70 75 80  
 Ala Lys Asn Asp Ala Gly Gln Ala Glu Cys Ser Cys Gln Val Thr Val  
 85 90 95  
 Asp Asp Ala Pro Ala Ser Glu Asn Thr Lys Ala Pro Glu Met Lys Ser  
 100 105 110  
 Arg Arg Pro Lys Ser Ser Leu Pro Pro Val Leu Gly  
 115 120

<210> 30  
 <211> 87  
 <212> PRT  
 <213> Human

<400> 30  
 Ala Pro Ala Phe Ile Leu Pro Pro Arg Asn Leu Cys Ile Lys Glu Gly  
 1 5 10 15  
 Ala Thr Ala Lys Phe Glu Gly Arg Val Arg Gly Tyr Pro Glu Pro Gln  
 20 25 30  
 Val Thr Trp His Arg Asn Gly Gln Pro Ile Thr Ser Gly Gly Arg Phe  
 35 40 45  
 Leu Leu Asp Cys Gly Ile Arg Gly Thr Phe Ser Leu Val Ile His Ala  
 50 55 60  
 Val His Glu Glu Asp Arg Gly Lys Tyr Thr Cys Glu Ala Thr Asn Gly  
 65 70 75 80

Ser Gly Ala Arg Gln Val Thr  
85

<210> 31  
<211> 119  
<212> PRT  
<213> Human

<400> 31  
Ser Asn Ala Gln Gly Gln Val Ser Cys Ser Trp Thr Leu Gln Val Glu  
1 5 10 15  
Arg Leu Ala Val Met Glu Val Ala Pro Ser Phe Ser Ser Val Leu Lys  
20 25 30  
Asp Cys Ala Val Ile Glu Gly Gln Asp Phe Val Leu Gln Cys Ser Val  
35 40 45  
Arg Gly Thr Pro Val Pro Arg Ile Thr Trp Leu Leu Asn Gly Gln Pro  
50 55 60  
Ile Gln Tyr Ala Arg Ser Thr Cys Glu Ala Gly Val Ala Glu Leu His  
65 70 75 80  
Ile Gln Asp Ala Leu Pro Glu Asp His Gly Thr Tyr Thr Cys Leu Ala  
85 90 95  
Glu Asn Ala Leu Gly Gln Val Ser Cys Ser Ala Trp Val Thr Val His  
100 105 110  
Glu Lys Lys Ser Ser Arg Lys  
115

<210> 32  
<211> 98  
<212> PRT  
<213> Human

<400> 32  
Gly Gln Arg Asp Ser Ala Phe Pro Lys Phe Glu Ser Lys Pro Gln Ser  
1 5 10 15  
Gln Glu Val Lys Glu Asn Gln Thr Val Lys Phe Arg Cys Glu Val Ser  
20 25 30  
Gly Ile Pro Lys Pro Glu Val Ala Trp Phe Leu Glu Gly Thr Pro Val  
35 40 45  
Arg Arg Gln Glu Gly Ser Ile Glu Val Tyr Glu Asp Ala Gly Ser His  
50 55 60  
Tyr Leu Cys Leu Leu Lys Ala Arg Thr Arg Asp Ser Gly Thr Tyr Ser  
65 70 75 80  
Cys Thr Ala Ser Asn Ala Gln Gly Gln Val Ser Cys Ser Trp Thr Leu  
85 90 95  
Gln Val

<210> 33  
<211> 82  
<212> PRT  
<213> Human

<400> 33  
Val Thr Ala Ser Leu Gly Gln Ser Val Leu Ile Ser Cys Ala Ile Ala  
1 5 10 15  
Gly Asp Pro Phe Pro Thr Val His Trp Leu Arg Asp Gly Lys Ala Leu  
20 25 30  
Cys Lys Asp Thr Gly His Phe Glu Val Leu Gln Asn Glu Asp Val Phe



(12) INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(19) World Intellectual Property Organization  
International Bureau



(43) International Publication Date  
23 May 2002 (23.05.2002)

PCT

(10) International Publication Number  
**WO 02/040683 A3**

(51) International Patent Classification<sup>7</sup>: **C12N 15/54**,  
9/12, G01N 33/573, C12Q 1/68, A61K 38/45

(21) International Application Number: PCT/US01/32616

(22) International Filing Date: 22 October 2001 (22.10.2001)

(25) Filing Language: English

(26) Publication Language: English

(30) Priority Data:  
09/711,134 14 November 2000 (14.11.2000) US  
09/858,664 17 May 2001 (17.05.2001) US

(71) Applicant: **PE CORPORATION (NY)** [US/US]; 761  
Main Avenue, Norwalk, CT 06859 (US).

(72) Inventors: **WEI, Ming-Hui**; c/o Celera Genomics, 45  
West Gude Drive C2-4#21, Rockville, MD 20850 (US).  
**KETCHUM, Karen**; c/o Celera Genomics, 45 West  
Gude Drive C2-4#21, Rockville, MD 20850 (US). **DI**  
**FRANCESCO, Valentina**; c/o Celera Genomics, 45  
West Gude Drive C2-4#21, Rockville, MD 20850 (US).  
**BEASLEY, Ellen, M.**; c/o Celera Genomics, 45 West  
Gude Drive C2-4#21, Rockville, MD 20850 (US).

(74) Agent: **MILLMAN, Robert, A.**; Celera Genomics,  
Chief Intellectual Property Counsel, 45 West Gude Drive  
C2-4#21, Rockville, MD 20850 (US).

(81) Designated States (*national*): AE, AG, AL, AM, AT, AU,  
AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CO, CR, CU,  
CZ, DE, DK, DM, DZ, EC, EE, ES, FI, GB, GD, GE, GH,  
GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC,  
LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW,  
MX, MZ, NO, NZ, PH, PL, PT, RO, RU, SD, SE, SG, SI,  
SK, SL, TJ, TM, TR, TT, TZ, UA, UG, UZ, VN, YU, ZA,  
ZW.

(84) Designated States (*regional*): ARIPO patent (GH, GM,  
KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZW), Eurasian  
patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European  
patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE,  
IT, LU, MC, NL, PT, SE, TR), OAPI patent (BF, BJ, CF,  
CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD,  
TG).

Published:

— with international search report

(88) Date of publication of the international search report:  
17 July 2003

*For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.*

(54) Title: ISOLATED HUMAN KINASE PROTEINS, NUCLEIC ACID MOLECULES ENCODING HUMAN KINASE PROTEINS, AND USES THEREOF

(57) Abstract: The present invention provides amino acid sequences of peptides that are encoded by genes within the human genome, the kinase peptides of the present invention. The present invention specifically provides isolated peptide and nucleic acid molecules, methods of identifying orthologs and paralogs of the transporter peptides, and methods of identifying modulators of the kinase peptides.

WO 02/040683 A3

## INTERNATIONAL SEARCH REPORT

onal Application No

PCT/US 01/32616

## A. CLASSIFICATION OF SUBJECT MATTER

IPC 7 C12N15/54 C12N9/12 G01N33/573 C12Q1/68 A61K38/45

According to International Patent Classification (IPC) or to both national classification and IPC

## B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

IPC 7 C12N C12Q A61K G01N

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

EPO-Internal, WPI Data, PAJ, SEQUENCE SEARCH, BIOSIS

## C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	EP 0 911 408 A (JAPAN SCIENCE & TECH CORP) 28 April 1999 (1999-04-28) the whole document	1-16, 19-23
X	GALLAGHER P J ET AL: "MOLECULAR CHARACTERIZATION OF A MAMMALIAN SMOOTH MUSCLE MYOSIN LIGHT CHAIN KINASE" JOURNAL OF BIOLOGICAL CHEMISTRY, AMERICAN SOCIETY OF BIOLOGICAL CHEMISTS, BALTIMORE, MD, US, vol. 266, no. 35, 15 December 1991 (1991-12-15), pages 23936-23944, XP000979557 ISSN: 0021-9258 the whole document	1-16, 19-23

☐ Further documents are listed in the continuation of box C.

Patent family members are listed in annex.

## \* Special categories of cited documents:

\*A\* document defining the general state of the art which is not considered to be of particular relevance

\*E\* earlier document but published on or after the international filing date

\*L\* document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)

\*O\* document referring to an oral disclosure, use, exhibition or other means

\*P\* document published prior to the international filing date but later than the priority date claimed

\*T\* later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention

\*X\* document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone

\*Y\* document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.

\*G\* document member of the same patent family

Date of the actual completion of the international search

12 November 2002

Date of mailing of the international search report

19/11/2002

Name and mailing address of the ISA

European Patent Office, P.B. 5818 Patentlaan 2  
NL - 2280 HV Rijswijk  
Tel. (+31-70) 340-2040, Tx. 31 651 epo nl,  
Fax: (+31-70) 340-3016

Authorized officer

Keller, Y

## INTERNATIONAL SEARCH REPORT

International application No.  
PCT/US 01/32616**Box I Observations where certain claims were found unsearchable (Continuation of item 1 of first sheet)**

This International Search Report has not been established in respect of certain claims under Article 17(2)(a) for the following reasons:

1. ☐ Claims Nos.:  
because they relate to subject matter not required to be searched by this Authority, namely:
  
2. ☒ Claims Nos.: 17, 18  
because they relate to parts of the International Application that do not comply with the prescribed requirements to such an extent that no meaningful International Search can be carried out, specifically:  
see FURTHER INFORMATION sheet PCT/ISA/210
  
3. ☐ Claims Nos.:  
because they are dependent claims and are not drafted in accordance with the second and third sentences of Rule 6.4(a).

**Box II Observations where unity of invention is lacking (Continuation of item 2 of first sheet)**

This International Searching Authority found multiple inventions in this International application, as follows:

1. ☐ As all required additional search fees were timely paid by the applicant, this International Search Report covers all searchable claims.
2. ☐ As all searchable claims could be searched without effort justifying an additional fee, this Authority did not invite payment of any additional fee.
3. ☐ As only some of the required additional search fees were timely paid by the applicant, this International Search Report covers only those claims for which fees were paid, specifically claims Nos.:
4. ☐ No required additional search fees were timely paid by the applicant. Consequently, this International Search Report is restricted to the invention first mentioned in the claims; it is covered by claims Nos.:

**Remark on Protest**

- ☐ The additional search fees were accompanied by the applicant's protest.
- ☐ No protest accompanied the payment of additional search fees.

## FURTHER INFORMATION CONTINUED FROM PCT/ISA/ 210

Continuation of Box I.2

Claims Nos.: 17, 18

Present claims 17, 18 relate to an extremely large number of possible compounds. Support within the meaning of Article 6 PCT and/or disclosure within the meaning of Article 5 PCT is to be found, however, for only a very small proportion of the compounds claimed. In the present case, the claims so lack support, and the application so lacks disclosure, that a meaningful search over the whole of the claimed scope is impossible.

The applicant's attention is drawn to the fact that claims, or parts of claims, relating to inventions in respect of which no international search report has been established need not be the subject of an international preliminary examination (Rule 66.1(e) PCT). The applicant is advised that the EPO policy when acting as an International Preliminary Examining Authority is normally not to carry out a preliminary examination on matter which has not been searched. This is the case irrespective of whether or not the claims are amended following receipt of the search report or during any Chapter II procedure.



# INTERNATIONAL SEARCH REPORT

Information on patent family members

International Application No

PCT/US 01/32616

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
EP 0911408 A	28-04-1999	JP 11098984 A	13-04-1999
		CA 2244928 A1	26-03-1999
		EP 0911408 A2	28-04-1999
		US 5958748 A	28-09-1999
		US 6171841 B1	09-01-2001